in the physician's opinion and the radiologist's report. One of these plates shows fluid as was diagnosed, but it also indicates the presence of a new growth which was not diagnosed. Whenever pleural effusion exists, an opportunity should be taken of examining the lungs by X rays, either when the fluid has been absorbed or soon after aspiration. We may then look for signs of tubercle or of new growth.

[A series of plates were given which showed a new growth demonstrated in this way.]

Bronchiectasis.—This condition is generally easily recognisable in X ray photographs, but occasionally the details are obscured and of doubtful interpretation. In such cases recourse should be made to the injection of lipiodol into the trachea; this sinks into and delineates the dilated bronchioles in a very striking fashion. This substance likewise often renders valuable assistance in defining the course and origin of sinuses in the chest wall.

TECHNIQUE IN RADIOGRAPHY.

I think I have said enough to demonstrate the exceedingly valuable service X ray examination of the lungs affords. The use made of this method of diagnosis in this hospital is very considerable; in fact, on an average, some 50 to 60 per cent. of the in-patients are X rayed at some time during their residence. The proportion would be higher with increased facilities, and we have under consideration a system whereby every case shall be X rayed on admission. I consider this routine examination very desirable, simply as an aid to clinical diagnosis.

To make this routine examination effective, however, a high standard of technique must be maintained. At present many X ray pictures are good, but too many are bad. The number of requests for repeats have been too large. These should be, I am certain, unnecessary, and I consider it should be as possible to produce uniformly good X ray photographs as it is to produce technically perfect photographic studio portraits—it is largely a matter of standardising all procedures. It is good technique which we physicians look for before anything else. There is a natural disposition to make our own interpretations, and an expert interpretation from a bad plate does not satisfy us.

There is another point of technique which, in my opinion, requires urgent attention—that is, the accurate standardisation of position. The point of view should always be the same, or only purposely varied—we then know what to look for. This is especially important when investigating the chest, for it is frequently desirable to note alterations in the size or position of various structures. Now, unless the point of view is exactly the same in each case, judgment becomes impossible. The incidence of divergent rays upon objects in different positions and planes produces a distortion in relative size and position so that it is impossible to compare photographs in which the position of the source

of the rays has varied even to a small degree. These remarks apply with special force to serial photographs.

Further, the source of the rays should be as far removed from the object as possible—the further away the more the rays approach to parallelism and the exaggeration of the shadows of the nearer objects is reduced; a more true picture is thus obtained.

Technical points such as these are of great importance in our special work in this hospital. They seem to be at present insufficiently appreciated, but their careful consideration, which I hope we shall achieve in time, will double the value of X ray work to us physicians.

"PROSTATIC" NEURASTHENIA.

BY

J. JOHNSTON ABRAHAM, C.B.E., D.S.O., M.D.,
F.R.C.S. ENG.,

SURGEON, LONDON LOCK HOSPITALS; SENIOR SURGEON,
KENSINGTON GENERAL HOSPITAL.

If there is one sort of patient more than another who is pre-eminently a source of infinite weariness to everyone with whom he comes in contact, it is probably that unfortunate individual known as the "prostatic" neurasthenic.

He is a constant worry to himself, to his family doctor, to the series of consultants—physicians, neurologists, urologists, electro-therapeutists, and probably, in a last despairing fling, to the psychoanalysts—to whom he has been sent in the hope that some one or other of them may light on a cure. Nobody loves him. Indeed, the only person who really appears to hail his advent with delight is the quack with his pills for "lost vitality," to whom he turns eventually in despair when orthodox medicine has failed him. A short account of how he may be treated successfully may therefore be of interest.

His history is generally as follows: At some unfortunate time he acquires an attack of urethritis, sometimes gonococcal, more often due to secondary organisms such as staphylococci or coliform bacilli. For this attack he is treated more or less adequately; he never develops any definite complications; and in the efflux of time his urine clears.

But in the interval he has suffered a mental trauma. He is naturally what is called a nervous man. He is generally very intelligent. He takes a keen interest in his symptoms. He reads up the subject, examines his urine for flakes, looks for signs of recurrence, and worries continually.

Occasional attacks of phosphaturia and stinging at the penile tip after micturition occur, and keep up his fear. He develops neuralgia along the cord

and in the testes. If he is a married man—and he frequently is—he fears to infect his partner, although, as the result of a series of tests, he has been assured that he is free from infection.

After this, in course of time, he finds that his virility is diminishing. He gets incomplete erections, has premature emissions, notices that the force in micturition becomes less, and occasionally he finds the stream of urine is forked. There is a constant dampness at the meatus, and the urethra does not empty itself completely in urination—a few drops dribble out afterwards.

He now complains of a constant dull ache in the perineum; and, after an action of the bowels, if the motion is at all hard, he may have what he calls a "loss" from the urethra—in other words, prostaticorrhœa. By this time he is thin and anxious-looking, pasty-complexioned, flabby-muscled. He sleeps badly, and is constantly thinking about his condition.

When he comes to see you, if you happen to shake hands with him, you will notice that his fingers are curiously clammy. On examination, the meatus will be found to be wet and rather purple. In hospital cases, as a rule, smegma is abundant. The testes will be found smaller, softer, and more tender than normal. The urine may be brilliantly clear and acid, with a few flakes, or amphoteric, or alkaline with phosphatic turbidity, or faintly opalescent from fine epithelial debris or mucus. Examination of smears from the meatus, the anterior urethra, the prostate, and the seminal vesicles, after massage, will be negative for organisms, and cultures of the urine following such massage are also sterile. The prostate itself, felt per rectum, may be larger and fuller than usual, but it is always smooth and compressible, and without vesicular thickening. Occasionally both lobes may be flattened and empty to the touch, but the urethra will then stand out as a circular cord, rather firm and thickened. On endoscopic examination of the penile urethra the mucous membrane may show its Littre's glands very prominently—a sign that mucorrhœa (the analogue of leucorrhœa) is probably present. There is no evidence of stricture on passing a sound, but more than usual discomfort is complained of when the instrument reaches the posterior urethra. Here one meets with a distinct feeling of obstruction, and, unless one has delicate hands, a little bleeding may be produced.

On passing the posterior urethroscope practically nothing can be seen except a red transversely lobulated mass—the verumontanum—which seems to fill the entire prostatic urethra, and prevents any sight of the lateral recesses. It is usually covered with mucus, is very friable, and may have a few small polypoid masses projecting into the lumen of the field of vision. Occasionally the sinus pularis is seen, and then generally it exudes a mucoid discharge. The clinical picture is that of a much hypertrophied verumontanum.

Everything else having been excluded already by the physical and bacteriological examination, and nothing abnormal having been found except this very much enlarged verumontanum, one now comes to the conclusion that this, in all probability, is the underlying cause of the condition.

FUNCTIONAL DERANGEMENT.

All neurasthenics have some derangement of function which, if discovered and treated, will ultimately lead to cure. Neurasthenia is probably only a name to cover our ignorance of a diversity of conditions produced by want of balance of the sympathetic nervous system caused by organisms of low toxic virulence or by traumata which would not cause pain in normal individuals, and would not be noticed by them, but none the less capable of producing symptoms in people with an unstable sympathetic nervous mechanism.

In prostatic neurasthenia the trouble seems to be produced by an error of function of the verumontanum. What use in the metabolism of the body is subserved by this organ is not known. Histologically it consists of a central glandular mass and a peripheral fibrous portion covered by squamous epithelium. The central part is made up of racemose glands lined with cubical or columnar epithelium, each lobule surrounded by well-marked elastic fibres. The glands apparently have no ducts; but, as noted, they have a very definite fibrous elastic capsule. They also have a markedly good vascular and lymphatic supply.

Why the verumontanum should hypertrophy in sexual neurasthenia is not known, but that the neurosis clears up when the hypertrophy is reduced should be recognised; and it is therefore on the lines of depletion that treatment is carried out.

GENERAL AND LOCAL TREATMENT.

General.—Having excluded all focal disease in the vesicles, prostate, Cowper's glands, and anterior urethra, we may now proceed to attack the condition. But to do so successfully it is absolutely essential to gain the patient's confidence. His story must be listened to with patience, his symptoms inquired into, his urine looked at regularly. Recognising that there is a large element of neurosis in the state, sedatives must be prescribed—bromides, valerian, belladonna. These serve a double function: they act systemically by dulling the general excitability; they also act locally by deadening the stimulus on the erection centre produced by the engorged verumontanum, and they quiet down the perineal and testicular neuralgia. Salines also are indicated, and to the saline prescribed 5–15 minims of ext. ergotæ liq. appears to be a useful adjuvant. Salines are used because it is essential to avoid the risk of a hard motion pressing on the prostate in defæcation and producing prostaticorrhœa. Such a symptom is exceedingly distressing to the patient's mind. He

is sure he loses "vitality" thereby. For similar reasons highly spiced food should be avoided, and to the list of things forbidden should be added coffee, especially at night.

Local.—Prostatic massage alone is useless, but, if a large cold sound is introduced into the bladder, and the posterior urethra massaged against this, improvement may be expected. Diathermy, with a metal electrode in the rectum, and a wide lead band round the pelvis, gives possibly better results—the element of suggestion is much more powerful with diathermy. But what produces the most rapid effect is silver nitrate painted or instilled on the verumontanum directly. This can be done either by an Ultzmann's catheter or a Guyon's syringe, starting with 15 minims of a $\frac{1}{2}$ per cent. solution and increasing weekly by $\frac{1}{2}$ per cent. until 5 per cent. is used. Or, instead, a posterior urethroscope can be passed and the "caput" painted directly with 5 per cent. silver nitrate once a week.

All the while treatment is being carried out cohabitation is contraindicated. The patient as a rule has a greatly diminished sexual desire, owing to the dulling of the trigger area in the posterior urethra caused by the treatment. But as he feels himself improving he may be tempted, and when he finds that he is impotent, as he will be owing to this treatment, he may become extremely depressed mentally by the fact thus suddenly forcing itself upon him. It is wise, therefore, to inform him of this temporary result of the treatment beforehand. Topical treatment ceases when the caput has diminished to such a size as to allow the lateral recesses to be seen easily. Six weeks' treatment with strychnine tonics will then restore sexual power again. The sense of fullness and pressure in the perineum goes when congestion of the verumontanum has been cured. The neuros-thenic symptoms built upon the congestion likewise diminish, and in the course of a few months the patient regains his normal outlook on life.

DISABILITIES OF THE HIP-JOINT IN CHILDREN.*

BY

ARTHUR ROCYN JONES, M.B., B.S. LOND.,
F.R.C.S. ENG.,

SURGEON, ROYAL NATIONAL ORTHOPÆDIC HOSPITAL;
ORTHOPÆDIC SURGEON, CHILDREN'S CLINIC.

DISABILITIES of the hip-joint may be conveniently divided into two groups: congenital and acquired. Of the former congenital dislocation and coxa vara are probably the commonest, and their similarity in symptoms and signs often present a difficulty in diagnosis.

CONGENITAL DISLOCATION OF THE HIP.

This rarely comes under observation until the child begins to walk, when lameness attracts attention; girls are more often affected than boys, the proportion being nine to one in a series of cases extending over 20 years at the Royal National Orthopædic Hospital. In a bilateral case the lumbar lordosis is greatly increased, and the gait is accompanied by a rolling of the body from side to side. In the unilateral case these characteristics are less marked, the repeated tilting of the body being confined to one side. Movement at the hip is perfectly free and painless, but abduction is limited by the taut adductors. The head of the femur is absent from Scarpa's triangle, but can be felt on the dorsum ilii with the top of the trochanter above Nélaton's line. If the limb is pulled downwards

and the hand then relaxes, the head of the femur telescopes into the buttock.

Another characteristic, also having its origin in the instability of the joint, is Trendelenberg's sign. If a child stands on one limb the other side of the pelvis is raised, if its corresponding thigh is flexed; but if the hip of the supporting limb is dislocated, the opposite side of the pelvis drops. The presence of all these signs should leave no doubt about the diagnosis. The common error is to mistake the condition for coxa vara and less frequently for infantile paralysis. But the fact that there is no wasting, muscular weakness, or circulatory disturbance present should be enough to exclude palsy. Coxa vara, however, has a distinct resemblance, and it will be well to consider this disability next.

COXA VARA.

Coxa vara is strictly a structural term, and denotes a diminution of the normal angle between the neck and shaft of the femur. It may be congenital or acquired, and unilateral or bilateral. The acquired form is often due to trauma causing either a separation or slipping of the epiphysis or a fracture of the neck with malunion. It is frequently associated with rickets, osteomalacia, and osteomyelitis. In these diseases the femoral neck is softened, and the body-weight is the mechanical factor altering the bony angle. The commonest symptom is lameness dating from birth or following an accident. But this symptom may be gradual in onset and follow repeated minor accidents of apparently little consequence. The joint becomes gradually stiffer and on examination flexion, abduction and internal rotation are particularly limited, and if the disability follows a recent accident muscle spasm may be present. There is some shortening of the limb, and the top of the great trochanter is above Nélaton's line.

* Abstract of a Post-Graduate Lecture delivered in connexion with the Children's Clinic, Marylebone-road. (Cases illustrating the various disabilities with the end-results of treatment were shown.)