Lesion of the Median Nerve.

The patient had been given an injection of calcium chloride six weeks previously. By some mischance the vein above the elbow had been missed and there was a healing scar in this region. He complained of weakness and numbness of the right hand. When he clasped his hands the right index finger failed to flex (Oschner's clasping sign). When the base of the thumb was held he was unable to flex the terminal phalanx of the thumb. The area of decreased sensation was quite typical—the injection had resulted in a lesion of the median nerve above the elbow.

Cases presented by HAMILTON BAILEY, F.R.C.S.

Correspondence.

To the Editor of the "Post-Graduate Medical Journal."

SIR,—In my recent course of Lectures on Migraine delivered at the Medical Society of London, I drew attention to the feeling of well-being that some migraine patients have the day before an attack, and suggested that it was probably to be explained by a general rise in blood-pressure. I drew attention to the parallel case of the victims of blood-pressure who so often feel that they were never better in their life the day before a hæmorrhage and to the known fact that a certain standard of blood-pressure often makes a patient more comfortable.

Since I gave that course of lectures I have seen a number of cases of migraine and have concentrated, among other things, on this point. I have been astonished to find how many of the patients are quite definite that they feel very much better the day before an attack; the percentage of these people is really very large. This has led me to suggest that as soon as ever this symptom comes on they should take a tablet of calcium diuretin, gr. $7\frac{1}{2}$ (Knoll Ltd.) three times a day for a couple of days. I have already found in one case that this has aborted what a patient was quite confident was going to be a severe attack of migraine. I would venture to suggest that others should give this treatment a trial, and would be grateful for their record of any cases in which it is done. May I suggest that the tablets should not be swallowed whole, but should be dissolved in a quarter of a tumblerful of water and an equal quantity of water drunk afterwards. This takes away the objectionable taste and prevents gastric disturbance. I have known one

case in which calcium diuretin taken as a tablet caused a severe passing attack of gastritis. This cannot be debited to the drug, but simply to a fault in the method of administration.

May I at the same time raise another point in connection with migraine? I have been following up very carefully in my recent cases the question of heredity, and have been profoundly impressed by the fact that one rarely, if ever, fails to get a history of other blood relatives suffering from the same thing or from one of the other allied vasomotor troubles such as asthma, urticaria, &c. I think that this is a point well worth the careful attention of all who are interested in the condition.

Dr. Julia Bell has most kindly been helping me in this matter of heredity and would be interested to hear of any cases. Communications could be sent to her at the Galton Laboratory, University College, Gower Street.

The incidence factor in the population is, of course, a very important point in connection with heredity, though a very difficult one to elucidate. My own strong impression is that true migraine is a comparatively rare affection, although, for reasons I gave in my paper, its rarity is masked and a wholly erroneous impression of its frequency prevails. I met with it 300 times in 5,000 cases, but it would not be fair to suppose that this gives its relative frequency in the population inasmuch as they had all come to consult an eye specialist. If they had been taken truly at random the probability is that the percentage would have been much lower. This is one of the points on which Dr. Bell is particularly anxious for information.

Yours truly,
R. H. ELLIOT.