

ADVERSE DRUG REACTION

Complete dysphagia after thrombolytic treatment for myocardial infarction

T Mosimann, L Terracciano, C Sieber, F Rappo

Abstract

An 82 year old man was admitted to hospital with unstable angina pectoris. There was a long history of minor symptoms suggesting reflux disease, with a small diaphragmatic hernia. One day after admission the patient complained of severe chest pain. An acute inferior-posterior myocardial infarction was diagnosed on ECG, and thrombolytic treatment with alteplase (rt-PA) was initiated. Within a few hours total dysphagia occurred, caused by haemorrhagic oesophagitis. The haematoma resolved spontaneously within about 10 days. The patient was discharged three weeks later after full resolution of the dysphagia.

(Postgrad Med J 2000;76:795-796)

Keywords: haemorrhagic oesophagitis; thrombolytic treatment; bleeding; myocardial infarction

Although gastro-oesophageal reflux disease with subsequent oesophagitis is very common, and thrombolytic treatment for myocardial infarction is a routine treatment, there has been only one case report of haemorrhagic oesophagitis with complete dysphagia after thrombolysis.¹ Given the high incidence of gastro-oesophageal reflux, it is possible that lesser degrees of haemorrhagic oesophagitis may occur in this situation more often than has been recognised.

Case report

An 82 year old patient was admitted to hospital with a diagnosis of unstable angina pectoris. He was being treated with aspirin 100 mg daily and there was no history of other non-steroidal anti-inflammatory drugs. There was also a history of infrequent dyspeptic symptoms.

On the next day the patient complained of sudden severe chest pain. The tentative diagnosis of an acute inferior-posterior myocardial infarction was confirmed by ECG. Thrombolytic treatment with alteplase (rt-PA) was initiated. A few hours later, the patient developed odynophagia, severe chest pain radiating into the right shoulder, and complete dysphagia. On day 5, parenteral feeding had to be started.

On day 7, an oesophagogastroduodenoscopy was performed. At 18 cm from incisors, the lumen was nearly blocked by necrotic debris and bleeding tissue, and 30 cm from incisors the lumen was totally obstructed. These findings suggested an oesophageal carcinoma.




Figure 1 Endoscopy performed before discharge showing grade 4 oesophagitis with ulceration and fibrin clots.

The differential diagnosis included exfoliative inflammation. Over the next 10 days the dysphagia improved steadily. Eleven days after the onset of the symptoms the patient was able to swallow a soft diet. The endoscopic biopsies showed typical severe oesophagitis with no malignant or dysplastic cells. Endoscopy performed before discharge showed grade 4 oesophagitis (fig 1). Seventeen days after the thrombolytic treatment the patient could eat a normal diet. He patient was discharged in good general condition after three weeks.

Discussion

Thrombolysis in acute myocardial infarction is a well established treatment. The complication of severe haemorrhage is a rare but also much dreaded event, and occurs in around 1-4% of all the patients undergoing thrombolytic treatment.^{2,3} However, the total number of haemorrhagic complications (including small bleeds) may be as high as 30%.³ Around 6% of the bleeding complications occur in the gastrointestinal tract.³ The three monthly prevalence of subjectively relevant dyspeptic complaints in Switzerland in 1997 was about 25%.⁴ According to published reports, oesophagitis can be endoscopically confirmed in 48-79% of this group of patients.⁵ Overall, the prevalence of endoscopically proven oesophagitis is around 10% in Switzerland, and around 5% in the international literature.⁵ If the prevalence of oesophagitis in patients with acute coronary thrombosis is not different from that of the population as a whole, one would expect to find potentially haemorrhagic reflux oesophagitis in every 10th patient in Switzerland. However, the patient's case history does not usually help

Department of Internal Medicine, Thierstein Hospital, 4042 Breitenbach, Switzerland
T Mosimann
F Rappo

Department of Pathology, University Hospital Basle, 4031 Basle, Switzerland
L Terracciano

Department of Gastroenterology, University Hospital Basle
C Sieber

Correspondence to:
Dr T Mosimann,
Department of Internal Medicine, Kantonsspital Luzern, 6000 Luzern 16, Switzerland
thierry.mosimann@datacomm.ch

Submitted 26 October 1999
Accepted 10 February 2000

Summary points

- Haemorrhages in the gastrointestinal tract account for 6.4% of all bleeding complications of thrombolytic treatment.
- The prevalence of endoscopically proven oesophagitis is about 10% in Switzerland. In the international literature, this figure is around 5%.
- Haemorrhagic oesophagitis can lead to total dysphagia.
- The incidence of dysphagia caused by haemorrhagic oesophagitis following thrombolytic treatment is probably much higher than previously reported.

the physician to assess the individual risk of haemorrhage resulting from gastro-oesophageal reflux disease. It is generally accepted that the intensity of reflux symptoms does not correlate with the endoscopic findings. On the contrary, it has been found that endoscopically proven ulcerative oesophagitis tends to cause few or no complaints. Given the figures mentioned above, one would expect a higher incidence of haemorrhagic oesophagitis following thrombolytic treatment than appears to be the case. However, milder forms of oesophageal haemorrhage without obvious dysphagia have been reported quite often.⁵

Our patient had oesophagogastroduodenoscopy six and 10 days after myocardial infarction. Given the serious and sudden onset symptoms of odynophagia and total dysphagia, endoscopy was required in this case, and the patient was considered medically stable. However, this procedure should be performed only while monitoring vital signs, and endotracheal intubation and mechanically assisted ventilation should be available.⁵

To the best of our knowledge this is only the second case where haemorrhagic reflux oesophagitis and total dysphagia developed so dramatically within a few hours after initiation of thrombolytic treatment.¹

- 1 Jishi F, Sissons CE, Silverstone EJ, et al. Oesophageal dissection after thrombolytic treatment for myocardial infarction. *Thorax* 1992;47:835-6.
- 2 Berkowitz SD, Granger CB, Pieper KS, et al. Incidence and predictors of bleeding after contemporary thrombolytic therapy for myocardial infarction. *Circulation* 1997;95:2508-16.
- 3 McLeod DC, Coln WG, Thayer CF, et al. Pharmacoeconomics of bleeding events after use of r-alteplase or streptokinase in acute myocardial infarction. *Am J Pharmacother* 1993;27:956-62.
- 4 Giger M, Brignoli R. Prävalenz von Oberbauchschmerzen und deren Einfluss auf Lebensqualität und Konsum von medizinischen Leistungen. *Schweiz Med Wochenschr* 1998;128:874-9.
- 5 Cappell MS. The safety and clinical utility of oesophagogastroduodenoscopy for acute gastrointestinal bleeding after myocardial infarction: a six-year study of 42 endoscopies in 34 consecutive patients at two university teaching hospitals. *Am J Gastroenterol* 1993;88:344-50.



Complete dysphagia after thrombolytic treatment for myocardial infarction

T Mosimann, L Terracciano, C Sieber and F Rappo

Postgrad Med J 2000 76: 795-796
doi: 10.1136/pgmj.76.902.795

Updated information and services can be found at:
<http://pmj.bmj.com/content/76/902/795>

These include:

References

This article cites 5 articles, 2 of which you can access for free at:
<http://pmj.bmj.com/content/76/902/795#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Bird flu](#) (5)
[Calcium and bone](#) (69)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>