DENTISTRY IN MEDICINE

By F. W. E. WAGNER, M.A., Sc.D.

Fellow of the Royal Academy of Medicine of Ireland. Fellow of the Royal Society of Tropical Medicine

I. The Differential Diagnosis of Pain

Whenever and wherever the professional training of a dental practitioner is discussed there is usually general agreement on one point, that he should possess more knowledge of general medicine and surgery than he does in fact possess. There are extremists who hold that the dental practitioner should go through the full curriculum leading to a qualification in medicine. With these not all will agree any more than they will agree with the extremists at the other end who hold that a dental training should be in dentistry and nothing else. But somewhere between these two extremes is to be found a large body of opinion that more medical knowledge is desirable. Not the least important of the reasons for this view is that such a knowledge would enable us to co-operate more fully and to greater mutual advantage with our colleagues in the medical profession.

On the other hand, within the dental profession we sometimes find ourselves wishing that the curriculum of our medical friends included more about dental surgery than it does. The point is one which I have discussed with medical colleagues, and they have usually agreed with me that even some elementary practical knowledge of dentistry would be very helpful.

This series of articles is an attempt to give to medical practitioners a brief, practical outline which may prove helpful to them in dealing with patients who come to them with trouble of dental origin, perhaps in places where no dentist is available.

This first article is concerned with pain that is of dental origin.

There are two conditions which give rise to this pain; one is when the patient can definitely say that a particular tooth (or teeth) is aching. That is the condition known as odontalgia or, more simply, just toothache.

The other is when the pain can be connected with the sensory nerves rather than with the teeth. On paper those look to be quite simple, cut-and-dried definitions but, like so many of the things that look simple on paper, in practice they are sometimes very difficult to distinguish.

The medical practitioner will do well to provide himself with a few simple diagnostic instruments—a couple of mouth-mirrors, two or three dental probes, a couple of metal burnishers with rounded ends, a spatula and a small pocket torch. The uses of most of these are self-evident. The spatula is necessary to foil the irresistible desire which most patients evince to wrap the tongue round the very tooth which you wish to examine, the burnishers are useful for percussing teeth, the torch for lighting up the mouth and also for transillumination.

Let me say a few words about simple methods of diagnosis. It will be necessary to apply heat and cold to individual teeth. Heat can be applied by picking up a ball of hot gutta-percha on the end of a burnisher and laying it on the tooth; cold by directing a fine spray of ethyl chloride on to the tooth; percussion by tapping the tooth with the rounded end of a burnisher. It is never necessary to tap hard, that only inflicts unnecessary pain whereas a gentle tap will elicit the symptoms just as well as a violent blow.

The pain of toothache varies according to whether it is caused by disease of the pulp or disease of the periodontal membrane.

I. Disease of the pulp

Here we may have a variety of causes, such as inflammation arising from decay of the tooth or from a fracture involving the pulp or from suppuration or exposure of the pulp. One needs to be careful about acting on the statement of a patient that a particular tooth is the

one at fault. It may be the one in which he feels the pain, but the actual lesion causing the pain may be in some other tooth. Pain in an upper tooth may be due to a lesion in a lower and vice versa, but pain is never referred across the middle line of the mouth. Examine the tooth which the patient blames, but examine the other teeth on that side of the mouth as well.

The pain is sharp and sudden in onset. Later it dies down to a dull ache. In the early stages it is not affected by either heat or cold. In what we might call the intermediate state the pain is of a throbbing type and occurs in bouts; at this stage it is relieved by cold and intensified by heat. In the later stages the pain is intensified by both extreme heat and extreme cold. The patient will usually complain of severe attacks during the night, and these are caused by the warmth of bed. Unless there is associated periodontitis the tooth is not usually sensitive to pressure. If the pulp is actually exposed the pain is very severe.

When a patient complains of a violent, throbbing pain, which is greatly intensified by heat the probability is that it is caused by supuration of the pulp in an enclosed cavity.

It frequently happens that a patient complains of paroxysmal pain caused when the tooth is touched by either sugar or salt. If, on examination it is observed that there is considerable recession of the gums from the necks of the teeth, this is the cause. If you even touch the tooth lightly with a strand of cotton wool it will excite quite severe pain. Though treatment is outside the province of this article, I ought to mention the treatment for pain of this kind. Dry the tooth and paint it with the following:—

Gum Mastic (Powder)	3 i.
Zinc Chloride	gr. v.
Chloroform	3 ss.

Keep the tooth dry until the solution hardens. May I also suggest a palliative for toothache, which a medical practitioner will find very useful. Mix zinc oxide, carbolised resin and oil of cloves to a stiff paste, about the consistency of putty. Clean out the cavity in the tooth as well as possible and pack it with this paste. The paste will harden.

After this unwarranted digression I return to diagnosis! But this paroxysmal pain aroused

by salt and sugar may be due to another cause. It often indicates the presence of hidden cavities of considerable size, frequently approximal. The tooth may look healthy enough, but the crown may be no more than a mere shell.

II. Periodontal Disease

Here the pain is of a different type, and may range from discomfort to considerable severity. It is localized, dull and throbbing, and is not affected by either heat or cold. The gums are swollen and of a dusky hue. Inflammation causes swelling of the periodontal membrane and, in obedience to the law of physics that two bodies cannot occupy the same space at the same time, something must move. It is the tooth that moves, and it rises slightly in the socket. The patient will tell you that the affected tooth seems longer than the others; and this is a characteristic symptom of the disease. So is tenderness on biting or pressure of any kind. The disease is more common in the lower jaw than in the upper.

Percussion is always an aid to diagnosis, for a diseased tooth gives a distinctly duller note than a healthy one.

We turn now to neuralgic pain, that is to say pain which is connected with the sensory nerves rather than with any individual tooth. True trigeminal neuralgia is happily of rare occurrence. It is usually an ailment of middle age. It is nearly always unilateral and is characterized by the sudden onset of violent, shooting pain, caused by the most trivial things. The unhappy sufferer is a pitiable object, for he dreads washing or shaving or touching his face lest it should provoke an attack. But, as I say, the disease is rare, and in twenty-four years I have only seen one case of it.

Malposed teeth should always be regarded with suspicion. By their position they are difficult to cleanse; particles of food get wedged in places where the brush cannot reach them, and decay follows. A careful examination of such teeth should be made with a fine probe for cavities between teeth and below the gum margin. The lower wisdom teeth are a fruitful cause of neuralgic pain. These erupt at varying angles and in all sorts of positions, so that the removal of one of

them can be the most difficult operation in dental surgery, calling for all the skill the surgeon possesses. If there is pain in the ear, it is almost certain that the wisdom is to blame. Frequently this tooth is partly covered with a thick flap of tough gum tissue and particles of food get under this flap and set up inflammation. Or the tooth may be erupting crookedly. In all cases where a wisdom tooth is suspect it is advisable to have a radiogram. Where there is inflammation, swelling and pus the tooth should be extracted without delay under a general anaesthetic. I do not agree with a policy of temporising and waiting until the condition clears up. It is much more likely that it will go from bad to worse. When a surgeon is confronted with an acute appendix he does not usually advise the patient to wait till it clears up; he operates without delay. The reasoning which holds good in such a case is equally applicable to the acute wisdom tooth.

Now, again, in diagnosing neuralgic pain one must not overlook the possibility of a com-

pletely unerupted tooth. Some time ago I was consulted by an edentulous patient aged 62. She complained of persistent infra-orbital neuralgic pain. A radiogram disclosed the presence of a completely unerupted canine tooth lying transversely in the palate. I removed this tooth with local anaesthesia, and there was no further pain.

Teeth that have been filled should be examined carefully. If they are loose or tender to pressure or respond to heat or cold or percussion, the fillings need removal followed by adequate treatment. If an apparently sound tooth is dark in colour or if it is less translucent than the others the pulp is probably dead.

Neuralgia may also be caused by general systemic conditions such as anaemia, rheumatism, diabetes, influenza, etc.

The nose and the accessory sinuses should be explored, and one must not lose sight of the fact that the pain may be symptomatic of tumour of the brain involving the root of the fifth nerve.

WRIGHT'S PUBLICATIONS

8½ × 5½ in.

444 pp.

338 illustrations.

50s. net; postage 7d.

EYE SURGERY

By H. B. STALLARD, M.B.E., M.D. (Cantab.), F.R.C.S. (Eng.)

7¼ × 4¾ in.

136 pp.

Illustrated.

12s. 6d. net; postage 3d.

A SHORT HANDBOOK OF PRACTICAL ANAESTHETICS

By HÖEL PARRY-PRICE, M.R.C.S., L.R.C.P., D.A. (R.C.S.)

With a Foreword by Sir CECIL WAKELEY, K.B.E., C.B.

Eighth Edition. Reprint. Fully Revised. 7¼ × 4¾ in. 1,233 pp. 30s. net; postage 7d.

SYNOPSIS OF MEDICINE

By Sir HENRY TIDY, K.B.E., M.D., F.R.C.P.

Bristol: JOHN WRIGHT & SONS LTD.

London: Simpkin Marshall (1941) Ltd.