ANNOTATION

Hiatus Hernia

In recent years there have been many important new contributions to our knowledge of diaphragmatic hernia of the non-traumatic type, particularly with regard to its incidence, diagnosis and methods of treatment. Although herniation of the stomach or intestines through the diaphragm following trauma had been recognised by clinicians for over 300 years, it was not until 1921 that the first radiological diagnosis of oesophageal hiatus hernia was made at the Mayo Clinic. Since then the increase in frequency of the diagnosis has been phenomenal. Harrington (1938) stated that whereas in 1921 a radiological diagnosis of hiatus hernia was only made once, in 1937 no less than 99 cases were diagnosed radiologically. Brick (1949), in a study of 3,448 patients, reported that it was the second commonest pathological lesion found in the upper gastro-intestinal tract, duodenal ulcer taking first place. If the examination be done by a radiologist experienced in the special technique of diagnosis, there seems no doubt that hiatus hernia will be found to occur even more frequently than it is at present.

In the past there has been little attempt to classify the various types of hernia encountered beyond calling them traumatic or non-traumatic, congenital or acquired. It is now generally agreed that a hollow abdominal organ may herniate into the thorax through a defect in the diaphragm in any one of the three following positions :—

- (1) The oesophageal hiatus.
- (2) The pleuro-peritoneal canal, or foramen of Bochdalek.
- (3) The foramen of Morgagni.

or there may be congenital absence of the posterior part of the diaphragm.

The commonest abnormality is for the cardia or part of the fundus of the stomach to herniate through the oesophageal hiatus. It may cause no symptoms in the patient until past middle life and so remain undiscovered. P. R. Allison of Leeds in his report of 204 cases (1951) gives a vivid descrip-

tion of the symptoms complained of by a typical sufferer ; in addition he describes the morbid anatomy of the conditions and offers a convincing explanation of how the pathological changes which are observed, occur. The account given here of incidence, symptomatology, pathological changes and treatment is based largely on his paper.

Allison describes two types of hernia through the oesophageal hiatus ; the paraoesophageal (or rolling) and the sliding hernia. Analysis of his patients show that 170 had sliding hernias, 21 had paraoesophageal hernias (with the cardia in normal position) and 13 had a combination of the two types of abnormality. There was only one patient with incomplete descent of the stomach associated with a congenital short oesophagus.

In regard to age incidence, there is now general agreement that a diagnosis of hiatus hernia in adults is usually made over the age of fifty years. Belsey has pointed out that in the Frenchay Hospital series of 94 patients, 14 of them were children. Sex incidence varies considerably ; in Harrington's series males appear to predominate whereas of Allison's patients, the majority were female.

The symptoms complained of are generally epigastric distress, heartburn, palpitations, acid regurgitation and dysphagia but the symptoms differ in the two types of hernia. Allison has emphasised the importance of distinguishing between the two types, in view of the difference in prognosis and treatment. In the sliding type of hernia, the patient's symptoms are due to a reflux oesophagitis, which develops in the lower part of the gullet, owing to incompetence of the cardiac sphincter. Allison argues that this alteration in the physiology of the cardia is due to loss of the pinchcock action of the right crus of the diaphragm on the oesophagus ; thus, the lower oesophagus is exposed to the regurgitation of acid gastric juice, especially when the patient lies down.

In the paraoesophageal type of hernia, the anatomy is quite different. The sphincter action

of the diaphragm is not lost ; hence the cardia is not incompetent, and it follows that reflux oesophagitis, with its consequent train of symptoms does not occur. Patients with this type of hernia may complain of distension after a meal, shortness of breath, and may then develop an ulcer in the herniated portion of the stomach.

Other presenting features have been noted. The patient may complain of severe spasmodic sub-sternal pain, radiating to the neck, arm or teeth ; such pain may be indistinguishable from that of angina pectoris or coronary occlusion ; anaemia may also be the presenting symptom. Belsey (1951) stresses the importance of recognising that in infants and children anaemia is almost invariably present in the Type 1 (sliding) variety. The anaemia is usually of the hypochromic type, but it is not certain that it is always due to recognised blood loss. In the writer's experience, sudden severe haematemesis in the absence of preceding dyspepsia was the presenting symptom in two patients ; both patients were found to be very anaemic and on subsequent investigation both were found to have a paraoesophageal hernia. No evidence of gastric or duodenal ulceration could be demonstrated.

Diagnosis of oesophageal hiatus hernia rests on the radiological demonstration that part of the stomach and, perhaps, the cardia lies above the diaphragm in certain conditions. Whenever the condition is suspected on clinical grounds, radiological confirmation should be sought, using the technique first described by Soresi in 1919. In addition to taking films with the patient in the prone position, it is sometimes useful to screen him standing in the lateral position with the body bending forward as if to touch the toes. Endoscopic examination of the oesophagus and stomach should always be carried out ; oesophagoscopy, in order to ascertain the position and function of the cardia and to determine if oesophagitis or ulceration be present ; gastroscopy in order to exclude co-existing disease of the stomach such as ulceration or carcinoma. It is not unknown for a patient to have had his hernia repaired and to return within a few months suffering from carcinoma of the stomach or other intra abdominal organ. A full blood count is always advisable,

especially if surgical treatment is under consideration. Also, although the anaemia associated with hiatus hernia is usually of the hypochromic type, it is not always so. The writer recently had a female patient with hiatus hernia and severe anaemia, which, when investigated, proved to be classical Addisonian anaemia. Her anaemia responded readily to full doses of vitamin B. 12, but would obviously not have improved if treated with iron only. Belsey recommends that estimation of the plasma proteins should be done if surgery is contemplated ; he also considers occult blood examination of the faeces, and a fractional test meal (with radiological evidence that the tube is in the stomach) are useful investigations. Other diseases in which similar symptoms may occur include cholecystitis, appendicitis and, as already stated, coronary artery disease.

As the age group in which these diseases occur is approximately the same as in hiatus hernia, it must be expected that in some patients the two may co-exist. Because a man has an hiatus hernia, it does not necessarily follow that he may not subsequently get an infarct of his ventricle. Careful evaluation of his symptoms and signs, the mode of onset, and the appropriate investigations will lessen the chances of error.

At present there is no general agreement regarding treatment of oesophageal hiatus hernia. Some physicians prefer to treat their patients conservatively, by the use of antacids, avoidance of a late evening meal, a drink of water before going to bed and sleeping in a semi-upright position. It is argued that many elderly patients are not in a condition to stand a major operation. The above measures often succeed in the paraoesophageal type of hernia.

Allison recommends the complete reduction and repair of hernias of the sliding type before reflux oesophagitis and ulceration has rendered the hernia irreducible. He states that the aim of any surgical repair should be to restore the physiologically active cardia, thus preventing the reflux of digestive juices from the stomach into the oesophagus. He achieves this by two main steps ; first, he divides the overstretched phrenoesophageal ligament and peritoneum around the cardia

(Continued on page 368)

Cholangio-hepatitis usually arises secondarily to obstruction of the biliary tract, and a mild degree is almost constant in cases of biliary obstruction. Rare cases have been described in which no obstruction of the biliary tract has been found. The clinical picture is very variable, but presentation as pyrexia of unknown origin (Charcot's intermittent biliary fever) is described and recognized, and Hobson and Rice-Oxley (1950) in a clinical study of the condition describe two such cases.

In the present case it is possible that the stenosis of the ampulla of Vater had resulted from the passage of a single stone, which would account for the severe attack of epigastric pain when the patient was sent to hospital as a possible perforated ulcer. It is noteworthy, however, that he was not jaundiced at this time, and no evidence of

gall bladder disease or scarring was found in the bile duct at operation. It would, therefore, appear that the primary abnormality in this case was at the ampulla of Vater. Cases of functional biliary obstruction have been reported, and hypertrophy of the muscle surrounding the ampulla has been found in many such cases (Berg, 1923; Newman, 1933).

I wish to thank Dr. R. S. Bruce Pearson, under whom the patient was admitted, for his help and permission to publish this case, Mr. Geoffrey Parker for details of the operation, and Dr. Hall for the pathological report.

BIBLIOGRAPHY

- BERG, J. (1923), *Arch. f. klin. Chir.*, **126**, 329.
 HOBSON, F. G., and RICE-OXLEY, J. M. (1950), *Postgrad. med. J.*, **26**, 267.
 NEWMAN, C. (1933), *Lancet*, **1**, 785.

ANNOTATION—Continued from page 334

and resutures these to the under surface of the diaphragm; second, he restores the oesophageal hiatus by lightly suturing the fibres of the right crus behind the oesophagus. He does this operation via the thorax. Of 33 patients with sliding hernias so treated 30 had relief of symptoms with a normally placed stomach and no X-ray evidence of gastric reflux. Other writers have recommended temporary paralysis of the diaphragm by phrenic crush; some advise this procedure as a preliminary to surgical repair of the hernia. The method has the merit of simplicity but it does not appear to relieve the patient's symptoms for an appreciable time. By the time that oesophagitis

has caused shortening of the oesophagus and stenosis of the lumen resection of the diseased portion of the oesophagus and stomach, with oesophagojejunostomy may be necessary. Dilatation of the stricture has not proved successful in the treatment of dysphagia.

It is beyond the scope of this short annotation to discuss fully the various methods of surgical treatment and their indications. Until reports of treatment of many more patients are available, individual physicians and surgeons will probably continue to treat their patients according to their own experience, supported by a critical follow-up of each.

HAROLD DAVIS, M.A., M.B., M.R.C.P.

RUTHIN CASTLE, NORTH WALES

A Clinic for the diagnosis and treatment of Internal Diseases (except Mental or Infectious Diseases). The Clinic is provided with a staff of doctors, technicians and nurses.

The surroundings are beautiful. The climate is mild. There is central heating throughout. The annual rainfall is 30.5 inches, that is, less than the average for England.

The Fees are inclusive and vary according to the room occupied.

For particulars apply to THE SECRETARY, Ruthin Castle, North Wales.

Telegrams: Castle, Ruthin.

Telephone: Ruthin 66.