rib, lateral to the attachment of the arcuate ligament of the diaphragm. The dissection must, therefore, be below the diaphragm and behind the peritoneum. Dissection upwards and forwards in this position must lead to the subphrenic space. This method can also be used on the left side for the approach to a posterior subphrenic abscess on that side.

Anterior subphrenic abscesses are approached by means of subcostal incisions which will lead directly to the abscess cavity extra-peritoneally, as adhesions will have walled off the abscess from the free peritoneal cavity below.

The importance of carrying out drainage by an extra-serous approach is shown by Ochsner and Graves. They found that the mortality of subphrenic abscesses drained by an approach which did not contaminate any serous cavity was about 21 per cent., while in those drained trans-pleurally the mortality was 39 per cent. and in those drained trans-peritoneally it was 35 per cent.

Summary

1. The anatomy of the subphrenic region has been described.
2. The aetiology of subphrenic infection has been discussed; most cases follow intra-abdominal suppuration.
3. The diagnosis has been considered. The

condition will only be diagnosed early if the surgeon is continually on the look-out for its presence. There is no single method of diagnosis although the radiological findings are of great help.

4. The course of the disease and its main complications have been described. It is to be noted that about 70 per cent. of cases will resolve on their own without going on to suppuration.

5. The use of antibiotics in the treatment of the infection has been described, together with the methods of drainage of the abscesses. The importance of an extra-serous approach to the abscess is stressed and the method through the bed of the resected 12th rib is described.

Acknowledgments

My thanks are due to my colleagues on the Staff of the Norfolk and Norwich Hospital for permission to make use of the notes of cases treated by them and for so kindly allowing me to reproduce some of the X-ray films from their cases.

BIBLIOGRAPHY

- BAILEY, H. (1948), *J. Int. Coll. Surg.*, **11**, 377.
 FRIEDMAN, P. S. (1950), *Radiology*, **55**, 36.
 GROSS, H. T. (1948), *Ohio St. Med. J.*, **44**, 1005.
 OCHSNER, A., and GRAVES, A. M. (1933), *Ann. Surg.*, **98**, 96.
 SHEA, P. C., JUN., and HOLDEN, W. D. (1948), *Arch. Surg.*, **84**, 843.
 THOREK, P. (1947), *Surgery*, **21**, 739.

RESUSCITATION OF THE NEWBORN

By IAN DONALD, M.B.E., M.D., B.A., M.R.C.O.G.

Reader, Institute of Obstetrics and Gynaecology, University of London

At no time is a baby likely to be as near death as it is within the first 15 minutes of extra-uterine life. The first few breaths demand a well co-ordinated effort to produce the necessary pressures to open up the lung alveoli. Experimental attempts to inflate collapsed lung artificially usually require pressures of over 40 cm. of water at first. After the lung has once been expanded, lower pressures will succeed in keeping the alveoli open.

For a child to be equal to this effort it must be fit enough to react adequately to the stimuli which the birth process inflicts. The stimuli are of two sorts. Firstly, there are those from skin and joints which crowd in upon the baby's consciousness. Secondly, these are reinforced by the physiological anoxia of delivery. In animals the

foetus often falls on to its snout and this provides a very powerful stimulus to respiration.

The carotid sinus, too, plays an important part in respiration as a receptor organ for stimuli, both pressor or chemical, and it is known that breathing is markedly depressed following sinus denervation in experimental animals. An accumulation of carbon dioxide will thus stimulate the carotid sinus, but only where it is functionally capable of reacting. Previous anoxia may diminish the response, especially in cases of prematurity and cerebral compression.

Aetiology of Neonatal Asphyxia

In extreme prematurity, alveolar development may not be sufficiently far advanced to sustain

respiration. Also, in less extreme cases, the immature respiratory centre may be relatively insensitive, while the effect of raised intracranial pressure or the presence of drugs and anaesthetics is also more pronounced. Anoxia of sufficient degree depresses the respiratory centre in all cases, but particularly in that of the premature infant.

No respiratory centre can function unless an adequate circulation of oxygenated blood is able to reach it. In a case of foetal shock following a difficult delivery these very necessary conditions may not obtain. Lastly, and very commonly, there may be some mechanical factor such as blockage of the respiratory passages by the aspiration of foreign material.

In brief, failure to breathe is usually attributable to one of three types of cause: firstly, a depressed respiratory centre; secondly, a defective blood circulation to it; and, thirdly, blockage of the air passages. The seeds of neonatal asphyxia are often sown before delivery. The placenta may become separated in ante-partum haemorrhage; moreover, where the placenta is praevia, there is the additional risk of compression by the presenting part during labour and therefore a further reduction in effective placental blood flow. Pre-eclamptic toxæmia and, above all, eclampsia are common causes of foetal asphyxia, either due to diffuse placental infarction or to spasm of the blood vessels supplying the placental site which is now known to occur in these conditions. Prolonged labour, especially with the membranes ruptured, increases foetal anoxia and during uterine contractions in labour the amount of oxygen transfer across the placental site is greatly reduced. Less commonly, circulation in the umbilical cord may be hampered by accidents which mechanically compress it.

Any precipitate fall in maternal blood pressure will produce anoxia in the foetus before delivery and maternal asphyxia, for example, during induction of anaesthesia, will inevitably asphyxiate the baby. The signs of foetal distress during labour are usually those of foetal anoxia and they often presage a correspondingly greater degree of difficulty in resuscitation at birth.

The choice of analgesic drugs during labour is largely determined by the likely effect that they will have upon the baby's readiness to breathe. Morphia and its derivatives are notorious offenders in this respect and should not be used in any case in which delivery is likely to occur within the next four hours; in the case of premature delivery this period should be much longer. To the premature foetus morphia is nothing less than poison. The barbiturates to some extent are also liable to contribute to neonatal respiratory depres-

sion, and paraldehyde is still one of the safest drugs to use.

All the general anaesthetics carry the risk of respiratory depression in the child, but the risk is largely proportional to the depth of anaesthesia achieved. The administration of nitrous oxide is so often accompanied by a reduced intake of oxygen that it should not be used in established foetal distress, except in the hands of an anaesthetist giving pure oxygen at the same time.

Causes of Asphyxia in the Baby after Birth

The aspiration of mucus, meconium or blood should always be thought of first. The stomach is likely to provide another source of mucus, especially in cases delivered by Caesarean section. This is regurgitated and often inhaled and the practice of gastric aspiration of mucus is a sound one.

A raised intracranial pressure depresses the respiratory centre directly. This may be caused by intracranial bleeding or the venous engorgement which accompanies asphyxia. Cerebral oedema acts likewise.

Congenital abnormalities such as laryngeal atresia are rare and will not be considered further.

Venous engorgement is the inevitable and immediate effect of asphyxia, especially in the brain, although the process is generalized throughout the body. Capillary haemorrhages then occur as a result. In addition, arteriolar constriction further reduces circulation. The vasomotor centres, too, suffer from the effects of asphyxia and are readily depressed so that the clinical signs of shock appear. Another result of anoxia is the accumulation of cerebrospinal fluid in excess. This is even more likely to occur after Caesarean section and with premature infants. The resulting pressure from it is therefore likely to increase the child's respiratory embarrassment. Anoxia is thus followed by vasomotor and respiratory depression which, in their turn, cause further anoxia. A vicious circle is established and within a short time irreversible changes set in.

Clinical Aspects

The old division of asphyxia into cases of 'blue and white' is now outmoded. It would be more appropriate to refer to the latter as foetal shock which, in fact, it is. The difference in colour is due to the difference in the state of the peripheral circulation and blue coloration of the skin simply denotes that there is still a sound peripheral blood flow.

The earliest signs of asphyxia are a full and powerful heartbeat which is slower than usual, a mechanism which to some extent conserves oxygen. When peripheral circulation fails, the

colour of the child turns grey and muscle tone is lost, the anal sphincter becomes toneless, the tongue falls backwards and the jaw is relaxed. The vocal chords lie separated in the paralyzed position; this at least facilitates intubation. Any respiratory attempts which may be made are very feeble, shallow and punctuated occasionally by gasps.

As a rule the changes which appear, especially in the central nervous system, cannot be reversed after 10 to 15 minutes of foetal shock (white asphyxia) and the longer it takes to oxygenate the baby and restore its circulation the greater will be the damage. When intracranial haemorrhage is the cause of the child's reluctance to breathe, vasomotor tone is poor from the beginning and no livid or blue stage is evident. The outlook here is less favourable.

Treatment of Neonatal Asphyxia

Four conditions must be satisfied:

1. Patent air passages.
2. Availability of a suitable atmosphere.
3. Adequate respiratory movements to make use of this atmosphere.
4. A reasonably good blood circulation to revive the vital centres with oxygenated blood.

Airway Clearance

This is the first step. Whenever possible, and before the first gasp is taken, the mouth, pharynx and nostrils should be cleared of mucus, etc., by suction. This should take priority to the usual wiping of the eyelids. The traditional mucus catheter is an unsatisfactory instrument, because one has to blow down it to clear it. This is a septic procedure. In hospital practice it is possible to provide mechanically operated suckers, of which the simplest and most readily available is one operated from a water suction pump attached to a tap. It is not necessary for the suction pressure to exceed 2 or 3 in. of mercury. After dividing the cord, the child should be placed, turned slightly on one side, in a head downwards position in a warmed tray or cot with a slope of 20°. A tray in which the head flap can be lowered is useful, because intubation, should it become necessary, can then be carried out without unduly disturbing the child.

If no signs of spontaneous respiration appear within three minutes of birth the glottis should be cleared of mucus with the help of an infant's laryngoscope. This alone may suffice, but if the glottic reflex is absent the trachea should be intubated forthwith with a small, curved, soft rubber tube about 3 in. long. If the child is well enough to resent this manoeuvre there is no need for it, but, in the cases in which it is indicated,

the child is so toneless that no difficulty should be met in carrying it out gently and quickly.

After aspirating any mucus which may be present in the trachea, intermittent positive pressure of oxygen (25 cm. of water) may be applied through the tube by means of a manually operated bag or some such device. Although the alveoli are not likely to be inflated by such low pressures at least the bronchial tree is flooded with oxygen which is readily absorbed and should the child take a breath spontaneously it cannot fail to benefit immediately.

It is a pity that obstetricians are often unskilled in intubation of the newborn. In good hands the procedure is harmless and is often live-saving. Extreme degrees of head downwards tilt, or worse, dangling the child by its ankles for a prolonged period, only serve to discourage inspiration due to the weight of the liver upon the diaphragm and, furthermore, the cerebral veins become still further engorged.

Intermittent tongue traction has often been preached as a method of artificial respiration by evoking a reflex of doubtful anatomical authenticity. The main point in drawing the tongue forwards is to improve the pharyngeal airway. This objective can be achieved more simply and with less trauma by using a small-sized airway.

Another simple and much-used method of supplying oxygen to the baby is to pass a fine rubber catheter into the nasopharynx through one of the nostrils. It is safe but inefficient.

Above all, whatever measure is taken, gentleness of handling is essential. The child's inability to respond to the stimuli of its own birth is due to its shocked condition. Further stimulation by direct methods is therefore both harmful and unnecessary and the sooner the child can be left in peace in a warm cot and a warm oxygen atmosphere the better.

Atmosphere

The child's greatest need is for oxygen. While in a state of asphyxia it already has more than enough carbon dioxide and it is therefore useless to supply it with more. Once respiration has become established the concentration of oxygen should be reduced to a level at which the colour remains satisfactory; from now on the administration of up to 5 per cent. carbon dioxide in a mixture will not do any harm and may encourage alveolar expansion.

The prolonged use of pure oxygen is to be deplored and concentrations of more than 50 to 60 per cent. are unnecessary. In time the child should be weaned off these concentrations gradually over a few days and not simply removed suddenly to breathe ordinary air. It is now

believed that it is these sudden changes in the concentration of oxygen breathed which are mainly responsible for the development of retro-lental fibroplasia in premature infants.

Aspiration of Gastric Contents

In the normal process of vaginal delivery the pressures applied to the trunk of the baby help to express mucus and swallowed fluid from the stomach, but in the case of the baby delivered by Caesarean section this does not obtain and the quantity of fluid in the stomach may be as much as seven times as great. Regurgitation of this mucus either produces laryngeal spasm with resulting cyanotic attacks should the glottic reflex be present or, in its absence, inhalation of foreign matter may occur. This phenomenon may be one of the reasons why babies delivered by Caesarean section more frequently encounter respiratory difficulties and it is possible that anaesthetists in the past have been unjustly blamed. It is often worthwhile passing a small rubber catheter down the oesophagus to remove the gastric contents by suction, particularly in premature infants. The vigorous baby, of course, needs no such treatment and is perfectly capable of dealing with any regurgitated material.

Drug Stimulation in Resuscitation

The two drugs in common use are alpha-lobeline, which is said to lower the threshold of the respiratory centre, and Nikethamide. The effects of both are only transient and are often followed by further respiratory depression. Lobeline has to be given intravenously and can be milked up the umbilical vein. The dose is 1/20 gr. In the past its usefulness has been greatly over-rated.

Coramine or Nikethamide is drastic and may actually be harmful, but in the baby that is narcotised from morphia or some such drug it is very often useful. Intracardiac injection of adrenaline constitutes no more than a *coup de grâce* and it is most unlikely that any good will come of it.

Synthetic vitamin K preparations are often injected, but there can be no reason for thinking that they could in any way control intracranial bleeding which is either traumatic or asphyxial, and vitamin K has no rational place in the treatment of shock.

Artificial Respiration

The older and traditional methods of Sylvester and Bird have now been abandoned. For one thing they involve exposure and handling of the child, which are most undesirable. These methods relied upon the elastic recoil of the thoracic cage after compression, but since these babies are in

a shocked condition and therefore toneless no such recoil is likely to occur. For the same reason intermittent squeezing of the chest with the fingers is not only useless but mischievous.

Mouth-to-mouth insufflation looks as though it will take a long time to die out. Most of the air so insufflated, of course, finds its way into the stomach and there can be few more certain ways of introducing infection into the air passages of the newborn.

Intratracheal insufflation with intermittent pressures of oxygen of 25 to 35 cm. of water is at least efficient, but the margin of safety is narrow. Pressures below 30 cm. of water are unlikely to inflate the alveoli and higher pressures may rupture them and produce emphysema. Under this treatment the chest will often appear to expand, but this is more due to the distention of the bronchial tree than to alveolar expansion. It is doubtful whether the alveoli can be safely and permanently opened up by this method and only spontaneous inspiratory efforts are likely to achieve this. The benefit of endotracheal oxygen is not questioned, for oxygen is undoubtedly absorbed from the bronchial mucous membrane and at least an efficient airway is guaranteed.

Rocking Methods of Artificial Respiration

The advantage of Eve's method is that circulation is first improved by increasing venous return to the heart, particularly from the head. The respiratory centre is not likely to recover however much oxygen is blown into the baby's lungs until circulation is restored so that it can receive a supply of well-oxygenated blood. By using the weight of the liver and the abdominal viscera as the pumping mechanism at least the physiological state is maintained whereby the movement of inspiration is accompanied by a reduced intrathoracic pressure and therefore increased venous return.

The method also has the advantage that it is perfectly simple. The baby's body is held across the chest of the operator who sways from side to side through an arc of 40 to 70°, about ten times a minute. The disadvantage is that it is difficult to administer oxygen at the same time. The Roberts-Talley rocker is a foolproof piece of apparatus which operates on this principle (Fig. 1). It is driven by the same supply of oxygen as supplies the baby. In it the baby can be easily intubated should it be necessary and positive pressures of oxygen can be delivered through the endotracheal tube synchronously with the descent of the diaphragm. Warmth is supplied by hot-water cans underneath the tray and a few minutes of this mechanical rocking will often witness a

remarkable improvement in the condition of a shocked baby.

Phrenic Nerve Stimulator

This works on the lines of the Sarnoff apparatus and has been introduced by Cross. An electrical stimulus is applied on either or both sides of the neck over the point of stimulation of the phrenic nerve at about 20-second intervals. In babies it is often difficult to select the exact point of application of the electrodes and other muscles supplied by the upper part of the brachial plexus are often stimulated; however, their contraction aids rather than hinders chest expansion. To judge from adult experience this is not comfortable. Obviously this treatment should be used almost at once after birth before irreversible damage has been caused to the central nervous system by anoxia.

Positive Pressure Oxygen Air Lock

Bloxson's apparatus is now much in vogue in the United States. The whole child is subjected

to positive pressures, usually of oxygen, alternating between 1 and 3 lb. per sq. in. The pressures are rhythmically varied to simulate the intermittent pressures which the child undergoes during the process of natural delivery. It is to be noted, however, that there is no differential pressure supplied either to the air passages, the thoracic cage or the child as a whole and it is therefore not easy to see why such generalized pressure should increase foetal oxygenation. However, it is claimed such rhythmic pressures are beneficial, that oxygen absorption through the skin is increased and that the 'expansion of gases' promotes bronchial drainage. The use of the apparatus in some American centres has been accompanied by a substantial reduction in foetal mortality, but this reduction is more or less universal in most modern centres throughout the world, where a variety of up-to-date methods are used.

Gastric Administration of Oxygen

Oxygen in the newborn is very rapidly absorbed from the stomach and intestinal tract. This well-known phenomenon has been made use of in Scandinavia where Akerren and Furstenberg have devised a double catheter through which oxygen is bubbled into the stomach via one limb and the excess gas is allowed to escape through the other, thereby preventing over-distension. A simpler method is to trickle oxygen very gently into the stomach through a fine polythene tube whose small calibre prevents the risk of over-distension. Many of us in England have now had some experience of the method and can testify to its immense value. It does nothing, of course, to expand lungs, but will often keep a baby sufficiently oxygenated to give it time to recover from the shock of delivery. (Yllo 1935).

Prophylaxis

Considering the many hazards that occur in the process of being born and the loss of foetal life which is still unnecessarily high, it is only natural to enquire into the possibilities of prophylaxis against neonatal asphyxia. There is, however, no specific measure. Good ante-natal supervision should reduce the incidence of dystocia, should practically eliminate eclampsia and play at least some part in reducing the risks of prematurity. Foetal distress should be noted early and acted upon. The judicious use of drugs, timely episiotomy where indicated and the more skilful use of the obstetric forceps will all help to reduce these hazards.

Late Prognosis

How much permanent damage is caused by

FIG. 1.—The Roberts-Talley resuscitator.

asphyxia at birth is still much debated. While it is certain that prolonged anoxia can do nothing but harm to the central nervous system and is likely to be responsible for many of the grosser handicaps of later life, it is not yet possible to relate the child's intellectual capacity to the amount of oxygen deprivation which it may have suffered at birth. Schreiber, in analyzing 900 mental defectives, traced evidence of birth asphyxia in 70 per cent. This is a sobering thought. In resuscitating the newborn one is not only, for the time being, a trustee for the child's life but also for its intellectual future.

Summary

The following is a useful guide to the resuscitative procedures in modern use and the order in which they may be applied:

1. Clear the airways by suction.
2. Lay the child head downwards on a slope of about 20° in a warm cot or some such apparatus as the Roberts-Talley rocker.
3. Supply oxygen either by tent or by mask.
4. If the child is limp insert an airway to prevent the tongue from blocking the pharynx. In the absence of a suitable airway draw the tongue forwards with a tongue forceps.
5. If there is no improvement within two or three minutes start the rocker in motion.
6. If this does not produce any benefit within a further two minutes intubate under direct vision. When white asphyxia is pronounced from the moment of birth it is often wiser to intubate at once.
7. If intubation is not feasible, or in cases of gross prematurity which stand intubation rather badly, pass a fine polythene tube into the stomach and supply a fine trickle of oxygen.
8. Aspirate mucus from the stomach particularly in premature babies and in cases that are apnoeic after Caesarean section.
9. If all these measures are unavailing, and particularly in the narcotized infant, inject Nikethamide or Coramine 0.25 to 0.5 ml. intravenously.

More harm than good is done by injudicious over-treatment and a shocked baby's dwindling chances of spontaneous recovery may be thereby wrecked. Three guiding principles must always be observed, namely warmth, oxygen and minimal handling.

Acknowledgment

Medical Pneumatics of 28 Thayer Street, London, W.1, have provided the illustration of the Roberts-Talley resuscitator.

BIBLIOGRAPHY

- AKERREN, Y., and FURSTENBERG, N. (1950), *J. Obs. Gyn. B.E.*, 57, (5), 705.
- BARCROFT, J. (1946), 'Researches on Prenatal Life,' Blackwell, Oxford.
- BLOXSOM, A. (1950), *J. Pediatrics*, St. Louis, 37, (3), 311.
- CROSS, K. W., and ROBERTS, P. W. (1951), *Brit. med. J.*, May 12, 1043.
- EVE, F. C., and FORSYTH, N. C. (1948), *Ibid.*, ii, 554.
- FLAGG, P. J. (1951), *Amer. J. Obs. Gyn.*, 21, 537.
- GELLIS, S. S., WHITE, PRISCILLA, PFEFFER, W. (1949), *New Eng. J. Med.*, 240, (14), 533.
- GIBBERD, G. F., and BLAIKLEY, J. B. (1935), *Lancet*, 229, 138.
- POTTER, EDITH L., and ROSENBAUM, W. (1943), *Amer. J. Obs. Gyn.*, 45, 822.
- ROBERTS, HILDA (1949), *J. Obs. Gyn. B.E.*, 56, 961.
- SCHREIBER, F. (1938), *J.A.M.A.*, 3, 1263.
- WILSON, R. A., TOMMEY, M. A., and JOHNSON, K. S. (1937), *S.G.O.*, 65, 601.
- Yllo, A. (1935) *Act. Paediatr.* 17, Suppl., 1, 122.

MEDICAL PNEUMATICS LIMITED

Manufacturers of the Oxygen Rocking
Resuscitator, the Aberdeen Oxygen
Incubator, Oxygen Therapy Apparatus,
Anaesthetic Apparatus, etc.

Full details on request

28 THAYER STREET, LONDON, W.1

Telephone: Welbeck 3216