the slow introduction of the anaphylactic substance.

SUMMARY.

(1) In eczema there is a special susceptibility to external or to absorbed substances.

(2) That this susceptibility may be congenital or acquired.

(3) That the susceptibility is of a similar kind, but not of such frequent occurrence as the allergy present in cases of asthma, hay-fever and urticaria.

(4) That there is a good deal of clinical evidence which indicates that this susceptibility is apt to be developed in those whose metabolic and eliminative functions are defective.

(5) That the best treatment is :—

(a) The removal of the patient from his usual surroundings and particularly from his special poison.

(b) A change in his food, habits and occupation, and, finally,

(c) A course of detoxication and desensitization.

REFERENCE.

- (1) "The Blood-vessels of the Human Skin and their Responses," Sir Thomas Lewis.

BRONCHIECTASIS.

A POST-GRADUATE LECTURE DELIVERED AT YORK.

(Under the auspices of the Fellowship of Medicine.)

By L. S. T. BURRELL,

M.D., F.R.C.P.,

Physician to the Royal Free Hospital, and to the Brompton Hospital for Consumption and Diseases of the Chest.

IN a severe case of bronchiectasis, when many of the bronchi have large saccular dilatations, the diagnosis is quite simple. The patient coughs up a quantity of sputum which may be extremely offensive, and which contains pyogenic organisms but no tubercle bacilli. Physical signs vary according to whether the bronchial tubes contain sputum or have been emptied. The bases of the lungs are usually affected and not the

apices, as in tuberculosis, and there is almost always considerable clubbing of the fingers.

In less severe cases the diagnosis is often a matter of difficulty, and in some very slight cases it is possible to arrive at a correct diagnosis only after injecting lipiodol into the bronchi and making an X-ray examination.

There is one form of bronchiectasis, described by certain French physicians as the *forme sèche hémoptoïque* in which the patient suffers from periodic attacks of hæmoptysis, but between the attacks remains quite fit and free from symptoms.

The most common error is to confuse bronchiectasis with pulmonary tuberculosis. In every sanatorium it is common to find a bronchiectasis patient who has been wrongly diagnosed.

This is a serious mistake, not only because the patient is not having the proper treatment for bronchiectasis, but because he is unnecessarily exposed to the restrictions and expense of a consumptive's life. There are three important facts to remember in making a diagnosis :—

(1) Although basal tuberculosis of lung does occur, it is uncommon. If therefore signs are found at the base, the case is unlikely to be one of tuberculosis.

(2) In the early stage of pulmonary tuberculosis, when the sputum first appears, tubercle bacilli may not be found. When, however, the sputum becomes purulent tubercle bacilli are practically always present if the case is one of tuberculosis. Any case of a few months standing with purulent sputum, but without tubercle bacilli, is probably not one of tuberculosis. It is important to bear this in mind, as it is not generally appreciated and one often hears it said that failure to find tubercle bacilli means nothing. Sir James Kingston Fowler collected 186 cases of pulmonary tuberculosis, and of these tubercle bacilli were found in the sputum at the first examination in 167.

(3) Clubbing of the fingers is almost

always present in bronchiectasis and is of the drumstick type. In tuberculosis clubbing is rare, and when it does occur it consists in a mere curving of the nails and not the definite drumstick clubbing seen in bronchiectasis or congenital heart disease.

In making a diagnosis one must also remember that hæmoptysis is very common in bronchiectasis. In a series of cases collected by Dr. Acland, hæmoptysis of a pint or more occurred in 28 per cent. of the patients. Seventeen out of twenty-four cases collected by Osler had hæmoptysis. Finally, the diagnosis can be made with certainty by X-ray examination after the injection of lipiodol. I inject it through the cricothyroid membrane, as this causes less discomfort to the patient than if given through the larynx. It is a very simple method of giving lipiodol; does not make the patient choke or cough and is perfectly painless. Half a c.c. of 5 per cent. cocaine is injected into the trachea through the cricothyroid membrane. This makes the patient cough so that the cocaine spreads and anæsthetizes the trachea. When the needle is in the trachea the cocaine should be injected quickly and the needle withdrawn, or it may be broken when the patient coughs. After the attack of coughing has stopped, the patient is propped up with pillows and made to turn slightly towards the affected side, so that the lipiodol may flow chiefly into the right or left bronchus, as desired. Twenty c.c. of lipiodol are then injected through the cricothyroid membrane. If the lipiodol is warmed and the syringe used has a piston worked by a screw, the injection can be easily and quickly made. X-ray examination is made directly afterwards, and films should be taken in the recumbent and erect positions.

CAUSE.

Inflammation of the bronchial wall leading to weakness causes a bulging of the bronchial tube, especially under pressure. Any condition therefore which weakens the bronchial wall is likely to lead to

bronchiectasis. If the surrounding lung is healthy and expands normally with inspiration, the bronchial tube is protected to a great extent, but if, owing to fibrosis or some other condition, part of the lung does not expand with inspiration, there is a strain on the bronchial tubes and a tendency for them to dilate in their weakest parts. It is plain, therefore, that an ideal condition for the production of bronchiectasis would be partial obstruction of a bronchus with diminished air entry, and consequent impaired expansion of the part of the lung supplied by that bronchus, and also pent-up secretion in the distal part of the bronchus, causing inflammation and weakening of the bronchial tubes.

TREATMENT.

The first step before starting treatment is to find a cause for the disease. A bronchus may be obstructed by neoplasm, aneurysm, foreign body, or occasionally gumma, and if the obstruction can be removed the disease will be cured. I saw one patient with a small fibroma growing in a bronchus. An X-ray examination, after injecting lipiodol, showed exactly where the obstruction was present, and the fibroma was subsequently removed through a bronchoscope. A foreign body is by no means an uncommon cause of bronchiectasis, and it is remarkable how large an object can get into the bronchial tubes unknown to the patient. A man who complained of nothing more than a little bronchitis had a violent fit of coughing one morning after a cold bath, and coughed up a piece of pipe stem over an inch in length. Some eighteen months previously he had fallen off his bicycle, and on picking himself up had found that he had broken the stem of the pipe which he was smoking. He was unhurt, and had no idea that the missing pipe stem was in his bronchi until he coughed it up.

The throat, nose and nasal sinuses should be examined and treated if necessary. Nasal sinusitis is especially common in

bronchiectasis, which often improves after treatment of the sinuses.

If one is unable to find any cause which can be removed, one must then treat the bronchiectasis and symptoms to which it gives rise.

The first principle in the treatment is to empty the bronchial tubes, for not only does this make the patient less toxic, but it also tends to prevent the disease from spreading, and to lead to a gradual fibrosis, and even occasionally to a natural cure. Posture is the best method of drainage, and the patient should lie with his hips in bed and his head and shoulders bending over the floor. Sometimes it is better to bend to the right side, and sometimes to the left. This will depend on the site of the disease. Each patient will soon find in which position the flow of sputum is most readily started. In mild cases it is sufficient to empty the tubes in this way every morning, but in severe cases it will be necessary to do so two or more times during the day.

Drugs are of little value. If the sputum is offensive, creosote given by the mouth may render it less so, and potassium iodide will loosen it in the rare cases in which it is difficult to expectorate. To deodorize the sputum and encourage expectoration creosote given by inhalation is a far more satisfactory procedure. Commercial creosote is put into a metal bowl, which is heated by a spirit lamp so that creosote fumes are given off. It is best to have a special small room for this, and patients can be seated round the bowl and inhale the fumes. At first their eyes should be covered up, as the creosote is irritating, but after a time patients can remain in the room for a long time without any inconvenience.

Those who advocate more drastic treatment argue that it is irksome for a patient to "stand on his head" once or twice a day, and impossible for him to carry a portable creosote chamber about with him for the rest of his life. I would, however, repeat

and emphasize that these simple methods do tend to diminish the disease and may lead to an actual cure. If, however, the disease persists in spite of such treatment, further steps may be considered, but one must weigh the dangers and possible, or rather probable, failure of further treatment against the discomforts and dangers of the patient's present condition. If he is able to lead a fairly normal life and to attend to his work, in spite of the disease, he may prefer to leave matters where they are rather than embark on treatment of doubtful value.

If, however, it is decided that something further should be done, there are three methods which should be considered first, because they are not dangerous, cannot make matters worse and often give good results. These are:—

- (1) Artificial pneumothorax.
- (2) Phrenic evulsion.
- (3) Treatment through a bronchoscope.

ARTIFICIAL PNEUMOTHORAX.

This treatment is very successful in unilateral cases without adhesions. As many cases are bilateral and most have a considerable amount of adherent pleura, there are only a few cases suitable for pneumothorax treatment. Out of twenty-one cases in which I have tried to induce a pneumothorax for bronchiectasis, I have been able to obtain a full collapse in two only, and both of these are now quite well. If the symptoms disappear when pneumothorax is induced, the lung should be kept collapsed for six months and then allowed to re-expand. If the symptoms return as the lung expands, it should again be completely collapsed. Eventually it will be found that the lung can expand without causing symptoms. Sometimes when adhesions prevent a complete collapse the partial collapse is sufficient to relieve the symptoms. In these cases, as time goes on, the adherent surfaces tend to spread and obliterate the pneumothorax cavity by drawing out the lung. If this happens before a satisfactory

fibrosis of the diseased part has occurred, the symptoms will return, but in many cases it is possible to keep up the pneumothorax long enough to allow sufficient fibrosis to reduce if not actually to eliminate the symptoms.

PHRENIC EVULSION.

This is a simple and safe operation which leads to paralysis of the diaphragm on the affected side, and so gives additional rest to the lung and encourages fibrosis. I have not been impressed by the results of the operation, but good results have been recorded by others, and in some of my cases there has been slight improvement. The operation is chiefly indicated in cases where there are adhesions to the diaphragm preventing a satisfactory collapse of lung. It is also performed in many cases as a preliminary to thoracoplasty.

TREATMENT THROUGH A BRONCHOSCOPE.

Pus may be aspirated from the dilated bronchi by means of a suction pump and bronchoscope. In some cases, where the mucous membrane of a bronchus is congested and the lumen of the bronchus is obstructed, a large amount of pus is pent up in the distal dilated bronchial tubes, and the patient is much relieved when this is aspirated. It is also possible to wash out the tubes through the bronchoscope and, if desired, to paint or apply some drug to the diseased bronchial wall. The proceeding is very uncomfortable, but patients become quite used to it and will undergo the discomfort for the benefit derived from the treatment.

If, in spite of any method of treatment along the lines mentioned, the patient remains unrelieved, toxic and unable to work or enjoy life, surgical treatment may be justified. Lobectomy has been performed, but the mortality from this operation is so great that one may reasonably doubt if it is ever justifiable. Two opera-

tions, however, are far less dangerous, offer a very real chance of improvement or cure, and one or other of them should certainly be considered in severe cases of bronchiectasis:—

(1) Thoracoplasty.

(2) Cauterization of the diseased portion of lung.

THORACOPLASTY.

This can only succeed in unilateral cases, and before deciding on the operation the condition of the better or normal lung should be investigated by X-ray examination, after injecting lipiodol. Sometimes phrenic evulsion is performed before thoracoplasty, in order to make the fixation of lung more complete.

Cases most likely to succeed are those where considerable improvement followed artificial pneumothorax, but where symptoms returned as the collapsed lung was drawn out by spread of adherent pleura. I have known very bad cases improved or even cured by thoracoplasty. In one case a woman who was toxic and had copious offensive sputum had thoracoplasty performed in 1917, and since then has been quite well except for an occasional mild attack of hæmoptysis. In some cases thoracoplasty fails to compress the bronchiectatic cavities, which can be seen after lipiodol has been injected to be as large as before the operation.

Of the more serious methods of treatment thoracoplasty is certainly the best. The mortality is small if a surgeon of experience is chosen and, even if the operation fails to relieve the symptoms, it does not apparently make matters worse, although of course a great deal of lung tissue is permanently put out of action.

CAUTERIZATION.

E. A. Graham (*Journal of the American Medical Association*, 1923, lxxxi, 1010), described a method of treating bronchiectasis

by cauterizing the diseased tissue. It is done in several stages, and a bronchial fistula forms leading to drainage of the bronchiectatic cavities. The diseased portion of lung gradually sloughs away and the bronchial fistula heals. He states that hæmorrhage, if it occurs, can be readily controlled. In severe cases, when other methods have failed, I think this procedure holds out the best chance for the patient. It is more satisfactory than simple drainage, and has not the serious dangers of lobectomy. Of course it is chiefly indicated when the disease is localized to one portion of the lung, and cannot be used in cases where the bronchiectatic cavities are scattered over a large part of one or both lungs. Before embarking on this treatment lipiodol should be used and an X-ray examination made, as one frequently finds that the disease is far more widespread than one believed from clinical examination.

It is easy nowadays to follow the effects of treatment by X-ray examination after injecting lipiodol. It will sometimes be found that after phrenic evulsion the bronchiectatic cavities are not diminished in size, but simply raised higher up in the thorax as the dome of the diaphragm is raised. In artificial pneumothorax we may see considerable reduction in the size of the cavities, but occasionally it is found that the lung tissue collapses though the bronchiectatic cavities are unaffected. Similarly, after thoracoplasty one may find no diminution in the size of these cavities.

In conclusion, I would repeat the necessity of looking for the cause of the disease before embarking on treatment. Do not forget the possibility of a foreign body. Always determine the extent of the disease, and whether it is unilateral or bilateral, by means of X-ray examination after injecting lipiodol. Lastly, avoid the more serious surgical methods of treatment, unless the medical ones have failed and the condition of the patient is sufficiently bad to justify the risk.

EDITORIAL NOTES.

"THE SERVICES."

IN his Presidential Address to the War Section of the Royal Society of Medicine, Surgeon Vice-Admiral Arthur Gaskell, F.R.C.S., described some of the "Professional Opportunities of the Service Medical Officer." From his wide experience and official position his observations should be of especial interest to recent post-graduates as well as to undergraduates in medicine, their friends and advisers. The following were among the professional opportunities that he emphasized. One most valuable was the opportunity and time for reading, whether for advanced and special diplomas, which the doctor in active general practice could not usually obtain. Unlike the doctor in general practice he had at more less regular intervals of his proper work on board ship or with his regiment, ample opportunities for study and, what was even more valuable, for attending hospital practice. In the Services a medical officer while enjoying full pay might not only find time for such work, but might even obtain special leave for the purpose. Post-graduates belonging to the Services had also exceptional access to the study of disease in the Service hospitals, which provided clinical material large in amount and varied in type. He pointed out what we think is not generally known or appreciated, that the patients in the large Service hospitals included not only the men on the active list, but also men from the retired list. There were exceptional opportunities for major surgery and exceptional opportunities in certain directions, notably in venereal diseases and in tropical disease. A wise specialism was encouraged in the Services, and those who had had the experience which enabled them to obtain special diplomas were given specialist employment with a corresponding increase in stipend. It might be said that the pay of the Service medical officer was not so high as that of the practitioner in