and in which foci of myelomatosis were discovered in the skeleton at post-mortem examination; while Divertie (1959) has described a case of oesophageal obstruction due to amyloid deposits complicating multiple myeloma.

Although peripheral neuropathy, as a complication of myelomatosis, has been described (Hassan and Yousef, 1959; Barron, Roland and Zimmerman, 1960), it is improbable that the intestinal obstruction in the case reported could have been due to a neuropathy involving the autonomic nerve supply of the bowel, because the clinical features of the obstruction were not those of a paralytic ileus.

The information available from the study of this case is not sufficient to enable one to put forward a convincing explanation for the obstruction. It is reasonable to suppose, however, that the presence of myelomatosis was somehow related to the pathogenesis of this obstruction. Although there was no evidence of an extra-medullary plasmacytoma, nor was there any macroscopic evidence of amyloid infiltration, it is conceivable that the in-

testinal obstruction was due to the oedema and the subserous hæmorrhages, as is sometimes seen in cases of Henoch's purpura, and that these changes in the bowel wall were unknown manifestations of diffuse myelomatosis.

Summary

A case of myelomatosis, presenting with acute intestinal obstruction, has been described. At operation, only oedema and subserous hæmorrhage in the bowel wall were discovered, without evidence of mechanical obstruction. It appears from the literature, that all cases of myeloma presenting with intestinal obstruction previously reported were due to the presence of plasmacytoma in the bowel. The pathogenesis of the obstruction is discussed.

I am grateful to Professor A. G. R. Lowdon for permission to publish this case, and wish to thank Mr. F. C. Walker for his encouragement and Dr. E. A. Cameron for his guidance. I am indebted to Dr. R. B. Thompson for kindly examining the bone marrow.

REFERENCES

- ANNAMUNTHODO, H., and ROBERTSON, W. (1959): Primary Myeloma of the Stomach, *Brit. J. Surg.*, **449**.
 BARRON, K., ROLAND, L., and ZIMMERMAN (1960): Neuropathy with Malignant Tumour Metastases, *J. nerv. ment. Dis.*, **131**, 10.
 DIVERTIE, M. (1959): Oesophageal Obstruction due to Amyloid Deposits Complicating Multiple Myeloma, *Proc. Mayo Clin.*, **34**, 529.
 DANCATER, C. P., HUSSEIN, O. A., and JACKSON, W. (1959): Clinical Manifestations of Multiple Myelomatosis, *Postgrad med. J.*, **35**, 662.
 HAMPTON, J., and GRANDY, J. (1957): Multiple Myeloma of the Gastro-intestinal Tract, *Ann. Surg.*, **145**, 415.
 HASSAN, A. H., and YOUSEF, A. (1959): Neurological Complications of Multiple Myelomatosis, *J. Egypt. med. Ass.*, **42**, 386.
 HEFFERMAN, A. (1947): Plasmacytoma of Pancreas and Duodenum causing Acute Intestinal Obstruction, *Lancet*, **i**, 910.
 INNES, J., and NEWALL, J. (1961): Myelomatosis, *Ibid.*, **i**, 239.
 LOWDON, A. G. R. (1951): Deflation of Distended Bowel at Operation, *Ibid.*, **i**, 1103.
 MARTIN, N. A. (1961): The Incidence of Myelomatosis, *Ibid.*, **i**, 237.
 OSSERMAN, E. F. (1959): Plasma Cell Myeloma—Clinical Aspects, *New Engl. med. J.*, **261**, 952.
 SHARMER, K. D., SHRIVASTAV, J. D. (1961): Extra-Medullary Plasmacytoma of Gastro-intestinal Tract and Review of the Literature, *Arch. Path. (Chicago)*, **71**, 229.
 SNAPPER, I., TURNER, L., and MOSCOWITZ, H. (1953): Multiple Myeloma. New York: Grune and Stratton.

JUGULAR PHLEBECTASIA

E. LEIGHTON JOHN, M.B., B.S., F.R.C.S.(Edin.)

Neath General Hospital, Glamorganshire

JUGULAR phlebectasia is a rare anomaly of the neck veins and is of interest and importance in the differential diagnosis of swellings in the neck. Very few cases are reported in the literature, and the following case occurred in a young child.

Case Report

A 7-year-old girl was seen in January, 1961, regarding a swelling in the neck which had been noticed intermittently since birth. The story was a curious one, in that the swelling varied in size at different times, and apparently became most prominent when the child was

emotionally upset, while at other times it was not visible. There were no previous ailments.

On Examination. A soft swelling, fluctuant in character, was visible and palpable on the right side of the neck below the angle of the jaw, and over the upper third of the sternomastoid muscle. The swelling was smooth and rounded and approximately 5.0 cm. x 3.5 cm. in size.

It was noticed to vary in size during the examination and at times disappeared completely (Fig. 1). Direct pressure could empty the swelling completely and it would reappear slowly.

It was also observed that the swelling enlarged

FIG. 1.

FIG. 2.

markedly when pressure was applied below it at the root of the neck above the clavicle (Fig. 2). Also, to a lesser degree the swelling became prominent on rotation of the neck to the right.

There was no pulsation; no bruit was heard. It did not transilluminate. There was no movement on swallowing.

A diagnosis of a venous cyst freely communicating with the venous system in the neck was made.

Investigations. It was thought wiser to defer puncture of the swelling until immediately prior to operation, when it could be carried out under anaesthesia in the operating theatre.

A plain X-ray of the region showed no air visible and therefore the swelling was not an aerocoele or laryngeal diverticulum.

Carotid angiography: Right common carotid puncture. 'No abnormality shown in the arterial phase. There is an ovoid structure lying between the angle of the mandible and the hyoid bone which appears to fill from the facial veins. The internal jugular vein is normal. The exit vessel of the sac is not shown. No arterio-venous communication is present. The diagnosis suggests a venous abnormality of the neck' (Fig. 3).

Operation (E.L.Y.). Under general anaesthesia, the cyst was first aspirated and dark blood withdrawn. An operative angiogram was performed by injecting Hypaque solution into the cyst and taking an X-ray; this showed a dense opacity conforming in size and shape with the swelling in the neck. The external jugular vein was kept compressed at the root of the neck and the sac remained filled, suggesting that drainage was via the external jugular vein (Fig. 4). A curved incision was made below the right mandible. The swelling was dissected free and found to connect terminally with the external jugular vein. It appeared clearly as a dilated fusiform swelling of the vein itself and a dozen separate venous tributaries were arising from the cyst, necessitating ligation and division (Fig. 5).

The wound was closed with fine sutures and a firm dressing applied. Healing was uneventful.

Pathological Report

'The excised tissue is part of an imperfectly formed large vein. In parts of the wall there is a well defined intima, a muscular tunica media and a collagenous adventitia.

'In other parts of the wall the medial coat is repre-

FIG. 3.

FIG. 4.

FIG. 5.

sented by a few haphazardly arranged smooth muscle fibres and in adjacent areas of the wall of the vessel the media is replaced by irregular islets of adipose tissue. A few small groups of fat cells even penetrate the outer layer of the tunica intima.

'The histology suggests a vascular developmental abnormality? a vascular hamartoma?'

Discussion

The term 'phlebectasia', according to Gerwig (1952), is used to describe an abnormal fusiform dilatation of a vein, and differs from the term varix or varicose vein which implies tortuosity and dilatation

of a vein. It is a rare condition and was first described in 1928 by Harris of Toronto.

It may be seen in either the internal or external jugular vein (more commonly the latter), and appears as an abnormal spindle shaped swelling which is due to a developmental abnormality of the vein wall itself.

Such dilatations have been referred to as 'venomas', thereby likening the process to cystic hygroma arising in the lymphatic system in the neck.

It was possible in this case to establish a diagnosis of venous cyst by physical signs in the first place, later confirmed by X-rays. The swelling was of soft consistency and cystic, but cystic hygroma or lymph cyst could be excluded however, as it did not transilluminate. Any arterial connection was excluded by the absence of pulsation and bruit.

Pataro, Crosbie and Martinez Conde (1961) state that the one characteristic present in their series of four cases, was the increase in size of the tumour when the patient made an effort (e.g. performing the Valsalva manoeuvre) or to a lesser degree on lying down. They mention three types of swellings in the neck which show these features as follows:

- (1) A tumour of the superior mediastinum. This can be demonstrated by a chest X-ray.
- (2) A laryngeal diverticulum. This can be excluded by laryngoscopy and by normal X-ray appearances of the larynx and trachea.
- (3) The neck venous tumour.

REFERENCES

- GERWIG, W. H. (1952): Internal Jugular Phlebectasia, *Ann. Surg.*, **135**, 130.
 HARRIS, R. I. (1928): Congenital Venous Cyst of the Mediastinum, *Ibid.*, **88**, 953.
 PATARO, V. F., CROSBIE, J. C., and MARTINEZ CONDE, R. (1961): Jugular Phlebectasia, *J. Cardiovasc. Surg.*, **2**, 3.

MYXOEDEMA FOLLOWED BY ADDISON'S DISEASE AND DIABETES MELLITUS

G. FARRER-BROWN, M.A., M.B., B.Chir.

Late House Physician, Central Middlesex Hospital, Park Royal, London, N.W.10

BERNSTEIN (1948) described a patient with diabetes mellitus, Addison's disease and hypothyroidism, but there was no evidence of hypopituitarism found at post-mortem. This rare combination of diseases associated with apparently normal pituitary function has also been reported in cases by Howard, Mills and Mathews (1952), by Crispell and Parson (1952), and by Christy, Holub and Tomasi (1962), the last two patients also having hypogonadism. Beaven, Nelson, Renold and Thorn (1959) reported a case of a woman who had been treated for 'incomplete hypothyroidism' since childhood and who

Present Address: Bland-Sutton Institute of Pathology, Middlesex Hospital, London, W.1.

The diagnostic feature in the present case was the remarkable way in which a swelling could be made to appear, by pressure on the neck in the supra-clavicular fossa, with the patient erect, and the slow subsidence of the swelling on withdrawing the hand. This manoeuvre could be constantly repeated and proved the swelling to be a cyst communicating freely with the venous system in the neck.

If this sign occurs, it would seem to establish the diagnosis of phlebectasia.

Summary

A case of jugular phlebectasia is described, and the few accounts in the literature of this anomaly are quoted. The diagnosis is discussed and the clinical signs that were present are described.

As well as Valsalva's manoeuvre, simple pressure low down at the root of the neck followed by sudden withdrawal of the hand may be sufficient to demonstrate this condition. It can be confirmed by needle aspiration and injection of radio-opaque material followed by X-ray immediately prior to operation.

My thanks are due to Dr. D. F. Street for the X-rays and angiography and to Dr. D. B. Richards for the pathological report on the specimen.

I also wish to thank Dr. F. J. Davis for the photographs.

later developed Addison's disease and diabetes mellitus. Two other cases have also been traced in the English literature (Breslaw, Lashof and Klein, 1953; Beaven and others, 1959), but in each case with the hypothyroidism following treatment of thyrotoxicosis.

The patient described here had myxoedema for several years before developing Addison's disease and later diabetes mellitus.

Case Report

M.F., a housewife, aged 52, first attended the Out-patients' department in 1951, complaining of two recent attacks of vomiting, diarrhoea and abdominal pain for which no cause could be found, although a barium meal