whilst a leisurely (and almost certainly incomplete) chest aspiration is performed, and the relatively spent solution of Mustine and its produces are then instilled through the aspiration set. Under these circumstances satisfactory results will not be obtained. Published reports give an average of 50–60% of patients adequately controlled with Mustine, for example, Levison (1961). Personal experience suggests that 20 mg of freshly prepared Mustine introduced into a nearly empty pleural space in 200 ml normal saline controls 70% of effusions. Haematological disturbance is usually not marked, but systemic symptoms may be prominent for up to 48 hours. The very high figure of 87% with in-life control was reported by Kinsey, Carter & Klassen (1964), who used 30 mg of Mustine in 150 ml normal saline which was withdrawn after 4 hours, presumably in the hope of minimizing the systemic disturbance. In this series the lung cancer patients had an average survival time of 4.8 months. The local irritative effect of Mustine is probably at least as important as its antineoplastic effect; this may explain why Thio-Tepa, which produces little serositis, is less effective than Mustine. A further disadvantage of Thio-Tepa is that repeated administration is necessary.

Regrettably, cases are still referred for treatment after the (predictable) failure of intrapleural cyclophosphamide to control an effusion. Cyclophosphamide is activated in the liver, and is unsuitable for regional use.

4. *β -Emitting radio-isotopes.* Radioactive colloidal gold is obsolete in the management of pleural effusions, although it is still sometimes useful for ascites, especially when the ascites is caused by ovarian carcinomatosis peritonei. MacKay (1957) reported benefit in only 50% of cases, and noted that although there was a marked and consistent fall in the concentration of malignant cells in the fluid after radiogold, there was often no effect on the rate of formation of fluid. This is consistent with the hypothesis that subpleural lymphatic blockage is the major cause of fluid accumulation, since the finite

range of the β -particle is less than the depth of many of the lymphatics. The 10% γ -ray component of radiogold can have little therapeutic effect, but causes avoidable radiation exposure, which may reach 50 mr per hour at the foot of the patient's bed. Radioactive Yttrium is a pure β -emitter, and Walter (1960) reported benefit from its use in colloidal form, but it has not become popular. Radio-isotopes are an expensive and relatively ineffective way of irritating the pleura.

It must be conceded that many patients can be satisfactorily managed by repeated simple aspiration, but this approach can be troublesome when it fails, since loculation and encystment are not uncommonly the results, and they make subsequent treatment difficult.

In conclusion, the author's personal preference for the control of malignant effusion in lung cancer is Mustine, because it is effective, expeditious, and cheap. For effusions associated with other primary neoplasms, other approaches are often more appropriate.

References

- BORDA, I. & KRANT, M. (1967) Convulsions following intrapleural administration of quinacrine HCl. *Journal of the American Medical Association*, **201**, 1049.
- DOLLINGER, M.R., KRAKOFF, I.H. & KARNOFSKY, D.A. (1967) Quinacrine in the treatment of neoplastic effusions. *Annals of Internal Medicine*, **66**, 249.
- JONES, E.R. (1969) Treatment of recurrent malignant effusion by iodised talc pleurodesis. *Thorax*, **24**, 69.
- KINSEY, D.L., CARTER, D. & KLASSEN, K.R. (1964) Simplified management of malignant pleural effusion. *Archives of Surgery*, **89**, 389.
- LEVISON, V.B. (1961) Nitrogen mustard in the palliation of malignant effusions. *British Medical Journal*, **1**, 1143.
- MACKAY, N. (1957) Radioactive colloidal gold in the treatment of pleural and peritoneal effusions of malignant origin. *Lancet*, **ii**, 761.
- VASSEY, J.W., EDMONDS, J., IRVIN, J.L., GREEN, J.L. & IRVIN, E.M. (1955) Studies on the administration of quinacrine to tumour bearing mice. *Cancer Research*, **15**, 573.
- WALTER, J. (1960) Malignant pleural effusion treated by colloidal radioactive yttrium silicate. *British Medical Journal*, **2**, 1282.

Discussion

Chairman: DR J. R. MIKHAIL
(Central Middlesex Hospital)

DR MIKHAIL (Central Middlesex Hospital): You were talking about putting nitrogen mustard into pleural effusions. Do you advocate that the fluid is drained completely off just before you install the nitrogen mustard? It has been our practice always to leave some fluid there, then tipping the patient every 10 min to get complete

coverage of the pleura. It is a very painful procedure to put nitrogen mustard into a dry pleura and because of this I have always left a certain amount of fluid. I wondered what are your views on this.

DR SUTTON (Hammersmith): I quite agree it is difficult to get adequate coverage of the pleura and so what I

usually do is to put the mustine in about 200 ml of normal saline. I imagine that the normal saline not containing high amounts of protein is perhaps more likely to be re-absorbed back into the general circulation than would a high protein fluid deliberately left inside and perhaps containing malignant cells. I think it is a marginal issue. Yes, I agree it is painful and these people should have adequate premedication and analgesia following it.

DR G. POOLE (Hammersmith): Could I clarify just one point. When you began your talk you mentioned that the survival time with the definite malignant effusion from carcinoma of the bronchus was very short, about 4 months. I am not clear when you speak of the various treatments whether you are suggesting they make the life of the patient more comfortable and prevent further aspirations or whether you are suggesting they affect the length of life as well.

DR SUTTON: No, I am sorry but that was quite intentionally ambiguous. Unfortunately for the sake of brevity I have rather obscured the issue that the 90% response rate was an unselected series of cases. For instance, in that series there were twenty-three cases of carcinoma of the bronchus and about forty of various other disorders, mostly carcinoma of the breast. Twenty out of twenty-three of carcinoma of the bronchus cases did not require further aspiration in their lifetime and they survived between 4–6 months.

PROFESSOR BLEEHEEN: I think one should not lose sight of mepacrin here. On the whole I use this routinely rather than nitrogen mustard, and my impression is that the response is as good and is less disturbing to the patient. Mustine is quite disturbing to the patient over the subsequent 2–3 days.

DR SUTTON: I would not argue the point strongly at all. You may have to administer your mepacrin for 5 days in a row which is marginally more tedious than giving one shot of something.

DR SCADDING (Middlesex Hospital): I was going to mention mepacrin as I am an advocate of this. It is very seldom that we have to use it for 5 days. May I ask also in the breast cases to which you refer, could you make it clear to us whether they are having additional treatment, say Oestrogens or have you separated out those who have just had local treatment of the effusion.

DR SUTTON: The way these series are reported it is not possible to do that, and this a point I tried to make earlier, that in all three of those diseases, lymphoma, carcinoma of the ovary, carcinoma of the breast, systemic therapy is available, and this reduces the significance of conclusions you can draw from results in those diseases.

DR WADDELL: I was not terribly clear by what you meant by removing all the fluid first. Do you mean that you try to produce a normal X-ray or do you aspirate the fluid until you cannot get any more or do you aspirate until the patient is too distressed? I get the impression that a lot of these patients had rather rigidly adhered pleura either from metastasis or sub-pleural involvement by the time you first see them. And when you try to

aspirate as much as you can you are left with an X-ray which is still opaque and you don't really know what you are doing at that stage.

DR SUTTON: I think really the point I was trying to make, and perhaps I have over-stressed it, is that I do not think it is much good somebody sitting down with a needle and a syringe behind the patient, pulling enough off until the patient begins to wonder if he is getting short of breath and then banging some agent in. I would have thought it better to put in a drain and leave it in if necessary for 24–36 hr, re-inflate that lung gently over that period and put your treatment in when you have got as much out as you were likely to get out.

It seems to be a rather controversial suggestion. One or two people look distressed when I said that. I would be interested to hear objections.

DR STRADLING: Just a point of clarification. You said at the beginning you felt that the radio therapeutic effect or 'ray' effect of any of the drugs you put in was not important, and it is the mechanical effect that you were after really. And yet you say it is mustine that you prefer. Is there a 'non-ray' effect of mustine or are you using the 'ray' effect?

DR SUTTON: Can I substitute cytotoxic or cell-killing for 'ray'. Again perhaps the emphasis is misplaced. Obviously the use of mepacrine is wider than I suspected. We don't use it at the Hammersmith, and to a degree I am only entitled to talk about things about which I have experience myself; I refer to the results. This has a very minor cytotoxic action and yet as you have heard it is very effective. I would have thought, not having used it and being under the impression it was necessary to give mepacrine for 5 days and that there were side effects to that also. It seemed more convenient to use mustine, of which I have most experience.

DR STRADLING: It is the cell killing effect you are after?

DR SUTTON: Yes. It is a marginal benefit if you like.

DR STRADLING: Has it other effects?

DR SUTTON: Oh yes.

DR STRADLING: Cell-killing effect is quite incidental?

DR SUTTON: Incidental, yes.

DR MCHARDY YOUNG: I was also confused as to why radio-active gold does not seem to work. It seems to work on intraperitoneal carcinomas, does it not? I just wondered if you had any further explanation. It is not just a beta emission it is also a gamma, although these may not be effective.

DR SUTTON: Only 10% of the emission is in fact gamma and it is not really likely to have a very useful therapeutic effect. I mentioned that we had only performed ten interpleurals at the Hammersmith in the last 5 years. We have performed something like thirty radio-gold instillations and these have been quite effective. It is a slightly different situation where it is carcinoma of the ovary with myriad miliary seedlings over a very large surface. For such small seedlings the beta-ray may very well be effective, and get through it and radiate it homogeneously.