Introduction

J. M. MEDLOCK

Eastern Hospital, Homerton Grove, London E9 6BY

THE corner stones upon which the control and treatment of infectious disease have been built are isolation, immunization and chemotherapy. The need to prevent dissemination of infection as a rational procedure amongst the mumbo-jumbo of irrational therapeutics was acknowledged in the fourteenth century by the Knights Hospitallers of St John of Jerusalem in Rhodes by the construction of single rooms and separate sewage systems (Hume, 1938). In Britain, the distinctive dress and clapper to identify the leper when he left the lazar house are part of history and the disappearance of the disease from this country by the seventeenth century may be a tribute to the effectiveness, if not the humanity, of these measures. In the absence of widespread immunization, isolation remained of paramount importance to the community health until the twentieth century when the decline of the major epidemic diseases followed by the introduction of specific anti-microbial drugs suggested to the unwary that the solution of the problems of transmissible infection might be in sight. To think that these problems, which have been with man as long as man himself, should be so easily solved was naïve and the appearance of transferable antibiotic resistance, the recognition of highly lethal and infectious viral diseases unresponsive to chemotherapy and, in particular, the continuing presence of bacterial cross-infection in intensive care units, operating theatres, etc., have emphasized the need for a firm restatement by those who treat infection of the basic techniques for the prevention of cross-infection, long understood and applied by infectious diseases physicians.

In the past, emphasis has been placed on the need to prevent the infectious patient infecting others. With the increasing use of immunosuppressive drugs has come the realization that the immuno-compromised patient must sometimes be protected from the microbes of those who attend him. In the first section of the proceedings of the British Society for the Study of Infection 1975 symposium, 'Modern

Concepts of the Prevention of Infection', we are fortunate to read of P. C. Trexler's development of isolators for such patients, and then of their use in practice by those who have experience of them.

The second section is devoted to the subject of chemoprophylaxis. In some respects the word itself implies therapeutic failure for if our techniques for the prevention of cross-infection could be guaranteed, and if initial therapy was always successful, drug prophylaxis would be unnecessary. In recent years, however, the attempt to keep the well patient well or the immuno-compromised patient free of infection by the use of anti-microbial drugs has been widely practised with varying success. Three important aspects are discussed in the section.

If there has been controversy over the use of anti-microbial drugs and doubt in the minds of some of the importance of isolation, there is no denying that a vaccine which is effective and safe should be used to protect the public against infectious disease. It is sometimes forgotten, however, that those who care for the infectious patient may need protection. This is achieved by the use of sensible protective ward techniques together with immunization. Some aspects of the latter are dealt with in the third section. The immunization scene is a changing one. The success of the World Health Organization programme for the eradication of smallpox deserves wide acclaim. The controversy over whooping cough vaccine might be better keyed in low profile until more precise knowledge of the hazards and efficacy of the vaccine is available. In the meantime there are other diseases, including some the aetiology of which is only now being suggested as microbial, against which vaccines might be developed. Prospects for new vaccines are considered in the last contribution.

Reference

HUME, E.E. (1938) The medical work of the Knights Hospitallers of Saint John of Jerusalem. *Bulletin of the History of Medicine*, 6, 398.