SESSION I

Chairman: PROFESSOR N. R. BUTLER

Skeletal development

J. J. PRITCHARD
M.A., D.M., F.R.C.S.

Department of Anatomy, The Queen's University, Belfast

NORMAL skeletal development may be considered as exhibiting hierarchies of organization levels, processes and controls (Table 1). This approach is apposite to the study of bone dysplasias because developmental failure at one level inevitably, and fairly predictably, leads to undesirable sequelae at higher levels, although prediction is complicated by the organism's attempts at functional compensation for morphogenetic deficiencies. Unlike a rising building, a skeleton must function and adapt to its environment throughout its growth.

TABLE 1. Skeletal development	
Process	Level of organization
Cell recruitment	Unspecialized cells
	↓
Primary ossification (I.M., E.C.)*	Specialized cells
	↓
Secondary ossification	Primary bone / membrane
	↓ / cartilage
Tertiary ossification	Secondary bone
	↓
Co-ordinated growth of fibrous, cartilaginous, bony, etc. tissues	Tertiary bone
	↓
Co-ordinated growth of bones (local)	A bone
	↓
Co-ordinated growth of bones (systemic)	A bone complex
	↓
Co-ordinated growth of bones and muscles, etc.	The bony skeleton
	↓
	The organism

* I.M., intramembranous; E.C., endochondral.

The sources and modes of recruitment of the specialized osteogenic, chondrogenic and fibrogenic cells which are involved in skeletal development are far from clear. A scheme of osteoblast recruitment in early centres of ossification is given in Table 2. Later on it is accepted that each growing

TABLE 2. Cell recruitment	
Osteoblast recruitment	
Migration	Cells from (a) Sclerotome of somite (axial) (b) Neural crest (head region) ? (c) Lateral plate (limbs)
	↓
Differentiation proliferation	Local competent cells
	↓
Modulation (reversible)	Determined cells (osteoprogenitor cells)
	↕
Maturation	Active cells (osteoblasts)
	↓
	Mature cells (osteocytes)

bone recruits its osteoblasts from a local pool of osteoprogenitor cells present in the depths of the periosteum and in the marrow spaces. Heterotopic ossification poses problems in osteoblast recruitment and invites experimental enquiry. A list of osteoblast-inducing tissues and agencies is given in Table 3. Cells in process of becoming osteoblasts exhibit intense alkaline phosphatase activity as they acquire a massive load of cytoplasmic organelles. The list of probable and possible osteoblast functions is impressive (Table 4).

Primary ossification is either intramembranous or endochondral, the former occurring at the expense of

TABLE 3. Osteoblast induction

(1) Hyperthrophic cartilage (alive or dead)
(2) ? Nervous tissue (brain, peripheral nerve)
(3) ? Pharyngeal entoderm
(4) Transitional epithelium (proliferating)
(5) Bone (BMP* Urist) (decalcified)
(6) Dentine (Urist) (decalcified)
(7) Mechanical stress (tendon)
(8) Muscle (→cartilage→bone)
(9) ? Calcium deposits (cartilages, arterial walls)
(10) ? 'Chronic irritation'

*BMP = bone morphogenesis product

growing fibrous tissue under tension, the latter at the expense of growing cartilage under compression, and results in the formation of either primary membrane (woven) bone or primary cartilage bone. Theoretically dysplasia could result from failure of proper recruitment and/or function of fibroblasts, chondroblasts or osteoblasts at this level.

Secondary ossification involves the compacting by lamellar bone of the network of primary bone trabeculae to form secondary bone (primary osteonal bone, laminar bone, surface bone). Failure

TABLE 4. Osteoblast functions

(1) Manufacture and secretion of collagen (tropocollagen)				
(2) Orientation of collagen fibres into domains (Boyde)				
(3) Manufacture and secretion of proteoglycans and glycoproteins (? at pre-osteoblast stage)				
(4) Mineralization of organic matrix <table> <tr><td>(a) Manufacture and secretion of alkaline phosphatase (pyrophosphatase)</td></tr> <tr><td>(b) Aggregation and secretion of calcium and phosphate ions (mitochondria, matrix vesicles)</td></tr> <tr><td>(c) Provision of energy (ATP) for initiation of mineralization</td></tr> <tr><td>(d) Regulation of mineral traffic between blood and bone matrix (shared with osteoclasts and osteocytes)</td></tr> </table>	(a) Manufacture and secretion of alkaline phosphatase (pyrophosphatase)	(b) Aggregation and secretion of calcium and phosphate ions (mitochondria, matrix vesicles)	(c) Provision of energy (ATP) for initiation of mineralization	(d) Regulation of mineral traffic between blood and bone matrix (shared with osteoclasts and osteocytes)
(a) Manufacture and secretion of alkaline phosphatase (pyrophosphatase)				
(b) Aggregation and secretion of calcium and phosphate ions (mitochondria, matrix vesicles)				
(c) Provision of energy (ATP) for initiation of mineralization				
(d) Regulation of mineral traffic between blood and bone matrix (shared with osteoclasts and osteocytes)				
(5) ? Degradation of collagen, proteoglycans and glycoproteins				

at this level characterizes some kinds of osteogenesis imperfecta and scurvy.

In tertiary ossification the bony framework is remodelled as a result of balanced bone resorption by osteoclasts and bone deposition by osteoblasts so

FIG. 1. Cartilage models developed from embryonic rat limb bud mesenchyme transplanted to mother's rectus sheath.

that tertiary bone with Haversian systems (secondary osteons) is formed. Osteoclast failure at this level has serious and rather bizarre consequences.

In the development of a whole bone (as opposed to the formation of a bony tissue) fibrous tissue, cartilage and bone grow in concert, and there are solid grounds for the belief that this is made possible by cartilage serving as the leader of the team, with the other tissues obeying its dictates. Cartilage is well equipped for such a role. Unlike bone or fibrous tissue its growth potential is determined at an early stage in embryonic life. Thus the pre-cartilaginous vertebrae of the newborn rat's tail develop almost normally when transplanted beneath the skin of the mother's back, while rat limb-bud mesenchyme forms respectable cartilage models and ossicles when placed in the mother's rectus sheath (Fig. 1). Cartilage, in addition, can push against its surroundings as it grows, while its vascular requirements are modest. Moreover, in its hypertrophic state, cartilage is able to induce bone formation around it. It is envisaged that as the epiphyseal plates of a long bone grow they tense and orientate the collagen fibres of the periosteum, thereby stimulating collagen growth which in turn stimulates and orientates intramembranous ossification. The tense periosteum for its part tends to put a brake on the growth of the epiphyseal plates, for if the periosteal sleeve around a growing bone is severed transversely the growth of the bone in length is accelerated (Crilly, 1972). These views are summarized in Fig. 2.

The cartilage-fibrous tissue pacemaker-feedback system just outlined also appears to be important for harmony between growing bones. Harrison (1958) produced predictable distortion of the rat's pelvis by cutting the sacro-iliac ligament in early life. This ligament normally pulls the ilium caudally as growth occurs at the iliac crest, so maintaining the relative position of the sacro-iliac joint half way down the ilium.

FIG. 2. Cartilage and bone formation. Cartilage of growth plate: when young, acts as a pacemaker; when old, as a bone inducer; when dead, as a bone scaffolding. Cartilage model as a whole; determination of bone size and shape.

To what extent dysplasias are caused, or accentuated, by disharmonies between growing cartilaginous, fibrous and bony tissues remains to be determined: but on the positive side one may conclude that the making of an elegant and efficient skeleton demands good genes and a good environment: good cells (osteoblasts, osteoclasts, chondroblasts, fibroblasts) in proper numbers at the right places at the right times; a good programme of controls; and optimal mechanical, vascular, nutritional and endocrine milieux.

References

- CRILLY, R.J. (1972) Longitudinal overgrowth of chicken radius. *Journal of Anatomy*, **112**, 11.
 HARRISON, T.J. (1958) An experimental study of pelvic growth in the rat. *Journal of Anatomy*, **92**, 483.