Even with this treatment there are a few rebellious cases. Implantation of radon seeds or radium itself may be tried, and to judge by the few reported cases radium is highly beneficial.

REFERENCES.

- [1] HARGER, J. R. *Surg. Gynec. and Obstet.*, 1920, xxxi, 305 (OCHSNER, A. J., in Discussion.)
- [2] SANFORD, A. H. *Journ. Amer. Med. Assoc.*, 1923, xxxi, 655.
- [3] BAILEY, HAMILTON. "Branchial Cysts," London, 1929.
- [4] CHRIST, A. *Schweiz. med. Woch.*, 1925, lv, 643.
- [5] BOYKSEN, A. *Zentralbl. f. Chirurg.*, 1928, lv, 1542.
- [6] NEW, G. B., and FIGI, F. A. *Surg. Gynec. and Obstet.*, 1923, xxxvii, 617.
- [7] CHITTY, H. *Brit. Med. Journ.*, 1926, i, 418.
- [8] *Idem. Ibid.*, 1929, i, 347.
- [9] HALE-WHITE, Sir W. "Materia Medica," 14th Edition, 1915, p. 253, London.

"HAMMER-TOE."

By S. L. HIGGS,

F.R.C.S.

Surgeon, Royal National Orthopaedic Hospital;
Assistant Surgeon, Orthopaedic Department, St.
Bartholomew's Hospital, &c.

THE term "hammer-toe" should be confined to that type of flexion deformity which cannot be overcome by simple stretching. It therefore excludes all types of "claw-toes" which are commonly associated with the wearing of short stockings and shoes, or with other foot deformities, such as pes cavus or loss of the transverse arches.

The true "hammer-toe" cannot be straightened by traction because this is prevented by actual contracture of the skin and soft tissues beneath the flexed joint.

CAUSE.

The condition is rarely found in infants. It is an acquired deformity and the chief factor in its development is an over-long

toe. If short socks or shoes are worn the projecting toe is pressed back into line with the others and has to assume the flexed position. As growth proceeds this deformity becomes fixed and later on pressure symptoms develop.

TYPES.

The typical "hammer-toe" shows a flexion deformity of the proximal interphalangeal joint whilst the metatarsophalangeal and distal interphalangeal joints are hyper-extended. The only fixed deformity in this type is the flexed middle joint. This is by far the commonest variety and the easiest to treat.

Occasionally the terminal joint is flexed instead of extended, producing a double flexion deformity.

A third type shows flexion of the terminal joint only.

SYMPTOMS.

Pain occurs in three situations. Firstly, in the common type, over the prominent joint from pressure of the shoe. A corn and underlying bursa develop, and attacks of acute inflammation are not uncommon. Secondly, pain may be complained of beneath the ball of the foot. The hyper-extended first phalanx presses downwards the head of the corresponding metatarsal, leading to loss of the transverse arch and consequent metatarsalgia. Lastly, if the end joint is flexed a painful corn may develop around the nail at the tip of the toe where pressure occurs.

TREATMENT.

The development of a "hammer-toe" can be prevented in children by the wearing of full-length socks or stockings. Mothers

rarely appreciate the fact that their children's stockings may be too short and cramp the toes and this, far more than short footwear, is the factor which leads to foot troubles in later life.

If an over-long toe is becoming “hammer” but is not yet badly contracted, then a sole plate, with slots and chamois leather tapes to strap down the toes, should be worn at night.

A fully-developed “hammer-toe” can only be treated satisfactorily by operation. Treatment by stretching, sole plates and the like are quite useless. If operation is refused the pressure symptoms can be avoided but only at the expense of time and money. Special shoes can be made with a deep bed in the insole to accommodate the end of the affected toe and thereby depress the prominent joint. Frequent visits to the chiropodist for attention to the corns and application of felt pads will be required.

OPERATION.

Amputation of a “hammer-toe” is to be condemned. If, as is usual, it is the second toe which is affected, then the development of hallux valgus is inevitable. If it is the third or fourth toe, then transverse arch troubles may ensue and the alignment of the great toe may also, in time, be upset.

The exception to the rule is where such a severe degree of hallux valgus exists that the second toe is not only “hammer,” but also displaced upwards to overlie the great toe. In these cases, especially in elderly people when extensive operative treatment is not desirable, simple removal of the toe may be advisable.

A true “hammer-toe” deformity does not affect the little toe, but the rule against the amputation of toes does not apply, as it may be removed with any subsequent disability.

The usual operation consists of excision of the flexed interphalangeal joint to adapt the length of the toe to the shortened structures beneath. Bony ankylosis between the first and second phalanx is aimed at, but

owing to difficulty in subsequent splinting, may not be secured, and the toe may be painful for a considerable time. If fibrous union results, relapse of the deformity may occur.

I have found the following method most satisfactory :—

Two transverse elliptical incisions are made across the dorsum of the toe to include the corn over the flexed joint. One incision should be carried right to the bone, dividing and removing a section of the extensor tendon. The incisions should not be carried too far laterally, otherwise the digital nerves and vessels may be cut across. The joint is opened and the lateral ligaments divided, so that the proximal phalanx can be thrust well out through the wound. With narrow-bladed bone-forceps, this phalanx is sharpened into a spike. The cartilage from the base of the second phalanx is then removed with bone-forceps and a hole bored up into the shaft of the bone. I have found a rawl-plug borer a most satisfactory instrument for doing this. The pointed first phalanx is then spiked into the second phalanx. The divided extensor tendon is sutured with a catgut stitch, and the skin incision closed transversely with a few interrupted stitches. No splinting is required.

The foot should be rested until the stitches are removed, but the patient need not lie up entirely, as there is no objection to a few paces being taken with the heel on the ground. As a rule, an ordinary shoe can be worn in three weeks' time, and in some cases rather sooner.

If this procedure is carried out, post-operative pain is much diminished as, owing to the firm impaction of the bones, no movement occurs and firm internal splinting is secured.

In those types of “hammer-toe” where flexion deformity of the terminal phalanx is present, the best operative procedure is either to excise the head of the second phalanx, through a transverse elliptical

incision to shorten the skin over the dorsum of the joint or, where the nail is much thickened by chronic pressure, amputation of the terminal phalanx and removal of the whole nail should be performed. One long plantar flap from the pulp of the toe gives a good and comfortable stump.

DIVISION OF THE VAS DEFERENS IN PROSTATECTOMY.

By HAROLD DODD,

CH.M., F.R.C.S.ENG.

*Hon. Surgeon to the King George Hospital, Ilford,
and the Royal Hospital, Richmond.*

THE evolution, various techniques, management and complications of prostatectomy were exhaustively dealt with during the past year by Sir John Thomson-Walker, in his Lettsomian Lectures [1], by Young of Baltimore [2], and by the leading genito-urinary surgeons at the Royal Society of Medicine [3].

In this most interesting review, the complication of epididymo-orchitis and its sure prevention by ligation of the vas deferens, was stressed. Even Sir John Thomson-Walker, with his finished technique and mature experience, admitted the incidence of it in 6.4 per cent. of his cases, which is quite appreciable, but low compared with the experience of other genito-urinary surgeons, whose percentage of occurrence will probably be smaller than that of general surgeons. Against this must be placed the nine-year experience of Mr. Clifford Morson and other urological surgeons, with scarcely a case of epididymitis or orchitis. Morson makes a striking statement when he says, "I guarantee my patients that, whatever other complications they may have, they will never suffer from orchitis"; this means the [4] exclusion of a very irritating and debilitating condition, which occurs chiefly during the

first year after prostatectomy in at about 30 per cent. of cases. It frequently happens whilst patients are recovering in hospital or, as many practitioners will relate, after their discharge home, when it is explained by the statement to the patient that he "has caught a cold"; thus, on the day after his return home, one of my patients walked round his garden, and developed epididymitis next day; I think it was probably a coincidence.

This affection is excited by urethral trauma, e.g., after passing a bougie for difficulty in micturition after prostatectomy, by excess of alcohol, by exposure to cold, by the necessity of delay in emptying a full bladder and constipation.

It is a most painful and incapacitating complication, with marked pyrexia, and, unfortunately, recovery from one attack by no means confers immunity to its subsequent occurrence. Frequently, it recurs on the same or the opposite side or, occasionally, as described later, it is bilateral. Again, it often suppurates, necessitating further treatment by aspiration, or incision and drainage, or it may be necessary in a chronic case, where troublesome and persistent sinuses exist, to excise the testicle and part of the scrotum.

The origin of the infection is the prostatic bed, which quickly becomes septic after the operation and the organisms pass by the ruptured ejaculatory ducts to the vesiculæ seminales. These latter are larger than one ordinarily supposes from the impressions derived from the dissecting and post-mortem rooms and anatomy books; anyone who has seen the seminal vesicles in life, for instance during an abdomino-perineal resection of the rectum, will obtain a clear idea of the excellent but huge culture tubes which these blind, dilated vesicles make. Sepsis passes from them along the vas deferens, excited by the factors already mentioned, to produce the epididymitis.

A method of radical prevention would be to excise the vesiculæ at or before prostatectomy, as suggested by Sir J. Thomson-