Leading Article

Malignant fibrous histiocytoma

J.R. Salisbury

Department of Morbid Anatomy, King's College School of Medicine and Dentistry, Denmark Hill, London SE5 8RX, UK.

Malignant fibrous histiocytoma (MFH) was the name proposed for a group of soft tissue sarcomas on the basis of tissue culture studies which suggested the tumour cells were of histiocytic origin.¹ These sarcomas had been previously classified as poorly differentiated sarcomas of other types (particularly fibrosarcoma). As new histological criteria were established for both MFH and fibrosarcoma, many sarcomas were diagnosed as MFH with the result that MFH is now considered the commonest soft tissue sarcoma.² Further work, largely by Enzinger and his co-workers at the Armed Forces Institute of Pathology, introduced the sub-types of soft tissue MFH that are recognized today—pleomorphic-storiform, myxoid, inflammatory, giant cell-rich and angiomatoid. The histopathological features of the various sub-types have been well illustrated by Enzinger and Weiss.²

The spectrum of fibrous histiocytic tumours was expanded to include benign tumours of dermis and tendon sheath, such as dermatofibroma and benign synovioma, and soft tissue tumours of intermediate malignancy such as dermatofibrosarcoma protuberans. Bone tumours with a similar histological appearance were included as benign and malignant fibrous histiocytoma of bone.^{3–5} With increasing recognition by pathologists of the histological pattern of MFH, tumours with this appearance were diagnosed as MFH at more and more sites other than the soft tissues. Two further cases of MFH at uncommon sites are reported in this issue.^{6,7}

Grieco and his colleagues report a case of a 42 year old man with a 3-month history of pyrexia who was found to have a MFH of the sternum.⁶ The tumour was deemed inoperable and was treated with chemotherapy. Residual tumour could not be found in the sternum at autopsy. MFH is a rare bone tumour; for example, it represents only 0.8% of the 6514 primary bone tumours described from the Mayo Clinic.⁸ Sixty three per cent of tumours involve the long bones⁸ and the sternal location of the reported case is most

unusual. Like Grieco's case, the majority of osseous MFHs are of the pleomorphic-storiform type.⁹ The optimal treatment for these tumours is chemotherapy^{10,11} followed by surgical excision and, for long bone tumours, prosthetic replacement.

Aggarwal and colleagues describe a MFH of the posterior mediastinum in a 16 year old girl who presented with pyrexia.⁷ The tumour was surgically excised and the pyrexia subsided post-operatively. The mediastinum is another rare site for MFH. Since the initial description of two mediastinal cases by Mills *et al.*,¹² only single cases have been reported.¹³

A further unusual feature of the two cases of MFH in this issue is the prominent pyrexia. Many neoplasms are associated with pyrexia, notably gastric, pancreatic, hepatic and renal carcinomas and leukaemias and lymphomas. Most cases are due to associated infections or the products of tissue injury. Pyrexia in association with MFH has been noted previously¹⁴ and was present in 2 out of 200 cases of soft tissue MFH.¹⁵ Other occasional associations with MFH have included episodic hypoglycaemia¹⁵ and rapid enlargement of the tumour during pregnancy.¹⁶

Despite the evident attractiveness to pathologists of fibrous histiocytoma as a diagnosis, cracks are beginning to appear in the concept of fibrous histiocytic tumours. The diagnosis of pleomorphic MFH (the commonest sub-type) in particular is coming under strong attack with the question being posed whether it represents a real biological entity or has it become a 'diagnostic dustbin' like the poorly differentiated fibrosarcoma of old.

Doubt has been cast on the original tissue culture studies¹⁷ and it now seems probable that the tumours studied were malignant lymphomas. The finding that some cells within MFH stained with the first 'histiocytic' markers used in immunohistochemistry—polyclonal antisera against alpha-1 antitrypsin, alpha-1 antichymotrypsin and lysozyme^{18,19}—had been taken to support histiocytic differentiation but this support has been eroded by subsequent studies showing these enzymes to be rather ubiquitous,^{20,21} and by studies using newer monoclonal antibodies (with greater macrophage specificity) revealing that only a very small number of true histiocytes are present within

Correspondence: J.R. Salisbury, B.Sc., M.B., B.S., M.R.C. Path.

Received: 18 May 1989

these tumours²²—cells thought to be a transient macrophage population²³ as can be seen in many tumours.²⁴ That fibrous histiocytomas do not contain neoplastic histiocytes may be unsatisfactory to terminological purists, but it does not matter in any practical sense—what's in a name?—and our concepts of tumour histogenesis from, rather than towards, cell type may be at fault.²⁵

The accusations being levelled at pleomorphic MFH are more serious. That the histological pattern of pleomorphic MFH can be seen in the dedifferentiated parts of a number of tumours has been recognized for some years. What then, if tumours currently diagnosed as pleomorphic MFH are more thoroughly examined? By sampling the tumour even more widely than has been current practice, by using a panel of immunohistochemical markers and by sub-

mitting a portion of each tumour for electron microscopy, would evidence be found to make a diagnosis of another type of tumour? Some pathologists assert this to be so for many pleomorphic MFHs of soft tissues but most pathologists find that, even after such procedures, there remain some tumours for which pleomorphic MFH is the only creditable diagnosis.

As treatment for sarcomas becomes increasingly sophisticated, this is important; for only by knowing what exactly is being treated can we hope to improve outcome. It may be that sarcomas consisting mostly of MFH-like tumour will behave in the same biological manner but, at present, we do not really know. Because of the rarity of these tumours at some sites, case reports such as those of Grieco *et al.* and Aggarwal *et al.* will continue to have their place.

References

- Ozzello, L., Stout, A.P. & Murray, M.R. Cultural characteristics of malignant histiocytomas and fibrous xanthomas. *Cancer* 1963, **16**: 331–344.
- Enzinger, F.M. & Weiss, S.W. Malignant fibrous histiocytoma. In: *Soft Tissue Tumors*, 2nd edition. C.V. Mosby, St. Louis, 1988, pp. 273–300.
- Feldman, F. & Norman, D. Intra- and extraosseous malignant histiocytoma (malignant fibrous xanthoma). *Radiology* 1972, **104**: 497–508.
- Spanier, S.S., Enneking, W.F. & Enriquez, P. Primary malignant fibrous histiocytoma of bone. *Cancer* 1975, **36**: 2084–2098.
- Destouet, J.M., Kyriakos, M. & Gilula, L.A. Fibrous histiocytoma (fibroxanthoma) of a cervical vertebra. A report with a review of the literature. *Skeletal Radiol* 1980, **5**: 241–246.
- Grieco, A., Caputo, S., Silvestri, E., Caradonna, P., Bertoli, A. & Greco, A.V. Malignant fibrous histiocytoma of sternum: an usual cause of fever of undetermined origin. *Postgrad Med J* 1989, **65**: 926–928.
- Aggarwal, P., Sharma, S.K., Chattopadhyay, T.K. & Mathur, M. Malignant fibrous histiocytoma of the mediastinum. *Postgrad Med J* 1989, **65**: 929–931.
- Dahlin, D.C. & Unni, K.K. Malignant (fibrous) histiocytoma. In: *Bone Tumors*, 4th edition. Charles C. Thomas, Springfield, Illinois, 1986, pp. 357–365.
- Wold, L.E. Fibrohistiocytic tumors of bone. In: Unni, K.K. (ed) *Bone Tumors*. Churchill Livingstone, New York, 1988, pp. 183–197.
- Weiner, M., Sedlis, M., Johnston, A.D., Dick, H.M. & Wolff, J.A. Adjuvant chemotherapy of malignant fibrous histiocytoma of bone. *Cancer* 1983, **51**: 25–29.
- Urban, C., Rosen, G., Huvos, A.G., Caparros, B., Cacavio, A. & Nirenberg, A. Chemotherapy of malignant fibrous histiocytoma of bone. A report of five cases. *Cancer* 1983, **51**: 795–802.
- Mills, S.A., Breyer, R.H., Johnston, F.R. *et al.* Malignant fibrous histiocytoma of the mediastinum and lung. A report of three cases. *J Thorac Cardiovasc Surg* 1982, **84**: 367–372.
- Venn, G.E., Gellister, J., DaCosta, P.E. & Goldstraw, P. Malignant fibrous histiocytoma in thoracic surgical practice. *J Thorac Cardiovasc Surg* 1986, **91**: 234–237.
- Kyriakos, M. & Kempson, R.L. Inflammatory fibrous histiocytoma. An aggressive and lethal lesion. *Cancer* 1976, **37**: 1584–1606.
- Weiss, S.W. & Enzinger, F.M. Malignant fibrous histiocytoma. An analysis of 200 cases. *Cancer* 1978, **41**: 2250–2266.
- Weiss, S.W. & Enzinger, F.M. Myxoid variant of malignant fibrous histiocytoma. *Cancer* 1977, **39**: 1672–1685.
- Fletcher, C.D.M. Commentary: Malignant fibrous histiocytoma? *Histopathology* 1987, **11**: 433–437.
- Meister, P. & Nathrath, W. Immunohistochemical markers of histiocytic tumors. *Hum Pathol* 1980, **11**: 300–301.
- Du Boulay, C.E. Demonstration of alpha-1 antitrypsin and alpha-1 antichymotrypsin in fibrous histiocytomas using the immunoperoxidase technique. *Am J Surg Pathol* 1982, **6**: 559–564.
- Roholl, P.J.M., Kleyne, J., Elbers, H., Van der Vegt, M.C., Albus-Lutter, C.H. & Van Unnik, J.A.M. Characterization of tumour cells in malignant fibrous histiocytomas and other soft tissue tumours in comparison with malignant histiocytes. I. Immunohistochemical study on paraffin sections. *J Pathol* 1985, **147**: 87–95.
- Leader, M., Patel, J., Collins, M. & Henry, K. Anti-alpha-1-antichymotrypsin staining of 194 sarcomas, 38 carcinomas, and 17 malignant melanomas. Its lack of specificity as a tumour marker. *Am J Surg Pathol* 1987, **11**: 133–139.
- Wood, G.S., Beckstead, J.H., Turner, R.R., Hendrickson, M.P., Kempson, R.L. & Warnke, R.A. Malignant fibrous histiocytoma tumor cells resemble fibroblasts. *Am J Surg Pathol* 1986, **10**: 323–335.
- Lawson, C.W., Fisher, C. & Gatter, K.C. An immunohistochemical study of differentiation in malignant fibrous histiocytoma. *Histopathology* 1987, **11**: 375–383.

24. Wood, G.W. & Gollahon, K.A. Detection and quantitation of macrophage infiltration into primary human tumors with the use of cell-surface markers. *JNCI* 1977, **59**: 1081–1087.
25. Gould, V.E. Histogenesis and differentiation: a reevaluation of these concepts as criteria for the classification of tumors. *Hum Pathol* 1986, **17**: 212–215.