Reviews in Medicine

Oncology

J.S. Malpas

Imperial Cancer Research Fund, Department of Medical Oncology, St Bartholomew's Hospital, West Smithfield, London EC1A 7BE, UK.

Introduction

Given the intense interest in all aspects of oncology, any review is inevitably selective and must be strongly influenced by the interests of the author. In an attempt to avoid bias, this article has been written for the general physician, and has tried to cover the background to some modern developments in oncology, and at the same time highlight investigations or treatments reported recently that might be helpful in practice. The astonishing pace at which molecular biology is adding to our knowledge of the normal and malignant cell, and the rate at which this knowledge is being applied to diagnosis and therapy, makes this one of the most important aspects to consider.

Chromosomal abnormalities and cancer

The classic abnormality of the t(9;22) translocation resulting in the Philadelphia chromosome consistently associated with chronic myeloid leukaemia (Figure 1), and the more recently described translocations in Burkitt's lymphoma t(8;14) (Figure 2) or promyelocytic leukaemia t(15;17) have emphasized the central role of translocations, inversions and deletions of chromosomal material in tumour pathogenesis.

The way in which these abnormalities convey their survival advantage to the malignant cell is the subject of a number of hypotheses. Activation of genes which promote the production of growth-promoting substances and deletions of suppressor genes which normally prevent such activity, are being increasingly convincingly demonstrated. Allelic deletions in chromosome 13 in retinoblastoma (Figure 3), chromosome 11 in Wilms', embryonic, breast and bladder tumours, chromosome 22 in acoustic neuromas, chromosome 3 in renal cancer and lung carcinomas, are good examples of tumours in which deletions on one chromosome are associated with malignancy.

Figure 1 t(9;22)(q34;q11) showing Philadelphia chromosome.

Figure 2 t(14;18)(q32;q21) in follicular lymphoma.

Figure 3 del(13q) in retinoblastoma.

The association of chromosomal abnormality with tumorigenesis would be even further strengthened if it could be shown that an increasing incidence of chromosomal abnormality occurred as a tissue underwent malignant change. Vogelstein *et al.*¹ investigated the genetic alterations during colorectal tumour development, and were able to study the DNA in adenomas of increasing dysplastic grade and in colorectal cancer. RAS gene mutation, deletion on the long arm of chromosome 5, chromosome 18 and 17 were observed with increasing frequency as the tissue became increasingly dysplastic (Figure 4). The conclusions from

Figure 4 Genetic alterations during colorectal tumour progression. After Vogelstein *et al.* Reprinted by permission of the *New England Journal of Medicine* 319, 525, 1988. ● – RAS mutation; ○ – 5q allelic deletion; ■ – 18q allelic deletion; □ – 17p allelic deletion.

Vogelstein's work are that accumulated alterations affecting at least one dominantly-acting oncogene, and possibly several tumour suppression genes, are responsible for the development of colorectal tumours. In studying the initiating events of colorectal cancer that develop in familial polyposis, there is no doubt that the further deletion on chromosome 5 with loss of a suppressor gene is of great importance in the development of this malignancy.^{2,3}

While it would appear that molecular biology is bringing us nearer and nearer to the initiating event in cancer, a recent discovery, the polymerase chain reaction, may be of crucial importance. One of the most difficult problems for the management of clinical cancer is the detection of residual disease. If it were possible to say without equivocation that all tumour had been removed either by surgery, radiotherapy, or chemotherapy, many patients would be spared further treatment, which at the least is unpleasant, and at worst may be life-threatening. In the case of leukaemia and lymphomas, histopathology and the more refined cytological techniques have been inadequate to detect minimal residual disease. In other tumours – with the possible exception of germ cell tumours, in which α -fetoprotein and β -human chorionic gonadotrophin (HCG) have been helpful – no good markers of disease exist. The belief that the detection of minimal residual disease is important is supported by the fact that germ cell tumours in adults are among the most successfully treated of all malignancies.

Molecular biology has furnished a very powerful tool in the form of the polymerase chain reaction (PCR). It was noted earlier that the outstanding

cytogenetic abnormality in chronic myeloid leukaemia is the t(9;22) translocation. The area in which this translocation occurs is close to the *abl*-oncogene. DNA amplification can be utilized to produce many copies of the *bcr-abl* transcripts near the RNA junction (Figure 5). This provides a unique clonal marker for this tumour.⁴ Using this technique, malignant cells in concentrations as small as 1/100,000 can be detected. This technique is being extended to detect residual lymphomatous tissue. The majority of follicular lymphomas show a t(14;18) translocation, which results in removal of the immunoglobulin heavy chain gene into the *bcl-2* gene on chromosome 18.⁵ Sixty per cent of the translocation occurs within a narrow region on the *bcl-2* gene, known as the major breakpoint region, and the rest on a region known as the minor breakpoint cluster. Probes for these regions are available. PCR reactions across this major breakpoint region can be used to detect malignant cells with a sensitivity of up to 1/100,000 cells.⁶ At the moment, these techniques are most easily applied to the detection of minimal residual disease in the bone marrow of patients with acute leukaemia and lymphoma, but a wider application of the technique is now being sought so that it might be helpful in certain solid tumours.

Cell culture

Study of the effect of cytotoxic drugs – influence of growth factors

This section allows a brief look at new aspects of malignant cell culture *in vitro*, the practical application of such a system in myeloma and a look at the impressive number of regulatory molecules or cytokines that are now being identified and tried in clinical practice. It is a truism that new techniques open up new fields of pure and applied research. The difficulty of growing myeloma cells reliably has prevented research both on the nature of this B cell and its response to therapy. Using a double layer agar technique, Millar *et al.*⁷ have been able to culture myeloma cells so that they grow with a high plating efficiency without contaminating B, T or granulocyte macrophage colonies. Bone marrow containing as little as 2% infiltration consistently grew colonies of small lymphoid cells and larger plasmacytoid cells, which can be demonstrated to arise from the original myeloma cells. It is interesting to note that these colonies could even be grown from the peripheral blood. The drug sensitivity of myeloma cells compared with normal haemopoietic cells can be measured by using dose response curves in individual patients. This method can be used to detect resistant subpopulations of myeloma cells. In serial bone marrow samples the emergence

Figure 5 Polymerase chain reaction (PCR) is a means of increasing small quantities of known DNA or RNA sequence to an easily detectable level. The schematic diagram illustrates the process PCR. Initially double stranded DNA is placed in the reaction mixture consisting of 2 short sequences (Primers) of DNA at either side of the sequence to be copied, the four individual bases (adenine, thionine, cytosine and guanine) and Taq enzyme that is required to link the bases in the manufacture of DNA. A cycle of temperatures is followed. Initially the double stranded DNA is broken into 2 single strands (denatured) by heating to 94°C. The temperature is then lowered to around 55°C to encourage the 2 primers to attach to the complementary sequence of single stranded DNA (annealing). The temperature is then raised to 72°C, the ideal condition for Taq enzyme to incorporate the appropriate bases onto the starting primer sequence (extension) producing 2 double stranded segments. This cycle usually takes no longer than 3 minutes and is repeated 25 to 30 times to produce approximately 10¹² copies of the sequence.

of drug resistance can be determined. In Figure 6 the comparative survival of myeloma cells and normal haemopoietic cells from the same patient is shown. An example is given of a highly sensitive and a highly resistant cell. The theoretical importance of such studies cannot be overemphasized, because with our increasing awareness of the molecules that are responsible for controlling cells, this technique will give an opportunity to study regulatory mechanisms.

Regulatory mechanisms imply circulating molecules which in turn must have receptors if they are to function. There has been increasing interest in epidermal growth factor receptor (EGFr) in human breast cancer. Many prognostic variables have been described for breast cancer – for example, age at menarche or parity, size of tumour and presence of lymph node metastases, histological grade⁸ and more recently the beneficial import of the presence of oestrogen (ER) and progesterone (PR) receptors. EGFr on human breast cells was described by Sainsbury *et al.*⁹ They also pointed out that it occurred in an inverse ratio to the oestrogen receptor.¹⁰ The presence of EGFr is correlated with poor survival in breast cancer.¹¹ From these studies, and studies on the effect of growth in tissue culture, it would seem that EGFr plays an important part in regulating the growth of human breast cancer.¹² It has been suggested, because of a possible role in autoendocrine cell stimulation, that its presence identifies a group of more aggressive tumours with a higher metastatic potential. Battaglia *et al.*¹³ have investigated the numbers of EGFr, ER and PR in primary breast cancer and axillary lymph node metastases. They found that EGFr receptors were more common in the metastatic glands than in the primary tumour, and that there was a reverse relationship between the presence of EGFr and PR and ER, which was highly significant. The adverse role of EGFr is further strengthened by the latest report from

Figure 6 Comparative survival of myeloma and normal haemopoietic cells in a patient treated with melphalan. The effect of melphalan on the survival of myeloma (●) and GM-CFUc (○) cells *in vitro*. (Cells were exposed to melphalan in PBSA for 1 h at 37°C before plating.) Reproduced with permission from B. Miller.

Sainsbury's group,¹⁴ who have demonstrated, in a series of 264 consecutively excised breast carcinomas, a close relationship between the presence of EGFr and histological high risk tumours.

Haemopoietic growth factors

Continuing the theme of regulatory factors and receptors, the ability to produce peptide regulatory factors such as erythropoietin, granulocyte macrophage colony stimulating factor (GMCSF) and granulocyte stimulating factor (GCSF) must be considered to be one of the most exciting recent developments.^{15,16} The clinical role of erythropoietin has been explored most in the area of primary deficiency of this factor in patients with end-stage renal disease, so that further discussion would not be relevant, though the possibility that it, too, could be of use in initial stages of marrow transplantation is still to be explored.

The remarkable fact about GMCSF and GCSF is that they cause a many-fold increase in functional granulocytes without evidently depleting the stem cell pool. They can do this in humans in effective doses without causing toxicity. The increased levels of granulocytes are sustained over a number of days, but return to normal when the factor is no longer given (Figure 7). The application of these factors to clinical situations¹⁷ in which cytotoxic drugs have depressed the neutrophil count is still being investigated, but good clinical studies¹⁸ showed a reduction in the infection rate in patients with small cell lung cancer treated with chemotherapy when given GCSF using within-patient randomization. Using a similar within-patient study, in which the first course of therapy was accompanied by GMCSF and the second course was not, Antman *et al.*¹⁹ showed in 16 patients with inoperable metastatic sarcomas a

significant difference in the duration of neutropenia. The duration of neutropenia was only 3.5 days after the first cycle, whereas it was 7.4 days after cycle 2 ($P < 0.01$). However, the number of febrile episodes in the two cycles was the same, but as fever may have been related to the GMCSF, it is difficult to know whether infection rates were lower in the GMCSF cycles. Other studies in urogenital tumours have been reported.²⁰

High-dose chemotherapy accompanied by total body irradiation has been used in the experimental treatment of such devastating diseases as end-stage breast cancer and disseminated malignant melanoma. The profound neutropenia seen following such therapy makes an ideal situation for testing the value of these factors. Brandt *et al.*²¹ reported the use of GMCSF in 19 patients with breast cancer and melanoma, and compared the duration of neutropenia to an historical control group of 24 patients matched for age, diagnosis and treatment. Total leucocyte and granulocyte recovery were accelerated in the patients given GMCSF compared to the control group. Toxicity at lower level of GMCSF was very tolerable, but at the maximum doses used, generalized oedema, pleural effusions and hypotension developed. Two of the 19 patients died of infection during admission, one due to bacteria and the other fungal, but although there had been no in-patient deaths from infection in the controls, there was evidence of bacteraemia in only 3/19 treated with GMCSF compared to 8/23 in the controls. The authors do suggest that there was less hepatotoxicity and renal toxicity in the treated patients, and speculate that this was due to a lower incidence of bacterial infection. The fact that sudden death from overwhelming bacterial infection can occur in patients treated with haemopoietic growth factors raises the worrying possibility that these patients might have an impaired ability to respond to infections following stimulation by these substances.

It is very probable that the efficacy of these factors may not only be due to the rapid recovery of numbers of polymorphs (PMN). Kaplan *et al.*²² studied the PMNs from 10 patients with cancer treated with GMCSF, and were able to show that there was enhanced superoxide release in the PMN from the GMCSF patients treated with lower doses. Superoxide production is related to the effectiveness of bacterial killing, so that this may indicate a further reason for not escalating the dose of these factors to too high a level. Metcalf,²³ in a recent review, points out that both GMCSF and GCSF are probably produced as a response to natural infections, and further studies will be needed to study the combinations of these factors and the addition of other regulators such as interleukin-1. The role of these interesting products in cancer therapy is now a focus of intense activity.

Figure 7 Increased level of polymorphonuclear cells following the injection of GCSF compared with the level of neutrophils in untreated patients. Note initial rise followed by a short nadir.

There is now a requirement for larger randomized studies to confirm that the restoration of the PMN count is correlated with a reduction in the number of febrile episodes due to bacteria, and long-term studies will also be needed to confirm the absence of any deleterious effect on the white cell progenitors. These factors do hold open the process of reducing the dose-limiting effects of marrow suppression by cytotoxic drugs, which in theory, at least, should enhance the efficacy of intensive chemotherapy.

Mechanisms of drug resistance

Drug resistance has been a major obstacle to curing cancer with chemotherapy. Even when many drugs are given, either together or sequentially, drug resistance soon arises. In leukaemia and solid tumours that are curable, such as lymphoma or testicular cancer, malignancies that are highly responsive initially may relapse, and when they do so, the tumour cells are refractory to many drugs. Other tumours show resistance *de novo* – for example, adenocarcinoma of the colon and malignant melanoma. Both forms of resistance to drugs may involve more than one drug, with so-called 'cross resistance' and this is now termed 'multidrug resistance' or MDR. The study of cell lines which have developed MDR has enabled the mechanisms of such resistance to be studied, and the mechanisms are now known at a molecular level.

Although there are many ways in which drug resistance can develop, MDR appears to be effected by an increased ability of the cell to excrete the toxic drug by means of an energy-dependent efflux system (Figure 8). Circumstantial evidence suggests that membrane glycoproteins of 170 kDa are responsible, as these are to be found in increased quantities in drug resistant cells.²⁴ MDR genes encoding for these proteins are found to be amplified in resistant cell lines and reduced in sensitive cells.

One of the most telling pieces of evidence is that when genes for the P-glycoprotein are transfected into the DNA of cell lines, these lines then show enhanced expression of P-glycoprotein and increased drug resistance. While there may be other mechanisms for drug resistance, the P-glycoprotein at least offers a practical target for manipulation and possible reversal of clinical chemotherapeutic resistance, by such agents as calcium channel blockers (verapamil),²⁵ calmodulin inhibitors²⁶ and isoprenoids.²⁷ Mechanisms by which these drugs act are reviewed by Morrow and Cowan.²⁸

Recently, a single and practical means of studying the over-expression of P-glycoprotein using a monoclonal antibody has been described.²⁹ The development of increased P-glycoprotein in the

Figure 8 Diagrammatic representation of the cell wall with a molecule of P-glycoprotein engaged in excreting cyto-toxic drugs from the cytosol below into the extracellular fluid above.

cells of patients with myeloblastic leukaemia who become progressively resistant to cytotoxic drugs can be demonstrated by a monoclonal antibody C219, which is now commercially available. It has been evaluated using immuno-blotting techniques, immunocytology,²⁹ or flow cytometry.³⁰ Immunohistochemical staining of cells from patients with myeloma, lymphoma and breast cancer have shown excellent correlation of high P glycoprotein with resistance to doxorubicin.³¹

Chemotherapeutic agents which are known to be affected by multiple drug resistance are largely derived from natural products, and include vincristine, vinblastine, doxorubicin, daunorubicin, mitoxantrone, mithromycin, actinomycin D, etoposide, tenoposide and mitomycin. These drugs have been shown to be effective in malignancies such as leukaemia, ovarian cancer, breast cancer, lymphoma and myeloma. These are also cancers in which it has been shown that P-glycoprotein is associated with MDR. Consequently it would appear likely that measurements of the level

of P-glycoprotein would be helpful in planning rational therapy for relapsed patients.

A new chemotherapeutic agent for haematological malignancy

After decades of new therapeutic agents, the number of successful compounds entering clinical trials has tended to plateau in the last few years. The arrival of fludarabine, a relatively non-toxic but effective drug for a common malignant haematological condition which is difficult to treat, chronic lymphatic leukaemia, is welcome. A Phase I study of fludarabine phosphate,³² has outlined its use by a loading dose and continuous infusion schedule. An earlier drug, vidarabine, was synthesized as a potential anticancer agent, but was limited by its poor solubility and rapid deamination to a less active compounds. A fluorinated derivative of vidarabine was synthesized and was found to be resistant to deamination. The compound was 9-bd arabinofuranosyl 2-fluoroadenine (2f-ara-a), which was even more soluble when phosphorylated. Activity was noted against a wide range of tumours in early studies, and the compound was also found to be relatively non-toxic.^{33,34}

Leiby *et al.*³² used a 20 mg/m² loading dose followed by a continuous intravenous infusion of 30 mg/m² in 24 hours for a total duration of 48 hours. Courses were given every 21 days. The loading dose was not increased, but the infusion rate was increased to 45 mg/m² in 24 hours. At the latter dose leucopenia occurred. Other toxic effects were mild nausea, fatigue and diarrhoea. One patient with non-Hodgkin lymphoma became stable during the course of therapy.

Keating *et al.*³⁵ reported the use of fludarabine in chronic lymphatic leukaemia at a dose of 25–30 mg a day for 5 days every 3–4 weeks by intravenous infusion. Seventy five patients with chronic lymphatic leukaemia were entered into the study, of whom 47 (63%) were Rai stage III or IV. Stages, O, I and II were included if they had progressive disease. Forty two (56%) of the 75 previously treated patients achieved a complete or partial response. Ten (13%) showed a fall of the blood count to $<4 \times 10^9/l$ with less than 30% lymphocytes in the marrow and liver and spleen becoming impalpable. They had shotty residual lymphadenopathy. A total of 32 (43%) had a less complete response. Mean overall survival for the whole group was 16 months, with those achieving complete or good partial remission having median survivals in excess of 35 months. The major toxicity was infection. Thirty one episodes of pneumonia and 16 minor infections occurred. Neuropathy, myopathy, muscular weakness were all uncommon. Tumour lysis was noted in 3 patients. Further

studies of untreated patients have shown a good partial response or complete response in over 70% of those treated, and the response has been rapid. Severe neuropathy, which has been such a feature of the patients treated with the higher doses in acute leukaemia, has not been seen in these series.

Another way to increase the efficacy of fludarabine was investigated, with prednisolone being used in combination.³⁶ A total of 50 patients with chronic lymphatic leukaemia were given fludarabine in a dose of 30 mg/m²/day for 5 days by intravenous infusion over 30 minutes. For the duration of the 5-day course, prednisolone was given orally in a dose of 30 mg/m². All patients had been pretreated. Using the same criteria for response as given above, 7 (14%) achieved a complete response, and 23 (46%) a partial response. A rapid fall in the peripheral lymphocyte count was seen. It was of interest that the most important determinant for prognosis in these 50 patients was the level of the serum albumin: those above 40 g/l showing a significantly better survival than those with less than 40 g/l. In this study, comparing the results with those of their previous studies, prednisolone did not appear to improve either response or survival, but since many of their previously treated patients had already had steroid therapy, this finding may not be true in previously untreated patients. Further studies of fludarabine as a single drug and in combination with steroids in patients with untreated chronic lymphatic leukaemia will be of great interest.

What hope for solid tumours?

While it has been possible to be enthusiastic about some aspects of the management of haematological malignancy during recent years, when one turns to the management of solid tumours, particularly those that are common, the outlook has remained distinctly gloomy. Even when modest but statistically significant improvement in disease-free survival and overall survival has been demonstrated (for example in a multidrug adjuvant therapy trial in carcinoma of the colon³⁷) these responses have been regarded with attitudes that vary from 'sceptical' to 'hostile',³⁸ and have had little impact on clinical practice. It seems that failure is almost expected in these cancers, and it is important therefore for new and encouraging results in colon cancer treatment to become more widely known. It is 20 years since 5-fluorouracil (5-FU) was established as the most useful drug for advanced colorectal cancer. When given intravenously response rates of 15–20% were seen. Unfortunately these responses were relatively short in duration, and the longest were only 6 months. Many attempts were made, by adding another drug

such as cisplatin, vincristine, and methyl CCNU to 5-FU, to improve response rates, but these have generally failed.

5-FU acts by inhibiting DNA synthesis of cells in S phase. Colon cancer is a slowly proliferating tumour, so that opportunity for 5-FU to be active is relatively infrequent, particularly if it is to be present in adequate concentrations for only a short time. Constant infusion of 5-FU has been shown to be about twice as effective in producing responses as bolus drug in randomized studies.³⁹

5-FU acts by inhibiting thymidylate synthetase. Folinic acid administration produces an excess of intracellular reduced folate, which inhibits the enzyme and adds considerably to the efficacy of 5-FU. Thirty seven patients with advanced measurable colorectal cancer were treated with high-dose folinic acid (200 mg/m² over 2 hours by infusion) followed by 5-FU as a bolus (300 mg/m²), and then a continuous infusion on the next two days.⁴⁰ This was repeated 14 days later, and then 14 days after that. Escalation of the dose of 5-FU was carried out if there were no serious side effects on the white cell count. The successive escalation of 5-FU increased the dose to 400, then 500 mg/m². The use of implantable infusion devices and portable pumps made the approach tolerable. An overall response rate of 54% was seen, which in some sub-groups of patients rose to as high as 61%. This probably represents the best response rate ever reported with 5-FU, which should obviously be studied in a randomized trial comparing 5-FU alone, as the authors suggest. The question is whether this response rate helps the patient. In these small series, the median survival was 18 months, with responders having a median duration of survival of 21 months, approximately double that seen previously. Again, although it seems likely to be a result of the therapy, a randomized study is still necessary to establish this. Response rates and survival such as these must be confirmed, and if they are as they seem, this would suggest that a nihilistic approach is no longer justified.

Furthermore, with objective responses beginning to exceed the 50% level, it is likely that the role of 5-FU and folinic acid in an adjuvant setting after surgical resection should be reconsidered.

Intensive therapy supported by bone marrow transplantation

The use of increasingly intensive chemotherapy, with or without irradiation, supported by either allogeneic or autologous bone marrow transplantation, is being explored in a variety of malignancies. The underlying principle that there is a dose-related response to chemotherapy has driven this expensive and labour-intensive form of treat-

ment. Allogeneic bone marrow transplantation has been the technique most used in haematological malignancies, particularly acute myeloblastic and lymphoblastic leukaemia in both adults and children. It would be expected that by now a clear statement could be made about the efficacy of this procedure, and that principles could be deduced that might help with its application to other tumours.

Allogeneic bone marrow transplantation as a method of support is not available to more than approximately 35% of the population, because of the absence of a suitable HLA-matched donor. Higher figures reported by some American centres reflect referral bias. Secondly, graft-versus-host disease has not been eliminated, and still accounts for substantial morbidity and mortality, especially in older patients. Finally, in the young, dramatic growth failures, endocrinopathies and gonadal failure, respiratory and renal complications have been difficult to counter. When faced with a lethal disease, these complications might be accepted if it could be shown that the alternatives are less effective.

The evidence is difficult to evaluate at the moment. If long-term event-free survival is taken to equate with cure, there is no doubt that this may be seen in children and adults with both forms of leukaemia treated either with intensive chemotherapy or with intensive therapy supported by an allogeneic transplant. The difficulty is that comparisons are impossible because these groups of patients subjected to one modality of therapy or another are not directly comparable. It is not possible to compare adults with acute myeloblastic leukaemia who are treated with intensive chemotherapy with those adults treated in the standard and carefully controlled conditions which exist in transplant centres, since those treated away from the centres are likely to show greater variation in age, intensity of induction therapy, and supportive care. Even the most committed transplant enthusiast has to admit that patients are selected. They have already been shown to have responsive disease, they are in a good age group, and their general clinical condition is usually better than that of patients who are entered on alternative intensive chemotherapy programmes. Meyer⁴¹ in a thoughtful review compares the best data available on the alternative approaches to the management of adults with acute myeloblastic and acute lymphoblastic leukaemia. This very complex data is presented in Table I.

The overall impression given by Table I is that there is no great difference between those patients given treatment supported by bone marrow transplantation, and those subjected to intensive therapy. The three within-institutions comparisons⁴²⁻⁴⁴ of intensive therapy with allogeneic bone marrow

Table I Results from alternative approaches to the management of adults with leukaemia

Source	Median follow up (months)	Median age	No. of patients	No. of relapsed patients (%)	Probability of 3-year disease-free survival
<i>Acute myeloblastic leukaemia</i>					
Allogeneic B.M.T. Seattle	> 60	~24.5	231	33 (14)	~ 51
International Bone Marrow Transplant Registry	N/A	25.0	704	N/A (20)	~ 52
Intensive chemotherapy St Bartholomew's Hospital	42	44.0	118	73 (60)	~ 38
Wolff <i>et al.</i>	29	34.0	55	19 (35)	~ 55
<i>Acute lymphoblastic leukaemia</i>					
Allogeneic B.M.T., Seattle	47	22.0	34	8 (24)	38
Westminster Hospital	43	26.0	23	7 (30)	44
Intensive chemotherapy West Germany	> 60	25.0	272	156 (57)	
Memorial Hospital	> 60	27.0	167	92 (57)	
				After Mayer (<i>J Clin Oncol</i> 1988, 6: 1532–1536)	

N/A = not available

transplantation would suggest that the probability of 4-year disease-free survival is greater in the transplant group than the chemotherapy group, but it has to be remembered that treatment in the transplant group was rigorously controlled, whilst the chemotherapy given outside the unit would undoubtedly be less well controlled, and indeed, in one of the series, the patients who were transplanted included much younger patients than the group who were not transplanted, which therefore favoured the transplant group. Attempts to remedy this unsatisfactory position are now ongoing, and although large, carefully-controlled studies are not mature enough to give us an answer, there is a suggestion^{45,46} that this information will soon be available. Preliminary results in adult myeloblastic leukaemia suggest that patients who follow induction by intensive therapy together with total body irradiation and autologous bone marrow transplantation do no better than those treated with high-dose cytosine arabinoside as consolidation. Furthermore, if induction therapy had been further intensified, the intensive chemotherapy group might have done significantly better.

Where there is no clear benefit in terms of outcome, then a new look at cost/benefit ratio is important, and nowhere more so than in children. In a well-argued editorial Pinkel⁴⁷ considered that allogeneic bone marrow transplantation in children with acute leukaemia was 'a practice whose time has gone'. Whatever the outcome of the debate, it is certain that the next few years will see a greater effort to define the role of standard intensive therapy and intensive therapy supported by both allogeneic and autologous bone marrow transplantation.

Consequences of intensive therapy

It is probably appropriate to follow discussion of one of the most intensive forms of therapy known, i.e. the use of cytotoxic therapy together with irradiation, with a short discussion of long-term sequelae of therapy of malignant disease. A major concern for the physician, the parent and the child, is the consequences of cranial prophylaxis against the recurrence of leukaemia in the central nervous system. More than 50% of children with acute lymphoblastic leukaemia will relapse in the central nervous system if cranial prophylaxis is not given. The first effective procedure was cranial irradiation with 2400 cGy and 5 doses of intrathecal methotrexate.⁴⁸ Subsequently it has been shown that cytosine arabinoside, methotrexate and hydrocortisone given intrathecally, or high-dose methotrexate given intravenously with sequential intrathecal methotrexate, are effective prophylactic therapies.

It was first thought that cranial irradiation with intrathecal methotrexate was probably unlikely to give rise to decreased intellectual function.⁴⁹ However, subsequent studies identified impairment of intellectual function.^{50–53} Earlier studies are difficult to assess, because they contain children from a wide variety of prophylactic central nervous system programmes. They often included children who had had more than one course of therapy, and took little account of the socio-economic background of the child.⁵⁴ Indeed, in some studies there seemed to be no difference in intellectual function between children having prophylactic therapy for central nervous system leukaemia, and children who had been hospitalized for cancer therapy

which did not involve central nervous irradiation.⁵⁵ It is still not clear whether prophylaxis does significantly affect the intellect of the child treated with one course of cranial prophylaxis, while there is considerable evidence that this is probably true in the child under the age of three. It has been accepted that intellectual deficit is seen more frequently in patients treated with cranial irradiation and intrathecal methotrexate rather than intrathecal methotrexate alone, or high-dose methotrexate alone. Carefully controlled studies have been reported,⁵⁶ in which memory has been examined in children randomized to receive either cranial irradiation 1800 cGy with intrathecal methotrexate, or intravenous methotrexate together with high-dose methotrexate. The study groups were carefully selected so that they contained no children with central nervous system disease at the start of the study. They were all studied on the same protocol, and were matched for social background. The children were examined some 5 years after they had had cranial prophylaxis. No difference was detected between the groups in full scale IQ or a battery of 16 memory tests. Furthermore, there appeared to be no increased adverse effects with younger onset of disease, in contradistinction to most series. They chose to examine memory further, because this is a most sensitive indication of learning aptitude which depends less heavily on previously-acquired knowledge than IQs. They again could detect no difference between the two groups. However, when the groups of leukaemic children were combined and compared to their normal peers, there appeared to be a significant decrease in spatiovisual memory and four scales of verbal memory. The authors, however, concluded that adequate academic, social, behavioural and intellectual development is possible in long-term survivors of acute lymphoblastic leukaemia 5 years after central nervous system prophylaxis, whether or not they received 1800 cGy cranial irradiation as prophylaxis.

Studies using computed tomographic (CT) scanning of the brain have previously noted a number of abnormalities in children who have cranial irradiation as prophylaxis.⁵⁷ These abnormalities have been correlated with impairment of cerebration, such as memory disturbance;⁵⁸ other studies in children who have had cranial irradiation and intrathecal methotrexate as prophylaxis showed lower frequencies of CT scan abnormalities, and in one series⁵⁹ no abnormality was noted in 27/27 children. In other studies done recently, the incidence has also been very low.

The changes reported have mostly affected the white matter, and as magnetic resonance imaging has been shown to be superior to CT scanning in detecting white matter changes, the findings of

magnetic resonance imaging (MRI) in central nervous system (CNS) leukaemia are of great interest.⁶⁰ MRI was used to investigate ten survivors of acute lymphoblastic leukaemia who received intrathecal methotrexate and either 1800 or 2400 cGy to the brain. At the same time they combined neuropsychological testing. MRI was normal in 9/10 patients. White matter tracts were normal, although in their testing some cognitive functions were impaired. However, the study of the cognitive functions could be criticized in that it compared results in these children with national standards, and not with sibs or children who had experienced prolonged illness. It would seem increasingly evident that parents and the children who have received these courses of standard prophylactic CNS treatment can be reassured about the low incidence of any objective brain abnormality, and that intellectual, social or behavioural development will allow a normal life.

Imaging disease

The breathtaking speed of development of CT and MRI in recent years, and the clarity and accuracy of the images produced have perhaps made us tend to forget that these techniques do produce a world of shadows and with CT, at least, as far as cancer tissue is concerned, little hint as to whether the tumour is viable or not.

Techniques that have been available much longer may still have a useful role to play. Gallium 67 (GA) scanning has been the subject of controversy in the past insofar as the management of lymphoma is concerned. GA is taken up by active Hodgkin's tissue, and a sensitivity of detection of 66% is recorded for lymphoma in the chest using GA and plain imaging.⁶¹ These authors claim that this may be increased to 96% by the use of simple photon emission computed tomography.

Anecdotes of cases of human disease in which GA is a better indicator of tumour viability than the presence of a tumour mass on scanning may be found in the literature.^{62,63} It is many people's experience in treating Hodgkin's disease with mediastinal lymphadenopathy that when the disease appears otherwise to have responded completely, either plain chest X-ray or CT scanning reveals a persistent mediastinal mass. What does this represent, and how in particular can one determine, in an area which is not easily accessible for biopsy, what is actually present?

Israel *et al.*⁶⁴ have used GA to monitor lymphoma response in 25 patients with GA avid tumours, and compared body CT, chest radiographs and physical examination of tumour infiltrated with peripheral lymph nodes. GA imaging was negative in 95% (20/21) of the patients who were considered clinically to be in remission and in whom treatment

was stopped. The disease did not recur in 15 patients who were followed between 12 and 26 months. In the other 6 patients, disease recurred 3–12 months after stopping treatment, and on each recurrence the GA became positive at the time of discovery of disease. Based on CT examinations, only 11 patients would have been considered to have achieved remission. The specificity for CT was 57%. Similar specificities for chest X-ray of 55% were noted. Thus these studies confirm the views of many clinicians that many mediastinal masses detected on CT or chest X-ray after effective therapy are in fact fibrotic and necrotic tissue only, and this study⁶⁴ has been one of the first systematic studies to show this and to demonstrate that GA imaging is useful in monitoring lymphoma response to treatment.

In a recent study,⁶⁵ in which gallium scanning was used to investigate residual mediastinal disease in children treated for Hodgkin's disease, 37/46 (80%) were positive, and 9/46 (20%) were negative in the mediastinum. At the completion of 8 courses of MOPP ABVD prior to low-dose total nodal irradiation, 23/37 patients with initially positive gallium scans had them repeated as part of their clinical re-staging evaluation. Twenty of the twenty three (87%) had had a conversion of an initial positive scan to a negative scan, whilst 3 remained positive. In 13 of the 20 patients whose scans became negative, the result was consistent with a clinical complete remission, and compatible with the normalization of the CT scan of the chest. In 7, although the gallium scan was negative, the CT scan continued to demonstrate residual mediastinal disease. All 7 of these patients had a biopsy performed, and this showed no evidence of Hodgkin's disease. They conclude that the gallium scan was a more accurate predictor of residual mediastinal disease than the CT scan in the subset of patients who convert from gallium positivity to negativity. In the 20 patients whose scan became negative, 2 have had recurrence of their Hodgkin's disease. In the 3 patients whose gallium scan remained positive, and who had a mediastinal mass on CT scan, one had a biopsy that was positive for disease. The other 2 have either relapsed or died. Again, it would appear that gallium scanning is of considerable help in their therapeutic dilemma.

Supportive care

Antibiotic therapy

Infection is among the chief causes of morbidity and mortality with intensive chemotherapeutic regimens. The development of antibiotics which are effective against Gram-negative organisms, resistant Gram-positive organisms, and some

fungi, has enabled deaths from infection with these organisms to be considerably reduced. Viral infection with herpes simplex and zoster viruses has also been controllable with the successive introduction of vidarabine and, more recently, acyclovir. Intravenous administration of acyclovir, which is a selective and specific inhibitor of viral replication, has produced many advances in the control of herpes simplex and varicella.^{66,67} Cytomegalovirus is, however, one pathogenic virus that has remained resistant to therapy until recently.

Cytomegalovirus is a herpes virus which becomes latent after primary infection, but can be provoked into activity during periods of immunosuppression consequent on bone marrow transplantation, intensive chemotherapy for a long period, or from acquired immunodeficiency syndrome (AIDS). In these clinical settings, chorioretinitis, pneumonitis (often associated with *Pneumocystis carinii*), and infection of the gastrointestinal tract can occur. Sight may be impaired, and blindness result from cytomegalovirus infection of the retina. In cytomegalovirus pneumonitis, which proceeds to hypoxia, recovery is rare.^{68,69} Cytomegalovirus can affect the whole of the gastrointestinal tract, causing nausea, vomiting, diarrhoea and abdominal pain, and lesions can be seen in the wall of the oesophagus and colon. Ganciclovir [9-(2 hydroxy-1-hydroxymethyl) ethoxymethyl guanine], formerly known as DHPG, is cyclic analogue of thymidine active against all herpes viruses. It is structurally similar to acyclovir, differing only in the addition of a terminal hydroxymethyl group (Figure 9). Ganciclovir is converted first to the triphosphate, which is a potent inhibitor of viral DNA polymerase. The drug is converted very rapidly to the monophosphate by thymidine kinase, and then to the triphosphate, and this occurs particularly in cytomegalovirus infected cells, for reasons which are presently unknown. Whatever the reason, the effect is specific because normal cells are not affected. The drug is excreted virtually unchanged in the urine. The commonest toxic effect is neutropenia, occurring in 40% of patients treated. Thrombocytopenia occurs in about 20%, and various manifestations of central nervous system disorders occur in about 9% of patients, and these include headaches, dizziness, anxiety states, drowsiness, confusion and hallucination. In a very small number of patients, a wide variety of symptoms have occurred in the alimentary, cardiovascular and respiratory systems, making it a drug which should only be given with a degree of caution to probable or confirmed cases of the disease. Critical reviews are now available.^{70,71} There would seem to be good evidence that sight may be saved by its use in cytomegalovirus infection of the retina, and improvements in 30/41 patients with cytomega-

Figure 9 Structural formula of ganciclovir and its relation to the other antiviral agents such as acyclovir.

lovirus oesophagitis, proctitis and enteritis are recorded with ganciclovir.⁷² Responses in the lethal pneumonia remain largely anecdotal, because it occurs so frequently in association with *Pneumocystis carinii* and it is difficult to isolate cytomegalovirus from the lung and to be certain of the diagnosis. There is a need for carefully constructed controlled trials of therapy. Nevertheless, this drug

and another potent inhibitor of herpes virus DNA polymerase, trisodium phosphonoformate (Foscarnet) are opening the door to the control of this infective agent.

Treatment of hypercalcaemia

Hypercalcaemia is a complication of malignant disease which is now undergoing a significant change in management. A recent review⁷³ noted that malignancy accounts for about half the cases of hypercalcaemia, that 5% of patients with malignant disease develop it at some stage of their illness, and that carcinoma of the breast, carcinoma of the lung and myeloma are frequently complicated and may present with the usual cluster of non-specific symptoms such as fatigue, anorexia, constipation, polydipsia, nausea, vomiting and loss of strength. The multitude of therapies used so far have not always been effective, but more recently reliable agents have become available. These are the biphosphonates, etidronate disodium and clodronate, the former generally available and the latter on a clinical trial certificate. These agents inhibit bone resorption and prevent osteolysis due to a direct inhibition of the action of osteoclasts. Table II shows the features of etidronate and clodronate. Neither is particularly well absorbed orally, but clodronate, if given in sufficient quantity, can be effective by mouth and has no unpleasant side effects. In order to obtain the most rapid effect, etidronate should be given intravenously over a period longer than 2 hours. Pain relief can be obtained with both etidronate and clodronate, and prompt and sustained control of hypercalcaemia has been noted.

Etidronate used intravenously in a daily dose of 7.5 mg/kg body weight for 5 days was compared with a placebo in 20 patients with malignant disease complicated by hypercalcaemia.⁷³ The etidronate was given by infusion intravenously for 2.5–3 hours. All patients were given 3000 ml of intravenous saline daily and 40 mg frusemide daily. Eighteen patients completed the study. Eleven of the 12 patients (92%) treated with etidronate attained normo-calcaemia compared to 2/6 (33%) in the placebo group. This was significant ($P = 0.05$). Renal calcium excretion decreased in the etidronate group, but not in the placebo group. Heath⁷⁴ uses this regimen together with potassium supplements, and notes that it is nearly always effective. This is better than the results achieved with steroids, which are much less effective, frequently delayed in action, and give rise to side effects including the problems secondary to fluid retention when given to elderly patients. Disodium etidronate is not effective orally and almost certainly does not replace calcium lost from the bones. Clodronate was first introduced in the early

Table II Biphosphonate therapy in myeloma – etidronate and clodronate

	Admin.	Effect of pain	Hypercalcaemia	Bone resorption	Bone resorption
Etidronate	Intravenous	+	+	+	–
			(prompt)		
Clodronate	Oral and intravenous	++	+	+	++

'80s,^{75,76} and then withdrawn because of fears about carcinogenicity. Clodronate would not only be effective orally, but might produce restoration of calcium to the skeleton. Clodronate is now being re-evaluated following the reappraisal of the evidence for carcinogenicity, which could not be substantiated, and it may well in the future be the treatment of choice for malignant hypercalcaemia.

Conclusion

Cancer still presents an intractable problem. The bystander must feel, quite justifiably at times, that the vast investment that has been made in the scientific understanding of the cancer cell, and the various modalities which can kill it, have given little

practical reward so far. This might be a reason for writing an annual report, for reading even some of the references relating to the molecular biology of the cancer cell cannot but give an appreciation of how detailed and accurate our knowledge of this subject is becoming. It is inconceivable that the next few years will not show some major advance in our ability to manipulate the intimate control of these cells. Meanwhile, some cancers are being cured, and many palliated. This article has recorded some examples where, within the last year, there have been significant advances.

Acknowledgement

I would like to thank Barbara Gibbons for the cytopathology illustrations.

References

Chromosomal abnormalities

1. Vogelstein, B., Fearon, E.R., Hamilton, S.R. *et al.* Genetic alterations during colorectal tumour development. *N Engl J Med* 1988, **319**: 525–532.
2. Bodmer, W.F., Bailey, C.J., Bodmer, J. *et al.* Localisation of the gene for familial adenomatous polyposis on chromosome 5. *Nature* 1987, **328**: 614–616.
3. Leppert, M., Dodds, M., Scambler, P. *et al.* The gene for familial polyposis coli maps to the long arm of chromosome 5. *Science* 1987, **238**: 1411–1413.
4. Kurzrock, R., Gutterman, J.V., Talpaz, M. The molecular genetics of Philadelphia chromosome positive leukaemia. *N Engl J Med* 1987, **319**: 990–998.
5. Weiss, L.M., Warnke, R.A., Sklar, J. *et al.* Molecular analysis of the t(14;18) chromosome translocation in malignant lymphomas. *N Engl J Med* 1987, **317**: 1185–1189.
6. Lee, M.-S., Chang, K.S., Cabanillas, F. *et al.* Detection of minimal residual cells carrying the t(14;18) by DNA sequence amplification. *Science* 1987, **237**: 175–178.

Cell culture

7. Millar, B.C., Bell, J.B.G., Lakhani, A., *et al.* A simple method for culturing myeloma cells from human bone marrow aspirates and peripheral blood in vitro. *Br J Haematol* 1988, **69**: 197–203.
8. Bloom, H.J.G. & Richardson, W.W. Histological grading and prognosis in breast cancer. *Br J Cancer* 1957, **13**: 359–
9. Sainsbury, J.R.C., Sherbet, G.U., Farndon, J.R. *et al.* Epidermal growth factors are present on human breast cancers. *Br J Surg* 1985, **73**: 186–194.
10. Sainsbury, J.R.C., Farndon, J.R., Sherbet, G.U. Epidermal growth factor receptors and oestrogen receptors in human breast cancers. *Lancet* 1985, **i**: 364.

11. Sainsbury, J.R.C., Needham, G.K., Malcolm, A. *et al.* Epidermal growth factor receptor status as predictor of early recurrence of and death from breast cancer. *Lancet* 1987, **i**: 1398.
12. Imai, Y., Leung, C.K.H., Friesen, H.G., *et al.* Epidermal growth factor receptors and effect of epidermal growth factor on growth of human breast cancer cells in long term tissue culture. *Cancer Res* 1982, **42**: 4394–4398.
13. Battaglia, F., Scambia, G., Rossi, S. *et al.* Epidermal growth factor receptor in human breast cancer: correlation with steroid hormone receptors and axillary lymph node involvement. *Eur J Cancer Clin Oncol* 1988, **24**: 1685–1690.
14. Sainsbury, J.R.C., Nicholson, S., Angus, B. *et al.* Epidermal growth factor receptor status of histological sub-types of breast cancer. *Br J Cancer* 1988, **58**: 458–460.

Haemopoietic growth factor

15. Donahue, R.E., Wang, E.A., Stone, D.K. *et al.* Stimulation of haemopoiesis in primates by continuous infusion of recombinant human GM-CSF. *Nature* 1986, **321**: 872–875.
16. Welte, K., Bonilla, M.A., Gillio, A. *et al.* In vivo effects of recombinant human G-CSF in therapy induced neutropenia in primates. *Exp Haematol* 1987, **15**: 72 (abstract).
17. Clark, S.C. & Kamen, R. The human haemopoietic colony stimulating factors. *Science* 1987, **236**: 1229–.
18. Bronchud, M.H., Scarffe, J.H., Thatcher, N. *et al.* Phase I/II study of recombinant human granulocyte stimulating factor in patients receiving intensive chemotherapy for small cell cancer. *Br J Cancer* 1987, **56**: 809–813.
19. Antman, K.S., Griffin, J.D., Elias, A. *et al.* Effect of recombinant human granulocyte-macrophage colony-stimulating factor on chemotherapy-induced myelosuppression. *N Engl J Med* 1988, **319**: 593–598.

20. Gabrilove, J.L., Jakubowski, A., Scher, H. *et al.* Effect of granulocyte colony-stimulating factor on neutropenia and associated morbidity due to chemotherapy for transitional cell carcinoma of the urothelium. *N Engl J Med* 1988, **318**: 1414–1422.
21. Brandt, S.J., Peter, W.P., Atwater, S.K. *et al.* Effect of recombinant human granulocyte-macrophage colony-stimulating factor on haemopoietic reconstitution after high-dose chemotherapy and autologous bone marrow transplantation. *N Engl J Med* 1988, **318**: 869–876.
22. Kaplan, S.S., Basford, R.E., Wing, E.J. *et al.* The effect of recombinant human granulocyte-macrophage colony-stimulating factor on neutrophil activation in patients with refractory carcinoma. *Blood* 1989, **73**: 636–638.
23. Metcalf, D. Haemopoietic growth factors. 2. Clinical applications. *Lancet* 1989, **i**: 885–886.

Mechanisms of drug resistance

24. Juliano, R.L. & Ling, V. A surface glycoprotein modulating drug permeability in chinese hamster ovary cell mutants. *Biochem Biophys Acta* 1976, **455**: 152–162.
25. Tsuruo, T., Iida, H., Tsukagoshi, S. *et al.* Overcoming of vincristine resistance in P388 leukaemia *in vivo* and *in vitro* through enhancing cytotoxicity of vincristine and vinblastine by verapamil. *Cancer Res* 1981, **41**: 1967–1972.
26. Akiyama, S., Shiraishi, N., Kurajomi, Y. *et al.* Circumvention of multidrug resistance in human cancer cells by thioridazine, trifluoroperazine and chlorpromazine. *JNCI* 1986, **76**: 839–844.
27. Nakagawa, M., Akiyama, S., Yamaguchi, S. *et al.* Reversal of multidrug resistance by synthetic isoprenoids in the KB human carcinoma line. *Cancer Res* 1986, **46**: 4453–4457.
28. Morrow, C.S. & Cowan, K.H. Mechanisms and clinical significance of multidrug resistance. *Oncology* 1988, **2**: 55–67.
29. Ma, D.D.F., Scurr, R.D., Davey, R.R. *et al.* Detection of a multidrug resistant phenotype in acute non lymphoblastic leukaemia. *Lancet* 1987, **i**: 135–137.
30. Kartner, N., Evernden-Porelle, D., Bradley, G. *et al.* Detection of P-glycoprotein in multidrug-resistant cell lines by monoclonal antibodies. *Nature* 1985, **316**: 820–823.
31. Salmon, S.E., Grogan, T.M., Miller, T. *et al.* Prediction of doxorubicin resistance *in vitro* in myeloma, lymphoma and breast cancer by P-glycoprotein staining. *JNCI* 1989, **81**: 696–701.

Fludarabine

32. Leiby, J.M., Grever, M.R., Staubus, A.E. *et al.* Phase I clinical investigation of fludarabine phosphate by a loading dose and continuous infusion schedule. *JNCI* 1988, **80**: 447–449.
33. Grever, M.R., Kraut, E.H., Neidhart, J.A. *et al.* 2-fluoro-ara-AMP, a Phase I clinical investigation. In: 'The Fourth NCI-EORTC Symposium on New Drugs in Cancer Therapy'. *Invest New Drugs* 1984, **2**: 116–.
34. Hutton, J.J., Von Hoff, D.D., Kuhn, J., *et al.* Phase I clinical investigation of 9 b D arabinofluranosyl-2 fluoroadenine 5' monophosphate (NSC 312887) a new purine anti-metabolite. *Cancer Res* 1984, **44**: 4183–4186.
35. Keating, M.J., Kantarjian, H., Talpas, M. *et al.* Fludarabine therapy in chronic lymphocytic leukaemia. *Nouvelle Revue Française d'Hematologie* 1988, **30**: 461–466.
36. Keating, M.J., Kantarjian, H., O'Brien, S. *et al.* Fludarabine-Prednisone: a safe effective combination in refractory chronic lymphocytic leukaemia. *Proc Am Soc Clin Oncol* 1989, **8**: 201 (abs. 780).

Solid tumours

37. Wolmark, N., Fisher, B., Rockette, H. *et al.* Postoperative adjuvant chemotherapy or BCG for colon cancer: results from NSA BP protocol C-01. *JNCI* 1988, **80**: 30–36.

38. Wittes, R.E. Progress in a desert? Oasis or mirage. *JNCI* 1988, **80**: 5–7.
39. Seifert, P., Baker, L.H., Reed, M.L. *et al.* Comparison of continuously-infused 5-fluorouracil with bolus injection in the treatment of patients with colorectal carcinoma. *Cancer* 1975, **36**: 123–128.
40. Gramont, A., Krulik, M., Cady, J. *et al.* High dose folinic acid and 5-fluorouracil bolus and continuous infusion in advanced colorectal cancer. *Eur J Cancer* 1988, **24**: 1499–1503.

Bone marrow transplantation

41. Mayer, R.J. Allogeneic transplantation versus intensive chemotherapy in first-remission acute leukaemia. Is there a 'best choice'? *J Clin Oncol* 1988, **6**: 1532–1536.
42. Appelbaum, F.R., Fisher, L.D., Donnell-Thomas, E. & Seattle Marrow Transplant Team. Chemotherapy v. marrow transplantation for adults with acute non-lymphocytic leukaemia – a five-year follow up. *Blood* 1988, **72**: 179–184.
43. Champlin, R.E., Ho, W.G., Gale, R.P. *et al.* Treatment of acute myelogenous leukaemia: a prospective controlled trial of bone marrow transplantation versus consolidation chemotherapy. *Ann Intern Med* 1985, **102**: 285–291.
44. Zander, A.R., Keating, M., Dicke, K. *et al.* A comparison of marrow transplantation with chemotherapy for adults with acute leukaemia of poor prognosis in first complete remission. *J Clin Oncol* 1988, **6**: 1548–1557.
45. Champlin, R. & Gale, R.P. Bone marrow transplantation for acute leukaemia: recent advances and comparison with alternative therapies. *Semin Haematol* 1987, **24**: 55–67.
46. Cassileth, P.A., McGlave, D.P., Harrington, J.D. *et al.* Comparison of post-remission therapy in AML. Maintenance versus intensive consolidation therapy versus allogeneic bone marrow transplantation. *Proc Am Ass Clin Oncol* 1989, **8**: 197 (abs. 765).
47. Pinkel, D. Allogeneic bone marrow transplantation in children with acute leukaemia: a practice whose time has gone. *Leukaemia* 1989, **3**: 242–244.

Consequences of intensive therapy

48. Pinkel, D. Five year follow up of total therapy of childhood lymphocytic leukaemia. *JAMA* 1971, **216**: 648–652.
49. Soni, S., Marten, G., Pitner, S. *et al.* Effect of central nervous system irradiation on neuropsychological functioning of children with acute lymphocytic leukaemia. *N Engl J Med* 1975, **293**: 113–118.
50. Eiser, C. Intellectual abilities among survivors of childhood leukaemia as a function of CNS irradiation. *Arch Dis Child* 1978, **53**: 391–395.
51. Eiser, C. Effects of chronic illness on intellectual development: a comparison of normal children with those treated for childhood leukaemia and solid tumours. *Arch Dis Child* 1980, **55**: 766–770.
52. Meadows, A., Massan, D., Fergusson, J. *et al.* Decline in IQ score and cognitive dysfunction in children with acute lymphocytic leukaemia treated with cranial irradiation. *Lancet* 1981, **i**: 1015–1018.
53. Poplack, D.G. & Brouwers, P. Adverse sequelae of central nervous system therapy. In Nesbit, M.E. (ed) *Clinics in Oncology*, vol. 4. W.B. Saunders Company, London, Philadelphia, Toronto, 1985, pp. 263–285.
54. Williams, J.M. & Davis, K.A. Central nervous system prophylactic treatment for childhood leukaemia. Neuropsychological outcome study. *Cancer Treat Rev* 1986, **13**: 113–127.
55. Tamaroff, M., Salwen, R., Miller, D.R. *et al.* Neuropsychological sequelae in irradiated (1800 RADS r and 2400 r) and non-irradiated children with acute lymphoblastic leukaemia. *Proc Am Soc Clin Oncol* 1985, **4**: 165.

56. Mulhearn, R.K., Wasserman, A.L., Fairclough, D. & Ochs, J. Memory function in disease-free survivors of childhood acute lymphocytic leukaemia given CNS prophylaxis with or without 1800 cGy cranial irradiation. *J Clin Oncol* 1988, **6**: 315–320.
57. Peylan-Ramu, N., Poplack, D.G., Blei, C.L. *et al.* Computer assisted tomography in methotrexate encephalopathy. *J Computer Assisted Tomography* 1977, **1**: 216–221.
58. Brouwers, P., Riccardi, R., Fedio, P. Long term neuropsychological sequelae of childhood leukaemia: correlation with CT brain scan abnormalities. *J Pediatr* 1985, **106**: 723–730.
59. Day, R.E., Kingston, J., Bullimore, J.A. *et al.* CAT brain scans after central nervous system prophylaxis for acute lymphoblastic leukaemia. *Br Med J* 1978, **ii**: 1752–1753.
60. Kramer, J.H., Norman, D., Brant-Zawadzki, M. Absence of white matter changes on magnetic resonance imaging in children treated with CNS prophylaxis therapy for leukaemia. *Cancer* 1988, **61**: 928–930.

Imaging disease

61. Tumen, S.S., Rosenthal, D.S., Kaplan, W.D. *et al.* Lymphoma: evaluation with GA SPECT. *Radiology* 1987, **164**: 111–114.
62. Glass, R.B., Fernback, S.K., Conway, J.J. *et al.* Gallium scintigraphy in American Burkitt lymphoma: accurate assessment of tumour load and prognosis. *Am J Radiol* 1985, **145**: 671–676.
63. Anderson, K.C., Leonard, R.C., Canellos, G.P. *et al.* High-dose gallium imaging in lymphoma. *Am J Med* 1983, **175**: 327–331.
64. Israel, O., Front, D., Lam, M. *et al.* Gallium 67 imaging in monitoring lymphoma: response to treatment. *Cancer* 1988, **61**: 2439–2443.
65. Weiner, M., Leventhal, B., Brecher, M. *et al.* Gallium 67 scans as an adjunct to clinical restaging in pediatric patients with Hodgkin's disease. *Proc Am Soc Clin Oncol* 1989, **8**: 278 (abs. 1085).

Supportive care

66. Corey, L., Benedetti, J., Critchlow, C. *et al.* Treatment of primary first episode genital simplex virus infections with acyclovir. Results of topical intravenous and oral therapy. *J Antimicrob Chemother* 1983, **12** (Suppl): 79–88.
67. Shepp, D.H., Dandliker, P.S., Meyers, J.D. Treatment of varicella zoster virus infection in severely immunocompromised patients; a randomised comparison of acyclovir and vidarabine. *N Engl J Med* 1986, **314**: 208–212.
68. Henry, K., Cantrill, H., Fletcher, C. *et al.* Use of intra-vitreous ganciclovir (dihydroxy propoxy methyl guanine) for cytomegalovirus retinitis in a patient with AIDS. *Am J Ophthalmol* 1987, **103**: 17–23.
69. Reed, E.C., Dandliker, P.C., Meyers, J.D. Treatment of cytomegalovirus pneumonia with 9-(2-hydroxy-1 hydroxymethyl) ethoxymethyl guanine and high-dose corticosteroids. *Ann Intern Med* 1986, **105**: 214–215.
70. Whitley, R.J. Ganciclovir – have we established clinical value in the treatment of cytomegalovirus infections? *Ann Intern Med* 1988, **108**: 452–454.
71. Jacobson, M.A. & Mills, J. Serious cytomegalovirus disease in the Acquired Immuno-Deficiency Syndrome (AIDS). *Ann Intern Med* 1988, **108**: 585–594.
72. Chachoua, A., Dieterich, D., Krasinski, K. *et al.* 9-(13-2-propoxy methyl) guanine (ganciclovir) in the treatment of cytomegalovirus gastrointestinal disease with the acquired immuno-deficiency syndrome. *Ann Intern Med* 1987, **107**: 133–137.
73. Hasling, C., Charles, P., Mosekilde, L. Etidronate disodium in the management of malignancy – related hypercalcaemia. *Am J Med* 1987, **82** (Suppl 2A): 51–54.
74. Heath, D.A. Hypercalcaemia in malignancy. *Br Med J* 1989, **298**: 1468–1469.
75. Siris, E.S., Sherman, W.H., Baquiran, D.C. *et al.* Effect of dichloromethylene diphosphonate on skeletal mobilisation of calcium in multiple myeloma. *N Engl J Med* 1980, **302**: 310–315.
76. Delmas, P.D., Charhon, S., Chapuy, M.C. *et al.* Long-term effects of dichloromethylene diphosphonate (Cl₂ MDP) on skeletal lesions in multiple myeloma. *Metabolism Bone Disease and Rel Res* 1982, **4**: 163–168.