formation and breakdown. The administration of heparin in this situation may have swung the balance to a predominance of fibrinolysis over fibrin breakdown.

N.B. Argent
D. Veale
*Royal Victoria Infirmary,
Queen Victoria Road,
Newcastle upon Tyne
NE1 4LP, UK.*

Reference

1. Jacques, L.B. The new understanding of the drug heparin. *Chest* 1985, **88**: 751.

Metirapone-induced alopecia

Sir,

Metirapone, the adrenal 11-beta hydroxylase inhibitor is occasionally used to treat pituitary Cushing's syndrome when other means are considered inappropriate.^{1,2}

An 85 year old female with a history of tiredness, forgetfulness, facial pigmentation and hypokalaemia was investigated. Her 09.00 h cortisol was 977 nmol/l, mid-night was 952 nmol/l (normal range less than 250 nmol/l). The 24 hour urinary cortisol was 764 nmol/24 hours, (normal range less than 350 nmol/24 hours). A 3-day dexamethasone suppression test at 2 mg four times a day showed minor suppression with a 09.00 h cortisol of 766 nmol/l. A subsequent contrast enhanced cranial computed tomographic (CT) and abdominal CT scan showed adrenal hyperplasia more marked on the left. No pituitary tumour was demonstrated.

She refused pituitary irradiation or yttrium implants and was started on metirapone 750 mg four times/day

reducing to 500 mg twice daily. After a year her serum cortisol had fallen to 320 nmol/l. Following a subsequent illness (a complete heart block secondary to an anterior myocardial infarct), her metirapone was reduced further to 250 mg twice daily. Her urinary cortisol was subsequently 1427 and 1513 (normal range less than 350). Her metirapone was increased to 500 mg twice daily. Two months later her urinary cortisol had fallen back to 646 nmol/l, over 24 hours. She developed a frank alopecia at this time. Her only other drugs were frusemide and amiloride (Frumil) two tablets mane and paracetamol and dihydrocodeine (Co-dydramol) 2 tablets 6 hourly as required.

There is one other report linking metirapone with alopecia.³ This was dose related and reversible. Our patient died of an unrelated illness before the metirapone could be withdrawn. The cortisol response to metirapone suggests that she had Cushing's disease rather than ectopic ACTH. Metirapone is well known to cause nausea, vomiting, dizziness, headache and hypotension but alopecia is not a well recognized association. The exact cause remains unknown.

Richard Harries-Jones
Peter Overstall
*The General Hospital,
Nelson Street,
Hereford HR1 2PA, UK.*

References

1. Gold, E.M. Cushing's syndrome – changing views of diagnosis and treatment. *Ann Int Med* 1979, **90**: 829–844.
2. Jeffcoate, W.J., Rees, O.H., Tomlin, F. *et al.* Metirapone in long-term management of Cushing's disease. *Br Med J* 1977, **2**: 215–217.
3. Harris, P.L. Alopecia associated with long term metirapone use. *Clin Pharm* 1987, **5**: 66–68.