Leading Article

Immunopathology of Behçet’s disease

R.J. Powell and S. Dunstan

Immunology Department, University Hospital, Queen’s Medical Centre, Nottingham NG7 2UH, UK

Behçet’s disease (BD) is best considered as a systemic vasculitis in which recurrent orogenital ulceration is a prominent feature. In 1937 Hulusi Behçet described the syndrome that bears his name¹ but the term ‘Behçet’s disease’ is now used in preference to ‘Behçet’s syndrome’. Behçet, himself, recognized the problems related to its diagnosis for which there remains no diagnostic test. BD is rare in Britain but it affects up to 1:1000 of the population of Japan. The increased incidence in Far Eastern countries² and in the eastern Mediterranean,³ and an association with HLA B5^{4,5} suggest that environmental and genetic factors are important influences in the aetiology of BD.

Until recently, 5 sets of diagnostic clinical criteria for BD were variously used around the world, each with its own emphasis on symptoms and signs pertaining to a particular geographical region. In 1990 new criteria for the diagnosis of BD, proposed by an international study group, were published⁶ and these should facilitate comparisons between studies and hopefully foster collaborative work.

Criteria:-	
Recurrent oral ulceration	minor aphthous, major aphthous, or herpetiform ulceration which has recurred > 3 times in a one year period.
Plus 2 out of:-	
Recurrent genital ulceration	aphthous ulceration or scarring.
Eye lesions	Anterior/posterior uveitis, or cells in vitreous on slit lamp examination; or retinal vasculitis observed by ophthalmologist.
Skin lesions	erythema nodosum, pseudo-folliculitis, or papulopustular lesions;

	or acneiform nodules observed by physician in post-adolescent patients not on corticosteroid treatment.
Positive pathergy test	Read by physician at 24–48 h.
(All findings applicable only in absence of other clinical explanations).	

Therefore, recurrent oral ulceration is now an absolute criterion for the diagnosis of BD, requiring, in addition, 2 criteria selected from genital ulceration, ocular inflammation, certain skin lesions and the pathergy test to make a definite diagnosis. A positive pathergy test is the development of cutaneous nodular pustules at a site of trauma, after approximately 24 hours. A needle stab site from venepuncturing would qualify as an appropriate site to inspect for such pustules, but it must be noted that this test can vary considerably between geographical areas.

Although clinical features such as arthritis and cutaneous manifestations are compatible with an immune complex mediated process the overall disease pattern cannot be entirely explained on such a basis. It is not, therefore, surprising that characteristic serological abnormalities are absent in BD. Patients with active disease may generate an acute phase response leading to a raised erythrocyte sedimentation rate (ESR), increased serum levels of C-reactive protein (CRP) and elevated plasma complement components such as C3, C4, C9 and Factor B.^{7–9} The ESR, CRP and C9 seem to reflect the disease activity in some patients, but C9 is not available as a routine test in most laboratories. Polyclonal increases in serum IgA associated with certain clinical subtypes such as ocular, arthritic and neurological, have been reported¹⁰ but results from studies of immune complexes, both circulating and in tissues, have not demonstrated consistent changes. Reduced fibrinolytic activity is seen in BD and is usually due to decreased production of tissue plasminogen activator; however, this appears to be unrelated to the tendency to develop thromboses.¹¹

The cellular infiltrate in the mucous membrane

Correspondence: R.J. Powell, M.R.C.P.
Received: 7 December 1990

lesions (oro-genital ulcers) of BD is predominantly lymphocytes and monocytes, infiltrating both the dermis and epidermis and around the small blood vessels. Interestingly, as the lesion ages, so the neutrophil infiltrate increases.¹² The cutaneous lesions can be divided into 2 types, those resembling erythema nodosum (EN) and those starting as sterile vesicles/pustules which evolve into deep punched out ulcers. Histology of the EN lesions reveals a small vessel vasculitis involving particularly the venules with a perivascular lymphocytic and mononuclear cellular infiltration and fibrin deposition in the vessel wall, whereas in contrast the punched out ulcers are characterized by a leucocytoclastic vasculitis (neutrophil infiltrate) with fibrinoid necrosis.¹³

Studies of peripheral blood T cells in BD have often given conflicting findings but this may be related to differences in patient populations, laboratory techniques, disease activity and therapy. Using a fluorescence cell scanner (FACScan), we have recently confirmed previous reports of a decrease in total peripheral blood T cells (CD3 +)¹⁴ and this resulted primarily from a reduction in the CD4 + population (T helper/inducer). In addition we demonstrated an overall increase in the proportion of activated (HLA-DR +) T cells and it transpired that all the subsets of T cells contributed to this, indicating marked cellular activation. In patients with active disease an increase in natural killer (NK) cells in peripheral blood is reported; however, their killing activity against K562 target cells is reduced and this may be related to immature forms of NK cells being released into the circulation.^{15,16} Both *in vitro* and *in vivo* studies of neutrophils from patients with BD have demonstrated enhanced neutrophil migration, particularly with respect to the number of cells migrating; this effect was again most prominent in those with active disease.¹⁷

Although the aetiology of BD remains obscure, recent clinical observations suggest that exposure to streptococcal antigens may be a major provoking factor for disease activity in BD. Although cell-mediated immunity appears to be intact when judged by intradermal skin testing with PPD, Candida, and streptokinase – streptodornase antigens,¹⁸ an intense delayed type hypersensitivity (DTH) skin reaction is observed when other streptococcal antigens (streptococcal vaccines) are in-

jected intradermally.^{19,20} A recent clinical report describes the onset of severe symptoms of BD shortly after dental treatment in two otherwise stable BD patients.²¹ The authors also cite the onset of similar symptoms in 4 patients who received streptococcal antigen skin tests. The authors argued that it was unlikely that the association was coincidental, or induced non-specifically by dental manipulation. They suggested that it was most likely that the flares of BD were induced by streptococcal related antigens used for skin tests, or released by certain microorganisms which entered the circulation from the oral cavity during the dental procedures. Kaneko *et al.*²⁰ had previously reported that patients with BD have chronically infected foci before the onset of diseases such as dental diseases and tonsillitis. They not only isolated streptococci from these infected foci but also demonstrated binding of a streptococcal group D antiserum to skin and mucosal lesional biopsies from patients with BD, particularly to blood vessels and at inflammatory foci.²²

The possibility of viral rather than bacterial infections triggering BD must not be ignored and, in fact, a viral aetiology was suggested by Behçet in his original publication based on his observation of inclusion bodies in the ulcerations.¹ Herpes simplex virus type 1 (HSV1) has long received attention as a likely candidate and HSV1 genome has been identified by hybridization in peripheral blood lymphocytes and monocytes of BD patients.^{23,24} T cell responses to HSV1 show impaired ³H-thymidine uptake by CD4 + cells in BD when compared to seropositive HSV1 controls, indicating an impaired cellular response specific to HSV1.²⁵

Major aberrations in the cellular immune system have therefore been demonstrated in BD and the intriguing possibility of BD being triggered by streptococcal antigens is being actively pursued. The multifactorial aetiology of this condition was recently exemplified by a study from Leeds entitled 'Is Behçet's disease triggered by childhood infection?' The group showed that an increased risk of BD was associated with tonsillectomy, a history of cold sores, large sibship size, late birth order, travel to countries with high incidence of disease and the first sexual intercourse before 16 years of age.²⁶ One is left to wonder what were the features with which there were no correlations!

References

1. Behçet, H. Über rezidivierende, aphthöse, durch ein Virus verursachte Geschwüre am Mund, am Auge und an den Genitalien. *Dermatol Wochenschr* 1937, **36**: 1152–1157.
2. Aoki, K., Fujioka, K., Katsumata, H. *et al.* Epidemiological studies on Behçet's disease in the Hokkaido district. *Jpn J Clin Ophthalmol* 1971, **25**: 2239–2248.
3. Chajek, T. & Fainaru, M. Behçet's disease. Report of 41 cases and a review of the literature. *Medicine* 1975, **54**: 179–196.
4. Lehner, T., Batchelor, J.R., Challacombe, S.J. & Kennedy, L. An immunogenetic basis for the tissue involvement in Behçet's syndrome. *Immunology* 1979, **37**: 895.

5. Yazici, H., Chamberlain, M.A., Schreder, I., D'Amaro, J. & Muftuoglu, M. HLA antigens in Behçet's disease: a reappraisal by a comparative study of Turkish and British patients. *Ann Rheum Dis* 1980, **39**: 344–348.
6. International Study Group for Behçet's Disease. Criteria for diagnosis of Behçet's disease. *Lancet* 1990, **335**: 1078–1080.
7. Oshima, Y., Shimizu, T., Yokohari, R. *et al.* Clinical studies on Behçet's syndrome. *Ann Rheum Dis* 1963, **22**: 36–45.
8. Adinolfi, M., Beck, S.E. & Lehner, T. Serum levels of acute phase proteins, C9, factor B and lysozyme in Behçet's syndrome and recurrent oral ulcers. In: Lehner, T. and Barnes, C.G. (eds) *Behçet's Syndrome: Clinical and Immunological Features*. Academic Press, London, 1979, pp. 107–139.
9. Rumfeld, W.R., Morgan, B.P. & Campbell, A.K. The ninth complement component in rheumatoid arthritis, Behçet's disease and other rheumatic diseases. *Br J Rheumatol* 1986, **25**: 266–270.
10. Scully, C.M., Lehner, T. & Harfitt, R. Serum, salivary and lacrimal immunoglobulins in Behçet's syndrome and recurrent oral ulcers. In: Lehner, T. and Barnes, C.G. (eds) *Behçet's Syndrome: Clinical and Immunological Features*. Academic Press, London, 1979, pp. 77–89.
11. Aitchison, R., Chu, P., Cater, D.R., Harris, R.J. & Powell, R.J. Defective fibrinolysis in Behçet's syndrome: significance and possible mechanisms. *Ann Rheum Dis* 1989, **48**: 590–593.
12. Muller, W. & Lehner, T. Quantitative electron microscopical analysis of leukocyte infiltration in oral ulcers in Behçet's syndrome. *Br J Dermatol* 1982, **106**: 535–544.
13. Lakhanpal, S., Desmond O'Duffy, J. & Lie, J.T. Pathology. In: Plotkin, G.R., Calabro, J.J., Desmond O'Duffy, J. (eds). *Behçet's Disease: A Contemporary Synopsis*. Futura Publishing Co., Mount Kisco, N.Y., 1988, pp. 102–142.
14. Ahmed, A.R. Lymphocyte studies in Behçet's syndrome. *Dermatologica* 1982, **164**: 175–180.
15. Kaneko, F., Takahashi, Y., Muramatsu, R. *et al.* Natural killer cell numbers and function in peripheral lymphoid cells in Behçet's disease. *Br J Dermatol* 1985, **113**: 313–318.
16. Hamzaoui, K., Ayed, K.H., Hamza, M. & Touraine, J.L. Natural killer cells in Behçet's disease. *Clin Exp Immunol* 1988, **71**: 126–131.
17. Efthimiou, J., Addison, I.E. & Johnson, B.V. *In vivo* leucocyte migration in Behçet's syndrome. *Ann Rheum Dis* 1989, **48**: 206–210.
18. Lehner, T. Immuno-pathology of Behçet's syndrome. In: Lehner, T. & Barnes, C.G. (eds). *Behçet's Syndrome: Clinical and Immunological Features*. Academic Press, London, 1979, pp. 127–139.
19. Graykowsi, E.A., Barile, M.F., Lee, W.B. & Stanley, H.R. Jr. Recurrent aphthous stomatitis: clinical, therapeutic, histopathologic and hypersensitivity aspects. *JAMA* 1966, **196**: 637–644.
20. Kaneko, F., Kaneda, T., Ohnishi, O. *et al.* Behçet's syndrome and infection allergy. (1) Detection of chronic infectious foci and immune responses to bacterial vaccines *in vivo* and *in vitro*. *Jpn J Allergol* 1978, **27**: 440–447.
21. Mizushima, Y., Matsuda, T., Hoshi, K. & Ohno, S. Induction of Behçet's disease symptoms after dental treatment and streptococcal antigen skin test. *J Rheumatol* 1988, **15**: 1029–1030.
22. Kaneko, F., Takahashi, Y., Muramatsu, Y. & Miura, Y. Immunological studies on aphthous ulcer and erythema nodosum-like eruptions in Behçet's disease. *Br J Dermatol* 1985, **113**: 303–312.
23. Eglin, R.P., Lehner, T. & Subak-Sharpe, J.H. Detection of RNA complementary herpes-simplex virus in mononuclear cells from patients with Behçet's syndrome and recurrent oral ulceration. *Lancet* 1982, **ii**: 1356–1360.
24. Bonass, W.A., Bird-Steward, J.A., Chamberlain, M.A. & Halliburton, I.W. Molecular studies in Behçet's syndrome. In: Lehner, T. & Barnes, C.G. (eds). *Recent Advances in Behçet's Disease*. Royal Society of Medicine Services, London, 1986, pp. 37–41.
25. Young, C., Lehner, T. & Barnes, G.C. CD4 and CD8 cell responses to herpes simplex virus in Behçet's disease. *Clin Exp Immunol* 1988, **73**: 6–10.
26. Cooper, C., Pippard, E.C., Sharp, H., Wickham, C., Chamberlain, M.A. & Barker, J.P. Is Behçet's disease triggered by childhood infection? *Ann Rheum Dis* 1989, **48**: 421–423.