Reviews in Medicine

Hepatology

Phillip M. Harrison, Johnson Y.N. Lau and Roger Williams

Institute of Liver Studies, King's College Hospital and School of Medicine and Dentistry, Bessemer Road, London SE5, UK

Introduction

Recently the field of hepatology has seen remarkable advances in the identification and understanding of the agents causing viral hepatitis. The medical therapy of patients with fulminant hepatic failure continues to progress and this is reflected in the improved survival rates which are now being achieved; while for those patients in whom liver function will never recover with medical treatment alone, emergency liver transplantation is now an established therapeutic option. Graft survival continues to improve as developments in immunosuppression are made. This review will cover the recent advances in these areas and in the management of portal hypertension.

Viral hepatitis

Viral hepatitis continues to be a major global health problem but with the advancement of many scientific disciplines including molecular biology and immunology, rapid advances are being made in the understanding of the viruses and development of specific vaccines. We now know that there are at least five major types of primary hepatotropic acute hepatitis viruses (Table I). This following section reviews the progress made recently on these hepatotropic viruses. Acute hepatitis caused by other viruses in which involvement of other organs is more common and more prominent than evidence of liver involvement, notably cytomegalovirus and herpes simplex virus, is not described here.

Hepatitis A virus (HAV)

HAV was first discovered in faeces of infected patients by immune electron microscopy in 1973.¹

Today, it is still ranked by the World Health Organization as a prevalent infection worldwide; especially in countries where overcrowding and poor standards of hygiene and sanitation are prevalent as shown by the recent major outbreak involving 1.2 million persons in mainland China in 1988.

Virology

Although HAV is classified as a picornavirus, more and more evidence supports the view that HAV is different from the well-studied four genera of this virus family.² The genome of HAV is a linear, single-stranded ribonucleic acid of messenger-sense (positive) polarity. RNAs extracted from different HAV isolates have been cloned and sequenced.^{3,4} According to these studies, the molecule consists of a total of 7,478 nucleotides, and,

Table I Classification of hepatotropic and other viruses responsible for acute viral hepatitis

Hepatitis type	Previously named/description
<i>Hepatotropic viruses</i>	
Hepatitis A (HAV)	Infectious hepatitis
Hepatitis B (HBV)	Serum hepatitis
Variants:	1. Wide-type
	2. African HBsAg- anti-HBc + mild disease course
	3. Mediterranean HBsAg-HBV-DNA + aggressive liver disease
Hepatitis C (HCV)	Parenteral non-A non-B hepatitis
Hepatitis D (HDV)	Delta hepatitis (agent)
Hepatitis E (HEV)	Enteral non-A non-B hepatitis
<i>Others</i>	
	Cytomegalovirus (CMV)
	Epstein-Barr virus (EBV)
	Herpes simplex virus
	Coxsackie B virus
	Echovirus
	Yellow fever virus

Correspondence: R. Williams, M.D., F.R.C.P., F.R.C.S., F.R.C.P.E.

like the genome of the typical picornavirus, it can be functionally divided into a 5'-terminal untranslated region composed of 733 nucleotides, a single open reading frame of 6,681 nucleotides, and a short untranslated segment of 64 nucleotides at the 3'-end of the molecule.⁵ The open reading frame gives rise to a precursor polypeptide which is subsequently cleaved into four different polypeptides VP1-4 (Figure 1). Existing information suggests that the molecular sizes of these peptides are VP1 33 kD, VP2 27 kD, VP3 29 kD, VP4 (truncated) 17 amino acids. Around 60 copies of each of these four structural proteins, VP1 to VP4, with molecular weights in the range of 8,000–33,000, form the capsid of the typical mature HAV.^{2–4,6} The most important differences between HAV and the four conventional genera of the picornavirus family apparently concern composition and function of the viral capsid as evidenced by the size of its structural proteins, the outstanding overall stability of the virus, and the resolution of only one immunodominant neutralization site.^{2,7,8}

After years of frustrating search for a cell system in which HAV could be propagated *in vitro*, there

are now a number of cultures of established susceptibility.^{9,10} Independent of the culture system used for the primary isolation, all HAV strains have protracted replication and establish persistent infection.^{11,12} More recently, two HAV cell systems have been described in which replication of HAV seemed to be paralleled by virus-specific cytopathology.^{13–15}

Pathogenesis of liver damage

The pathogenetic mechanism of liver cell damage in acute HAV infection is still unclear. The absence of convincing evidence that HAV gives rise to true persistent infection *in vivo* in contrast to *in vitro* replication system, strongly suggests that a close interplay between the virally controlled and host immunological factors is necessary to cause a cytolytic infection with subsequent elimination of the virus.² The postulation that hepatocyte destruction is mediated by anti-HAV antibody with or without the help of complements is untenable^{2,16,17} and the interferon system is also unlikely to play a major role in the elimination of the virus *in vivo*.¹⁸

Figure 1 Structure of the hepatitis A virus genome and the tentative protein cleavage sites. VPg – genome linked viral protein.

Two recent studies provide strong evidence that cytotoxic T cells capable of lysing HAV- infected target cells develop in the course of HAV infection.^{19,20} This effect is also shown to be virus-specific and is functionally restricted by the major histocompatibility complex (MHC).

Clinical features and laboratory findings

HAV usually causes a minor or unnoticed illness in children and young adults. On a worldwide scale, less than 5% of the cases are recognized clinically.^{21,22} It can cause a high mortality in patients over the age of 60.²³ In a recent major outbreak in Shanghai, China, approximately one third of those shown serologically positive for acute HAV infection were asymptomatic, and only around 20% had overt clinical hepatitis.²⁴ Systemic complications like sinus bradycardia and transient T-wave changes on electrocardiography, arthralgia and palpable splenomegaly may occur in less than 15% of cases.²³ Although prolonged cholestasis is well described in HAV infection, it is relatively uncommon on the whole but just disproportionately frequent in patients referred to hospitals because of delayed recovery.²⁵

The occurrence of a relapsing course in some patients and the persistence of high titre IgM anti-HAV antibodies for up to 400 days in others, together with the demonstration of intestinal reinfection and prolonged viral excretion may help to explain the mechanism of viral perpetuation in highly endemic areas.²⁶ It has also been recently shown that relapse of HAV infection can occur together with recurrence of IgM anti-HAV.²³

The laboratory findings in HAV infection with elevated serum transaminases are often accompanied by non-specific rise of IgM, IgG, anti-smooth muscle antibodies and acute phase proteins.^{27,28} The raised IgM is mainly polyclonal and only a small proportion is directed against HAV antigen.²⁸ The extent of elevation of serum transaminases does not appear to be of prognostic significance.²⁷ The blood lymphocyte count is usually high initially, but then falls with disease evolution and there may be a pancytopenia phase during the recovery.²⁷ Minor coagulation defects and depressed fibrinogen levels are also common.^{23,27}

Passive and active immunization

High titre serum immunoglobulin preparations have long been recognized to be of value in both pre-exposure and post-exposure prophylaxis against HAV infection.²⁹ A recent leading article in the *British Medical Journal* suggested that not all travellers from developed countries to highly endemic areas required protective immunoglob-

ulins routinely as screening of patients for antibodies against HAV before giving immunoglobulins is economically sensible and results in effective use of a limited resource.³⁰

Although HAV infection does not lead to chronic hepatitis and cirrhosis, it is an important source of morbidity and economic losses.^{22,23} An effective, safe, and low cost vaccine would be beneficial, especially in the developed countries in which the incidence of HAV infection is low. Killed HAV vaccine has been shown to be safe and efficient in animals and clinical studies are ongoing.³¹ This vaccine, however, requires up to three injections to induce protective levels of antibody in recipients. Attenuated HAV vaccine and recombinant HAV vaccine have also been developed and are being investigated.³¹⁻³³

Hepatitis B virus (HBV)

It has been estimated that there are approximately 300 million chronic HBsAg (hepatitis B surface antigen) carriers worldwide.³⁴ The World Health Organization estimates suggest that 40% of them will eventually die of chronic liver disease and/or hepatoma.³⁵

Virology

The HBV genome has now been well characterized and consists of four overlapping open reading frames designated S/pre-S, C, P, and X. An enhancer region, a glucocorticoid responsive element as well as two direct repeats (DR1 and DR2) have also been identified (Figure 2).^{36,37} Each is flanked by start and stop codons and encodes a viral polypeptide (Table II). The S/pre-S region of the genome codes for the proteins of the viral envelope.³⁶ Recent evidence suggests that pre-S1 sequences are involved in the binding of virions to a putative receptor on the surface of hepatocytes which may be the first step in the infectious cycle.³⁸ Less convincing is the evidence that the pre-S2 region carries a binding site or receptor for polymerized human serum albumin (pHSA) which in turn binds to hepatocytes.³⁹⁻⁴⁴ This finding is interesting but pHSA is an *in vitro* product (albumin treated with glutaraldehyde) and has never been found in human subjects.^{45,46} Hence, it is unlikely that this is the mechanism of viral entry into the hepatocytes. Recently, W.H. Gerlich and his colleagues demonstrated that human monomeric serum albumin can bind irreversibly to cysteine of the small HBsAg protein, suggesting an alternative way of viral entry into hepatocytes.⁴⁷ There is also evidence that a lack of antibodies to this protein in the early phase of HBV infection may result in chronic HBV carriage in some

Figure 2 Structure of the HBV genome. The positions of the four open reading frames (ORFs) known to produce HBV proteins are shown in open boxes. The relative positions of the enhancer element (ENH), glucocorticoid responsive element (GRE), and the direct repeats (DR1 and DR2) are also shown. (+) and (–) denotes the positive and negative strands. Small, solid arrows show the 5' ends of the three major virus RNA transcripts of 2.1, 2.4, 3.5 kb.

Table II HBV polypeptides

HBV gene	Polypeptide
S	HBsAg (major protein)
pre-S2 + S	pre-S2 peptide (middle protein)
pre-S1 + pre-S2 + S	pre-S1 peptide (large protein)
C	HBcAg and HBeAg
pre-C + C	pre-core peptide
P	DNA polymerase
X	X protein

patients.⁴⁸ Both HBcAg (hepatitis B core antigen) and HBeAg (hepatitis B e antigen) are encoded by the C gene, which includes a pre-C region.^{37,38} Existing evidence suggests that HBcAg is the dominant target of the immune attack.^{49–54} Furthermore, the presence of cytoplasmic and membranous/submembranous HBcAg in addition to nuclear HBcAg induces immune lysis of the infected hepatocytes, which in turn may clear viral replication in some patients.⁵¹ When pre-C is transcribed, the core protein can also be targeted to the endoplasmic reticulum, where it is subsequently cleaved and HBeAg is secreted from the cell into the blood.^{55,56} The presence of HBeAg in the blood is therefore, an indicator of ongoing viral replication. Early results suggest that X protein may also be related to HBV replication.^{57–59}

Two important HBV variants have been identified recently (Table I). The first variant occurs most commonly in Africans with an incidence of between 15 to 40%.⁶⁰ These patients had positive serum anti-HBc but not HBsAg. It is not as yet defined whether this represents infectivity or immunity, but it is likely that this group comprises both categories. The second variant occurs most commonly in Mediterranean population. They have serum HBeAg negative, serum HBV-DNA positive, and often have severe progressive chronic hepatitis. It has recently been characterized that this HBV variant has a single point mutation in the pre-C sequence which generates a translational stop codon.^{61,62} The absence of pre-C protein excludes the production of HBeAg and this explains the persistence of HBV replication in some anti-HBe positive patients.

The development of the very powerful molecular biology technique, polymerase chain reaction, apart from helping sequencing of the variant viruses, allows the detection of very low levels of DNA in serum. With this technique, the frequent involvement of HBV in a proportion of HBsAg seronegative patients with chronic liver disease, including primary hepatocellular carcinoma, was revealed.⁶³

Pathogenesis of liver damage

The existence of HBV carrier state for many years without evidence of parenchymal liver damage suggests that HBV is not cytopathic in its natural host, man.⁶⁴ This may, however, not be true in animal models; for example, recent studies in the transgenic mouse have shown that high level expression of HBsAg is associated with the onset of hepatocellular degeneration and necrosis.^{65,66}

All the early studies, including immunohistochemical studies and autologous hepatocyte cytotoxicity, suggested that immune-mediated elimination of nucleocapsid antigen-containing hepatocytes from the liver caused many patients to develop permanent liver damage.⁶⁷ This was further supported by the association of CD8 positive T cells with liver damage and that this T cell cytotoxicity can be abolished by antibodies to either nucleocapsid antigens or HLA class I antigen, suggesting that nucleocapsid antigen is the target for T cell immune attack.^{50,51}

The target for non-T cell cytotoxicity has been less well studied. Previous studies have shown that this can be blocked by liver-specific membrane lipoprotein (LSP), aggregated IgG or the F(ab')₂ fragment of anti-human IgG, suggesting that non-T lymphocytes mediate antibody-dependent cell-mediated cytotoxicity against a component of LSP.⁶⁸

Immunological defects in chronic HBV infection

Various immune defects have been demonstrated in chronic HBV carriers which may have a role in underlying the persistence of HBV infection in these patients. Interferon-alpha production is active in acute but deficient in chronic HBV infection.^{69,70} The capacity of monocytes to activate lymphocytes is also abnormal in chronic HBV carriers.⁷¹ Interestingly, interferon-alpha can partially correct this abnormality.⁷² This raises the possibility of interferon deficiency in some patients that predispose them to develop chronicity of HBV infection. The strongest evidence against this hypothesis is that the proportion of adult males that become chronic carriers is high (~5%) and it is highly improbable that there is such an inherited defect of interferon-alpha in the absence of other clinical evidence of immune deficiency in chronic HBV infection.⁶⁷ Recently, it has been shown that HBV itself can inhibit interferon production in an *in vitro* system and this may explain the decreased interferon production observed in this group of patients.⁷³ In another experiment, a plasmid vector containing multiple copies of the hepatitis B virus genome was transfected into a cell line resulting in a reduced sensitivity to interferons.⁷⁴

Other defects include abnormal natural killer cell function and decreased *in vitro* anti-HBs production in active HBV replicators.^{75,76} The production of interleukin 1 and tumour necrosis factor-alpha have also been shown to be increased in patients with chronic HBV infection.^{77,78}

Clinical features and laboratory findings

The serological profiles of acute HBV infection and acute HBV infection progressing to chronic HBV infection are shown on Figure 3 and Figure 4. It has also been shown that histological chronic persistent hepatitis may not necessarily be a mild disease and can be a progressive disease when associated with sustained virus replication.⁷⁹

In the Far East and Middle East, most patients acquire the infection during the peri-natal period and become chronic carriers; whereas in countries like the UK, where most patients acquire their HBV infection in adulthood, more than 95% of the patients with acute HBV infection clear the HBV and less than 5% of the patients develop a carrier state. The basic mechanism underlying this difference is unknown but has been attributed to immune tolerance in childhood.⁸⁰ It is also known that male, intravenous drug users and homosexuals are more prone to develop chronic HBV infection.

Anti-viral therapy

Various approaches have been taken to the treatment of chronic HBV infection (Table III). The only truly promising mode of therapy at present is interferon-alpha.⁸¹⁻⁸⁵ However, the success rate of lymphoblastoid and recombinant interferon-alpha in these patients varies from only 20 to 40%. This variable response rate is not due to variable interferon-alpha receptor expression.^{86,87} Deter-

Figure 3 Clinical, biochemical and serological profile in patients having acute hepatitis B virus infection with recovery.

Chronic hepatitis B infection

Figure 4 Current view of serological and histological profile, and HBV-DNA integration in patients with chronic hepatitis B virus infection.

Table III Agents that have been studied in the treatment of chronic HBV infection

<i>Anti-virals</i>	<i>Immunosuppressive</i>	<i>Immunostimulators</i>
Interferons	Corticosteroids	BCG vaccination
Alpha		Levamisole
Beta		Interleukin-2
Gamma		Interferon-gamma
Adenine arabinoside (Ara-A)		Thymosin
Acyclovir, deoxyacyclovir		
Zidovudine		
Suramin		
Ribavirin		
Phosphonoformate		
Quinacrine		
(+)-cyanidanol-3		
<i>Phyllanthus amarus</i> (plant extracts)		

minants of a likely positive response are summarized in Table IV. As measured by serum HBV DNA, interferon-gamma is also an anti-viral to HBV.⁸⁸⁻⁹⁰ However, its clinical use, either alone or in combination with interferon-alpha, has been shown not to be advantageous.⁸⁸⁻⁹⁰ Further studies are being conducted to evaluate the role of new drugs, combination therapy, and continuous low dose interferon-alpha therapy in these interferon-alpha resistant patients.^{91,92}

Active immunization

The first generation plasma-derived HBV vaccines were licensed in 1981. The fear of transmission of blood-borne diseases as well as their cost had considerable effects in limiting their acceptance and use. With the second generation yeast-derived recombinant HBV vaccines launched in 1986, much more rapid progress is being made with vaccination of the high risk groups such as health

Table IV Patient-related determinants that are thought to influence the response to interferon-alpha therapy

<i>Determinants of positive response</i>	<i>Determinants of negative response</i>
Acquisition in adulthood	Acquisition in infancy/childhood
Female sex	Male sex
High pre-treatment ALT	Low pre-treatment ALT
Low pre-treatment HBV-DNA	High pre-treatment HBV-DNA
Active histology	Inactive histology
HIV antibody negative	HIV antibody positive
Heterosexual	Homosexual
Anti-delta negative	Anti-delta positive

ALT – serum alanine aminotransferase; HIV – human immunodeficiency virus.

personnel and babies born to HBsAg seropositive mothers in most developed countries.^{93,94} In some endemic areas, like Hong Kong and Taiwan, all newborn babies are routinely vaccinated. The World Health Organization has recommended routine HBV vaccination for countries that possess the economic capacity to purchase the vaccine and where the HBV carrier rate exceeds 2.5% of the population. The third generation synthetic peptide vaccines have already been developed; their safety, immunogenicity and protectivity against HBV infection are at present being evaluated. It is hoped that these developments can produce cheap, efficient vaccines to allow for a widespread vaccination programme in Third World countries.

The seroconversion rates of all these vaccines vary from 80 to 95%. Around 35% of healthy HBV vaccine responders will have their anti-HBs titre dropped to below 10 mIU/ml 5 years later. The Immunization Practices Advisory Committee at present does not recommend booster doses for these subjects. However, studies in homosexual men demonstrated that the life-table attack rate of new HBV infections in vaccinated responders was 1.8% annually.^{94,95} Although most of these new infections were subclinical seroconversions, one clinically apparent HBsAg event has been documented.⁹⁵ Recently, Carmen *et al.* showed that HBV mutants may be responsible for some of the patients who developed HBV infection after successful vaccination.⁹⁶

Host factors associated with suboptimal HBV vaccine responsiveness have also been identified (Table V). A supplementary dose may help a small proportion of these non/hypo-responders. New HBV vaccines, which include additional antigens such as pre-S1, pre-S2 and possibly HBc, are being developed.⁹⁷ Another preliminary report suggests that interleukin-2 may be of value for inducing response in some non-responders.⁹⁸

Use of liver transplantation in chronic HBV related diseases is addressed later in the transplantation section of this article.

Hepatitis C virus (HCV)

Following the isolation of HBV in the 1960s and the HAV in 1970s, it became obvious that there was a proportion of cases of clinically viral hepatitis without a defined aetiology. The majority of these cases occurred following blood transfusion of intravenous drug-users and led to chronic hepatitis in a proportion of individuals.^{99,100} In the 1970s, the frequency of transfusion-associated hepatitis measured in prospective trials was approximately 10% ranging from 7–12%.¹⁰¹ The term non-A non-B hepatitis (NANB) was coined to describe these cases.^{99–101} The clinical diagnosis of NANB hepatitis was similarly imprecise, and was based on the clinical exclusion of other causes of hepatocellular inflammation and the serological exclusion of other hepatotropic viruses including HAV, HBV, Epstein-Barr virus (EBV), cytomegalovirus (CMV) and herpes simplex virus (HSV).^{99,100} It has also been suggested that a proportion of NANB hepatitis represented infections with cryptic forms of HBV, in which serological markers of HBV were not expressed, but HBV DNA could be detected in liver and serum. Approaches used to identify HAV, HBV antigens and viral particles were employed in the search of NANB virus and multiple reports were published but none stood the test of the well known serum panel prepared by Dr Harvey Alter at the National Institute of Health.^{102–105}

However, several deductions could be drawn from these early experiments. Firstly, cross-challenge experiments in chimpanzees suggest that

Table V Suboptimal HBV vaccine responsiveness: host factors

• Age	• HIV infection
• Obesity	• Immunosuppression
• Alcohol	• Route of administration
• Chronic illness (e.g. chronic renal failure)	

HIV – human immunodeficiency virus

there are two blood borne NANB hepatitis agents.^{106–112} One of them is probably the predominant agent of transfusion-associated NANB hepatitis. It is chloroform-sensitive and induces tubular ultrastructural cytoplasmic changes in hepatocytes. The other one is chloroform-resistant and does not induce ultrastructural cytoplasmic changes. Secondly, the chloroform sensitivity of the predominant transfusion-associated NANB hepatitis agents is consistent with a lipid envelope in the virus. Thirdly, physiological chemical studies using pore-sized ultrafiltration experiments showed that the NANB hepatitis virus was between 30–60 nm in diameter. Therefore, even before the isolation of the transfusion-associated NANB hepatitis virus, it has already been well-characterized and was consistent with it being a member of a group of small lipid-coated RNA virus (alphavirus and flavivirus).^{107–112}

Identification of HCV

The recent identification of the major non-A, non-B virus which has now been termed HCV was the result of the major advances in molecular cloning techniques.¹¹³ Utilizing large volumes of plasma derived from NANB hepatitis virus-infected chimpanzees provided by the Centre for Disease Control (CDC, Atlanta), the fraction containing the NANB hepatitis virus was pelleted by repeated ultracentrifugation and total nucleic acid was extracted. Although an RNA virus was suspected, both RNA and single-stranded DNA were converted to complementary DNA (cDNA) and inserted into a phage gt11 expression vector used to infect *Escherichia coli* to form a cDNA library. The expression library was tested with sera from patients who had chronic NANB hepatitis, or were convalescing from NANB hepatitis to detect those clones expressing NANB or HCV antigens. After around 1 million clones had been screened, a clone was isolated which produced an antigen detected by the NANB hepatitis convalescent sera. Another overlapping clone from the library was identified subsequently by hybridizing with the initial clone identified. Both clones hybridized with nucleic acids extracted from infected sera but not with control DNA. At around the same time, Arima *et al.*, from Japan, using essentially the same method, except that they used 100 litres of pooled human sera from patients with presumed NANB hepatitis, isolated 29 clones, the sequences of 3 were later found to be similar to that cloned by the Chiron Corporation.¹¹⁴ Three overlapping cDNA clones from the HCV identified by Choo were sequenced and found to contain a single continuous open reading frame (ORF) which was expressed as a fusion polypeptide (c100-3) containing 363 amino acids. These polypeptides were then

used to coat microtitre plates in an antibody capture assay which was then tested against the NIH panel of sera.¹¹⁵ The assay detected high titres of antibody in 6/7 samples known to be NANB hepatitis virus positive but in 0/9 negative controls. Subsequent studies showed that most patients with post-transfusion NANB hepatitis were positive for this antibody by 6 months. Around half of the patients with sporadic NANB hepatitis were also found to be positive at various intervals following the acute infection, suggesting a common aetiology in a proportion of cases.¹¹⁶ Recently, using polymerase chain reaction (PCR) to detect HCV RNA in tissue and serum, half of the patients with presumed chronic NANB hepatitis were positive and there was a good overlap with the immunoassay (c100-3).¹¹⁷

Virology

Current understanding is that HCV is an approximately 10,000 nucleotide linear, single-stranded, positive-polarity RNA virus with proportions similar to those of flavivirus, which is a member of the group B arboviruses (Figure 5).¹¹⁸ Other members in the same group include yellow fever virus and dengue virus. Only one ORF has been identified so far; presumably, there is only one large poly-protein produced that is cleaved post-translationally.¹¹⁸ This virus bears no genomic homology to retrovirus, HBV, hepatitis D virus (HDV) or other known viral agents. Clinically, antibody to HCV appears in the circulation between 1 and 3 months after onset of acute illness, but in rare cases not for up to a year.¹¹⁹ Antigenaemia is so limited that circulating viral antigen is beyond the limit of detection of most conventional assay techniques.¹¹⁸

Currently available serological assays

Using the serological assay for the antibody to non-structural protein of the virus, it has been shown that HCV accounted for 80% of the post-transfusion hepatitis. It also has a high prevalence in patients with cryptogenic liver cirrhosis, hepatocellular carcinoma and autoimmune chronic active hepatitis, as well as in alcoholics, haemophiliacs receiving replacement therapy, intravenous drug users and haemodialysis patients (Table VI).^{118,120–127} Recent studies showed that anti-HCV antibodies only appear several weeks after the infection but persist for more than 10 years in most patients.^{118,119} It also appears that there is high anti-HCV positivity in areas near the Equator. However, two factors have been identified recently to be associated with possible false positivity with this ELISA assay. Firstly, keeping serum samples in hot weather and repeated freezing and thawing of the samples can give possibly false-positive results

Figure 5 Schematic alignment of the polyproteins encoded by the putative domains of the hepatitis C virus. The positions of HCV clones 5-1-1 and C100 discovered by Choo *et al.*¹¹⁸ as well as the three clones identified by Arima *et al.*¹¹⁴ are indicated.

Table VI Anti-HCV antibodies in different geographical areas according to risk of hepatitis and presence of liver disease^a

Groups	Anti-HCV positive (%)			
	Italy	Spain	UK	USA
<i>Group I (Normal population)</i>				
Random blood donors		0	0.66	1–2
Healthy pregnant women		1.2		
Community populations			4	
<i>Group II (High risk population)</i>				
Intravenous drug users	75	92	50–81	76
Haemophiliacs	82	64	85	79
Haemodialysis patients		20	1–3	8
Homosexual men		8	0–26	24
Female contacts of drug users		6		
<i>Group III (High risk patients with liver disease)</i>				
Post-transfusion NANBH	84	85		85
Sporadic NANBH	74	69	29	73
Chronic NANBH	74	62	50	79
Cryptogenic cirrhosis		78		42
Alcoholic cirrhosis		39		
Hepatocellular carcinoma	65	75	60	26
HBsAg +	54	56		
Alcoholic		77		
<i>Group IV (Autoimmune liver disease)</i>				
Autoimmune chronic active hepatitis ^b	78	44	70	43
Primary biliary cirrhosis		38	0	9

NANBH – non-A, non-B hepatitis; ^aExtracted from references 119–127; all these data should be interpreted with caution as there is still no confirmatory test available; ^bNote that a positive anti-HCV test is associated with a high globulin level, suggesting that this may be a false positive. See reference 128.

(Tibbs *et al.*, personal communication). Secondly, studies in this Unit have shown high incidence of anti-HCV antibodies in autoimmune chronic active hepatitis and there was an excellent linear correlation between the ELISA optical density reading and the serum globulin level, suggesting that high globulin levels might produce non-specific reactivity in the ELISA system.¹²⁸

Recently, companies like Abbott Diagnostics, Wellcome Diagnostics and Universal Biological Incorporation have also developed their HCV serological assays. Chiron Company, utilizing the Western blotting technique, has also established a recombinant immunoblot assay (RIBA) and they claim that this test is more sensitive and specific. At the same time, serum HCV RNA was successfully detected employing the techniques of polymerase chain reaction and molecular hybridization.¹¹⁷ The specificity of all these newly developed assays requires further confirmation. As most of these assays only test the antibodies to a non-structural protein, it is impossible to determine whether this represents infectivity or immunity. It is hoped that a viral structural antigen assay can also be developed in the near future.

Anti-viral therapy

Low doses of interferon-alpha (3 million units three times per week) have been shown to be effective in reducing serum aminotransferases and in controlling liver inflammation in chronic hepatitis due to parental non-A, non-B virus infection.^{129,130} Serum transaminase was normalized in around half of the

patients and this is associated with histological improvement. This improvement was, however, only sustained in about 50% of the patients following the cessation of treatment and hence treatment may have to be lifelong. Predictors of a favourable response seem to be short duration of illness and high transaminases. A positive HCV test did not predict favourable outcome nor did the mode of transmission. It has been shown that the titre of HCV antibody declines in some successfully treated patients and this may prove to be a useful response monitor in the future.

Acyclovir has produced no apparent benefit and corticosteroids have yielded disputed results.¹³¹

Hepatitis D virus (HDV)

In 1977 Rizzetto *et al.* reported the observation of a new viral antigen present in the nuclei of hepatocytes in patients with chronic HBV infection.¹³² Subsequent studies revealed that this antigen was, in fact, related to a new virus, now termed HDV.¹³³

Epidemiology

HDV infection is endemic in Italy, Middle East, some areas of Africa and South America, but is rare in Western Europe, North America and Asia, despite the existence of susceptible HBV carriers in many of these areas (Table VII).¹³⁴ Family clustering in endemic areas suggests viral transmission by close contacts, whereas intravenous drug use is likely to be the major route of transmission in nonendemic regions of the Western world.^{133,135} Placental transmission does not appear to be an important route.

Virology

HDV is an incomplete single stranded circular RNA virus that depends on the helper function of HBV to replicate.^{136–138} It is encased in HBsAg and the RNA genome, which is around 1,700 nucleotides long, of minus polarity, contains a very high intramolecular base-pairing, similar to the genomes of plant viroids.^{139,140} Although several open reading frames (ORFs) have been identified, only one (ORF5) has been demonstrated to code for protein (HDAg). Replication of viral RNA has been shown to proceed by a rolling circle mechanism and specific self-cleavage and self-ligation of both genomic and anti-genomic HDV RNA strands has been demonstrated *in vivo*.^{141–147} HDV appears to require HBsAg for its hepatotropism and propagation but is not dependent on HBV for replication.^{133,147}

Pathogenesis of liver disease

The presence of replicating HDV seems to be invariably associated with liver damage suggesting that HDV may be directly hepatocytotoxic,¹³³ although immune-mediated mechanisms are also likely to be important in progression of the liver damage.^{148,149} A form of liver-kidney microsomal autoantibody similar to that found in a subset of autoimmune chronic active hepatitis has been described in patients with chronic HDV infection, but its significance is unclear.¹⁵⁰ The pattern of T cell infiltration in chronic HBV and HDV infection is also similar and this suggests a possible similarity in their pathogenic mechanisms.¹⁴⁹

Clinical features and laboratory diagnosis

Infection with HDV may occur simultaneously with HBV infection (co-infection), or as a superinfection in chronic HBsAg carriers. These two conditions have different clinical courses and outcomes.¹³³ Co-infection is usually self-limiting as acute HBV infection, although morbidity may be higher. However, a massive infecting dose may be associated with a more severe outcome, either in the form of fulminant hepatitis, or as a biphasic illness with initial improvement followed by relapse. In contrast, superinfection of HBsAg carriers with HDV may cause severe acute hepatitis (sometimes fulminant) and lead to chronic hepatitis and cirrhosis in over 70% of cases. What favours chronicity in HDV superinfection is not well understood but HDV superinfection is usually accompanied by a decrease in HBV replication markers.^{151,152} A subacute, rapidly progressive form of HDV superinfection has also been reported.¹⁵³

The diagnosis of different forms of HDV infection relies on the detection of HDAg in liver biopsies and serum HDAg, HDV RNA, and IgM and IgG anti-HD antibodies.¹³³ In acute HDV infection, HDV RNA and HDAg appear transiently in serum in around 25% of cases, IgM anti-HDV occurred in 93%, lasting from around 2–10 weeks, and this is usually followed in over 90% of patients by a persistent IgG anti-HD response. In those patients who develop chronic HDV infection, serum HDV RNA and IgM anti-HDV usually persists and HDAg can be found consistently in the liver. It has been suggested that there is a very good correlation between serum IgM anti-HDV and both intrahepatic HDAg and liver inflammatory activities.^{154,155} A recent study, however, showed that serum HDV RNA may also be detected in up to 32% of patients with chronic HDV infection negative for IgM anti-HDV.¹⁵⁶ *In situ* hybridization has also been employed recently to detect and localize HDV RNA in paraffin liver sections.¹⁵⁷ A riboprobe was also developed for

Table VII Prevalence of HDV infection

Country	HBsAg carriers + liver disease	Healthy HBsAg carriers	HBsAg positive i.v. drug users
<i>Europe</i>			
Britain		< 1%	42%
Ireland			31%
S Italy	14–28%		20–36%
Greece		2.4%	35%
Spain	9%		25–60%
Portugal	25%		85%
Belgium		2.8%	
Yugoslavia	7%		
Hungary	6%		
Romania	83%		
Poland	6.6%	1.6%	
Turkey	25%	17%	
USSR (Europe)	14–28%	3%	
<i>Africa</i>			
Tunisia	18%	10%	
Algeria	15%		
Ethiopia		5.8%	
S Africa		0.6%	
<i>Asia</i>			
USSR (Mid Asia)	41%	15%	
Israel	20%		
Saudi Arabia	53%	11%	
India (Bombay)	25%		
Taiwan	13%	2.2%	
China		1.8%	
<i>America</i>			
USA	25%	1.4–8%	20–53%
Amazon Basin	85–100%	23–35%	
Venezuela	91%	33%	
Columbia (S Marta)		29–36%	
Columbia (urban)		0%	
Argentina	2–5%	1.2%	
Chile	10–14%	1.8%	
<i>Australasia</i>			
Australia	19%	0.5–8%	19%
West Pacific		0–31%	

better detection of serum HDV RNA.¹⁵⁸ It is also important to note that in chronic HBsAg carriers with HDV superinfection the suppression of HBV replication may lead to a transient absence of HBV markers in the serum and liver;^{151,152} unless hepatitis D markers are sought, the diagnosis of NANB hepatitis may be made erroneously.

Anti-viral therapy and immunisation

Interferon-alpha has been shown to have an inhibitory effect on HDV replication but beneficial effects appear to be only transient in most cases.^{136,159–161} A preliminary report suggested that

fulminant HDV infection may respond to treatment with trisodium phosphonoformate. This, however, needs further studies for confirmation. Control of HDV infections is by vaccination against HBV.¹⁶²

Hepatitis E virus (HEV)

Epidemiology

The first and most massive outbreak of an enteric non-A, non-B epidemic was reported in 1955 when drinking water was contaminated by the overflow

of an open sewer in Delhi. A total of 29,300 residents developed an acute hepatitis, which ran a benign course in the majority and was self-limiting.¹⁶³ However, 10% of women, affected in their third trimester of pregnancy, died of fulminant hepatic failure.¹⁶⁴ The possibility of HAV, HBV, and other hepatotropic viral infection was excluded by paired viral serology. The disease displayed an unusual histological pattern with no chronic implications in subsequent follow-up studies.¹⁶⁵ These reports confirmed the existence of a faecal-orally transmitted type of non-A non-B hepatitis with completely different characteristics to blood borne NANB hepatitis. Of 10 Indian hepatitis epidemics subsequently investigated, 9 have been found to be due to epidemic/enteric NANB hepatitis (HEV).^{166,167}

Following the initial reports of HEV epidemic in India, similar epidemics were also reported in various Asian, African, Central and South American countries.¹⁶⁸ Sporadic imported cases of HEV have also been reported in the USA and it is likely that HEV is responsible for a small proportion of patients with acute sporadic NANB hepatitis seen in the UK and Europe.^{168,169} The attack rate has been estimated to be around 2% but is up to 19% in pregnant women, with the majority of attacks occurring in young adults.¹⁶⁷ This pattern of disease is compatible with an infection which is endemic but produces a sustained period of immunity. A reservoir of HEV is therefore required to sustain the infection in the absence of the carrier state. Recently, research workers in the USSR have been able to transmit HEV to pigs with induction of an hepatitis illness. In addition, they have also found anti-HEV antibody in rats from areas where the disease is endemic.

Identification of the virus

In 1983, 27–32 nm virus like particles (VLPs) were demonstrated by immune electron microscopy in the stools of 3 of 9 cases of HEV in Tashkent, USSR.¹⁷⁰ A volunteer then ingested a dilute suspension of stools pooled from the patients and he developed an acute hepatitis, with antibodies detectable to the VLP by immune electron microscopy (IEM).¹⁷⁰ A similar experiment was performed using the cynomolgus monkey. Subsequently,

HEV was also transmitted to a number of primates. In the majority of studies, 27–34 nm VLPs aggregated by autologous acute phase serum have been found in stool samples. Using IEM techniques, it was found that sera and VLPs from the different regions cross reacted, indicating that HEV in various parts of the world is associated with the same non-enveloped 27–34 nm VLP.^{171–176} Recently, an antigen related to HEV (HEV Ag) was found in the liver of experimentally infected macaques.¹⁷⁴ It has a granular distribution in up to 90% of hepatocytes in the early phase of experimental infection.

The recent successful partial molecular cloning of the viral genome of HEV by Reyes *et al.* represents a giant leap forward.¹⁷⁷ Briefly, bile juice from infected cynomolgus macaques, which contained high virus particle count on electron microscopy, was used for the construction of recombinant complementary DNA (cDNA) libraries. Libraries in bacteriophage lambda gt 10 were screened by differential hybridization with cDNA probes prepared from RNA extracted from infected and uninfected bile preparations. One molecular clone was found to hybridize with a roughly 7.6 kilobase RNA species present only in the livers of infected cynomolgus macaques but not to any of the control DNA and RNA extracts. The translated nucleic acid sequence contains a consensus amino acid sequence consistent with an RNA-directed RNA polymerase, an enzyme that is present in all positive strand RNA viruses (Figure 6). Further studies along this line will certainly be important in taxonomic classification and understanding of the virus.¹⁷⁸

Clinical features and laboratory diagnosis

The majority of clinical data comes from the well characterized Indian epidemics.¹⁶⁷ The incubation period was between 3–9 weeks. The attack rate was higher in young adults (2.9%) than children (1.4%) or those aged over 40 (2.0%). A brief prodrome illness with anorexia, nausea, vomiting and abdominal pain is followed by jaundice. The disease is usually benign and self-limiting with no chronic sequelae. However, fulminant hepatic failure has been reported to occur in up to 2.8% of men and 22% of pregnant women in whom it is usually

Figure 6 Schematic alignment of the known putative domains of the hepatitis E virus. POL – RNA-directed RNA polymerase; IRE – immunoreactivity epitope.

fatal.¹⁷⁹ This severe form of the disease in pregnancy, particularly the third trimester is a feature of all HEV epidemics.

Histologically, cholestasis was prominent in around 50% of the patients with marked canalicular and intracellular bile stasis in glandular channels.¹⁸⁰ Moderate inflammatory infiltrates of mononuclear and polymorphonuclear leukocytes occurred in the portal and intralobular regions with prominent lipofuscin pigment in Kupffer cells.¹⁸⁰ With the recent success in identification of HEV-Ag in the liver sections of experimentally infected primates and the development of fluorescent linked anti-HEV Ag, an antibody blocking assay was developed as a prototype test for the identification and titration of specific anti-HEV Ag antibody in the serum samples.¹⁷⁴ This test has reconfirmed the immune electron microscopy findings that the enteric non-A, non-B hepatitis outbreaks in Asia, Africa and North America were found to be caused by HEV. Nevertheless, this test is still only available in specialized centres.¹⁷⁴

Treatment and control

As is the case with HAV, treatment is purely supportive. Control of the disease relies on the well established primary health objectives of clean water and adequate sanitation. Adaptation of the virus to cell culture or synthesis of viral antigens by recombinant technology in the near future may enable more simple diagnostic tests to be established and to facilitate the development of vaccines.

Hepatitis X virus (X = F, G, . . .)

There are at least two more types of hepatotropic viruses as determined by epidemiology, molecular biology and electron microscopy studies. The first is the short incubation blood-borne non-A, non-B hepatitis virus. It is quite uncommon and may not be epidemiologically important. Based upon the limited data available, this second, or minor, non-A, non-B virus is thought to be smaller than the HCV (around 25–30 nm), naked rather than enveloped, and incapable of producing the characteristic cytoplasmic tubular structures found by electron microscopy in the hepatocytes of chimpanzees experimentally infected with HCV. To date, no antibody specific for this second non-A, non-B hepatitis has been detected. The second is the sporadic, acute, non-transfusion transmitted non-A, non-B hepatitis observed in Western countries (USA, Italy). It is serologically unrelated to the well characterized HEV outbreaks in the developing countries. The aetiological agent for these sporadic cases remains to be identified.

Fulminant hepatic failure

Fulminant hepatic failure (FHF) is the term used to describe patients with a rapid clinical progression and is defined as liver failure where the onset of encephalopathy is within 8 weeks of the first symptoms.¹⁸¹ The most common cause of FHF in the UK is still paracetamol, usually taken in overdose although hepatotoxicity has occurred following therapeutic doses of paracetamol in patients taking enzyme inducing drugs¹⁸² and in those with a chronic high alcohol intake.¹⁸³ It is well known that N-acetylcysteine will prevent paracetamol-induced hepatotoxicity, in most cases, if given within 8 hours of the overdose. It is less efficacious after this time and it fails to avert the development of severe hepatotoxicity if given more than 15 hours after overdose by the intravenous route¹⁸⁴ and 24 hours after overdose by the oral route.¹⁸⁵

It has always been assumed that once hepatic necrosis develops prior treatment with N-acetylcysteine has no further influence on outcome. However, even if N-acetylcysteine does not prevent the development of hepatic necrosis it can still confer a later benefit. Recent work from this unit has shown that patients who develop paracetamol-induced fulminant hepatic failure have a better prognosis if they have been given N-acetylcysteine after overdose (even up to 36 hours) in comparison to those who never received it.¹⁸⁶ Despite having a similar disturbance in liver function, as assessed by prolongation of prothrombin time, patients who received N-acetylcysteine had a lower incidence of renal failure and fewer progressed to deep grades of encephalopathy, suggesting that N-acetylcysteine therapy had a later protective effect on other organs. A prospective controlled trial of N-acetylcysteine in patients with paracetamol-induced fulminant hepatic failure has just been completed, confirming its beneficial role not only late after overdose but also at a time when coagulopathy and encephalopathy have already developed.¹⁸⁷

The mechanism of action of N-acetylcysteine at this time is unclear but it has recently been shown that an N-acetylcysteine infusion improves systemic haemodynamics and enhances oxygen delivery and consumption in fulminant hepatic failure.^{187a} This could be beneficial as prospective studies have shown that optimizing oxygen transport reduces tissue hypoxia and improves outcome in multi-organ failure due to other causes.¹⁸⁸ Whatever the mechanism of action no other new medical therapy has had such an impact on the survival from paracetamol-induced fulminant hepatic failure in recent years.

The pathogenic mechanisms causing hepatic encephalopathy and brain oedema in fulminant hepatic failure remain incompletely understood.

Apparent amelioration of encephalopathy in animal models and in uncontrolled clinical studies by the intravenous administration of the benzodiazepine receptor antagonist, flumazenil, raises the possibility that an endogenous benzodiazepine receptor ligand with agonist properties may contribute to the manifestations of hepatic encephalopathy by allosterically potentiating GABA-mediated neurotransmission.¹⁸⁹ However, data from controlled studies are lacking and although the putative benzodiazepine agonists have been identified in brain in animal models¹⁹⁰ they remain elusive in man. Evidence for the role of toxins, such as ammonia, as causative agents in cerebral oedema continues to accumulate.¹⁹¹ Brain oedema is now known to be the cause of the raised intracranial pressure seen in deep grades of encephalopathy. It had been thought that high cerebral blood flow was a contributing factor to raised intracranial pressure but it has now been shown that cerebral blood flow is low in fulminant hepatic failure and inversely proportional to the grade of encephalopathy.¹⁹² The practice of nursing patients in the head upright posture to reduce cerebral blood flow has, consequently, been re-examined. It has been found that a posture of head-up greater than 20 degrees adversely effects cerebral perfusion pressure¹⁹³ and, therefore, could lead to cerebral ischaemia from hypoperfusion; it should now be the practice to nurse patients at risk of developing cerebral oedema flat in bed. Hyperventilation producing hypocarbia can reduce cerebral blood flow further and can cause cerebral ischaemia,¹⁹⁴ it should be abandoned as a treatment for cerebral oedema complicating fulminant hepatic failure. Mannitol remains the most effective therapy for cerebral oedema.¹⁹⁵ Thiopentone infusion may be a useful adjunct in patients with cerebral oedema unresponsive to other forms of therapy.¹⁹⁶

Renal failure is frequently seen in combination with cerebral oedema as a complication of fulminant hepatic failure. It is known that the treatment of renal failure with machine driven intermittent dialysis can lead to an increase in the intracranial pressure,¹⁹⁷ however, continuous arteriovenous haemodiafiltration (CAVHD) does not appear to be associated with surges in intracranial pressure and it is suggested that CAVHD is now the treatment of choice for patients with fulminant hepatic failure complicated by oliguric renal failure who are at risk of developing cerebral oedema.¹⁹⁸ Intermittent machine dialysis still has a role in the treatment of those patients usually post-paracetamol overdose who get predominantly renal failure and only mild liver damage.

Bacterial¹⁹⁹ and fungal infection²⁰⁰ with commensal organisms has been found to be a common and often unrecognized complication of fulminant

hepatic failure. Prophylactic antimicrobial decontamination of the nasopharynx and gut at the onset of liver failure might be effective in reducing the incidence of later infection. The results of controlled trials are eagerly awaited.

Liver regeneration

Liver growth is the key factor in recovery from fulminant hepatic failure and inadequate regeneration is thought to contribute to the high mortality seen in some aetiological variants, such as non-A non-B viral or drug-induced hepatitis.²⁰¹ Work with hepatocyte cultures and animal studies have now identified a number of factors that are able to stimulate hepatocyte DNA synthesis and mitosis in otherwise quiescent cells. Epidermal growth factor (EGF) was the first substance to be shown to have this effect²⁰² but whether it has a physiological role in liver regeneration is unclear. Transforming growth factor α (TGF α) shares the same receptor as EGF and is also mitogenic for hepatocytes. It has recently been shown that regenerating hepatocytes produce TGF α and it may be that the secretion of TGF α constitutes an autocrine loop.²⁰³ Human hepatocyte growth factor has recently been identified as a potent and specific new hepatotrophic factor and serum levels in fulminant hepatic failure appear to be high, irrespective of the aetiology or outcome.²⁰⁴ No deficiency of any of the known stimulators of liver growth has been found in fulminant hepatic failure and it is, therefore, unlikely that administration of these factors to patients will improve their liver regeneration response. Indeed, administration of insulin and glucagon, which are permissive hepatotrophic factors, to patients with fulminant hepatic failure in a controlled trial did not stimulate regeneration nor improve outcome.²⁰⁵ Circulating inhibitors of regeneration have been postulated in fulminant hepatic failure. Interleukins 1 and 6 have been shown to inhibit hepatocyte DNA synthesis and proliferation. Perhaps this effect reflects their known actions on directing hepatocytes toward an acute phase response which then takes precedence over the synthetic processes required for liver cell growth.²⁰⁶

Liver transplantation

Although advances in the medical management of patients with fulminant hepatic failure are being made, allowing time for liver regeneration to occur, it is known that in some patients liver function will never recover with medical treatment alone. These patients need to be identified quickly because liver transplantation is now a viable and successful

treatment option^{207–213} with a one year survival of 55% in the first 50 patients with acute liver failure transplanted at King's College Hospital. Universally applicable criteria for the selection of patients are needed to prevent unnecessary transplantation and to ensure that patients who may benefit are appropriately treated. On our unit, an aetiology of non-A non-B or drug-induced hepatitis, time from jaundice to onset of encephalopathy > 7 days, age < 10 or > 40 years, serum bilirubin > 300 $\mu\text{mol/l}$ and prothrombin time > 50 seconds have been found to correlate with poor prognosis, whilst pH < 7.3 and prothrombin time > 100 seconds with creatinine > 300 mmol/l and coma grade $\geq$ III are important predictors of outcome in paracetamol-induced fulminant hepatic failure.²¹⁴ In the latter group serial measurements of prothrombin time²¹⁵ and the ratio of the individual clotting factors, V and VIII, may provide a further refinement. However, these prognostic indicators have yet to find universal acceptance and a multi-centred prospective assessment of their ability to select patients for transplantation is required. Orthotopic grafting is currently the accepted technique for liver transplantation in fulminant hepatic failure but heterotopic grafting has potential long term benefits.²¹⁶ A heterotopic grafted liver could restore hepatic function, allowing the patient to recover and the diseased liver time to regenerate; once host liver function is re-established immunosuppression could be withdrawn allowing the grafted liver to atrophy and the patient to remain off long term immunosuppressive drugs. The introduction of the University of Wisconsin Solution to preserve the donor liver, now allows safe storage of the grafts for up to 24 hours compared with 6–8 hours previously.²¹⁷

Liver transplantation in hepatitis B virus-related diseases

On a worldwide basis, chronic hepatitis B virus infection is potentially one of the most common indicators for liver transplantation, either because of end-stage liver disease or the development of hepatocellular carcinoma. Recurrence of HBV infection is another factor contributing to morbidity and mortality after transplantation in this group of patients. The one and five year survival in patients with HBV in the Pittsburg series were 57% and 40% respectively,²¹⁸ the two year survival in the King's/Cambridge series is 45%.²¹⁹ In the King's/Cambridge series of patients transplanted for chronic HBV infection-related liver failure, 84% of the patients became positive for HBsAg at 4 to 6 weeks post-transplant, probably because of maintained viral replication in extra-hepatic sites.^{219–221} Virtually, all patients with evidence of active viral replication prior to transplantation developed

HBV infection in the graft. This was also found in 64% of patients without positive serology pre-transplant. Only 5 patients showed either no recurrence or spontaneously clearing HBV in association with an acute hepatitis illness.

This high recurrence rate, particularly in those with no evidence of active viral replication prior to transplantation, has three major implications. Firstly, the source virion may be extra-hepatic and this is consistent with a series of studies indicating that intact HBV DNA (but not replicating virus) is present in peripheral blood mononuclear cells and bone marrow cells.^{220,221} Secondly, modern technology, like polymerase chain reaction, may help in the detection of very low levels of serum HBV DNA. Thirdly, as patients with negative serology so frequently have their grafts reinfected, there is little point in treating with anti-viral agents like interferon-alpha pre- or peri-operatively since this is unlikely to reduce the rate of transmission of HBV and also may be at the expense of peri-operative neutropenia, thrombocytopenia, and graft rejection.

The implications for the HBV reinfected graft are variable, ranging from normal histology to loss of graft at 4–13 months.^{219,222} The latter is usually the result of a rapidly progressive syndrome with a histological lesion of peri-sinusoidal bands of fibrosis extending from the portal tracts to surrounding plates of ductal type epithelium, with prominent cholestasis, and massive cytoplasmic expression of viral antigens, particularly HBsAg¹⁸⁵ (Lau *et al.*, unpublished data). This syndrome, which has been termed fibrosing cholestasis hepatitis (FCH), presents as jaundice together with a relentless prolongation of prothrombin time, but with only moderate increase in serum transaminases, leading to loss of the graft within 4–6 weeks of onset.²¹⁹

The similarity between this pattern of damage and that seen in the transgenic mouse model, where massive HBsAg expression was associated with hepatocyte damage and the relative absence of inflammatory cells, suggests that in this unique syndrome, damage is not immune-mediated but related to massive expression of viral protein.^{65,66,222} Glucocorticoid steroid has been shown to have a direct pro-viral effect by acting through a glucocorticoid responsive element on HBV DNA.^{222,224} Using primary hepatocyte cultures, we have recently demonstrated an increase in intracellular HBsAg production modulated by prednisolone; azathioprine and cyclosporin A have no direct viral modulating effect (Lau *et al.*, unpublished data). Observations in HBsAg positive patients with malignant lymphoma treated with intensive chemotherapy suggested that immunosuppression may be associated with reactivation of viral replication.^{225,226} Interestingly, co-existing delta infection

appears to be associated with less aggressive liver disease after transplantation.^{219,222} According to the Paris and Munich groups, HBV reinfection and its effects on the grafts appear to be reduced by long-term immuno-prophylaxis with immuno-globulin combined with active immunization, particularly when the patients are in a non- or low-replicative state at the time of transplantation.^{227,228} They also recommend restriction of liver transplantation to this category of patients only. This HB-immunoglobulin therapy has not been standardized and the initial peri-operative doses used ranged from 500–128,000 units. Long-term therapy is adjusted to maintain serum HB-immunoglobulin levels of more than 100 IU/l, and must be carried on indefinitely, thus adding considerably to the transplant procedure. The anticipated availability of human monoclonal antibody preparation of very high potency to the HBV envelope proteins in the near future may alter the outlook of these patients.

Transplantation for hepatic malignancy

The initial enthusiasm for recommending patients with primary hepatic malignancy for liver transplantation has declined, as it has become apparent that the rate of tumour recurrence is high, despite careful screening prior to transplantation for macroscopic extrahepatic spread. The size of a hepatocellular carcinoma at time of liver grafting appears to be the critical factor; in the King's series, the incidence of tumour recurrence was 0% for tumours up to 4 cm in diameter, 44.4% for tumours 5–8 cm in diameter and 68% for larger and multifocal tumours (J. O'Grady, personal communication). Patients transplanted for cholangiocarcinoma complicating primary sclerosing cholangitis whether diagnosed pre-operatively or at the time of surgery are considered to have a poor prognosis.²²⁹ Outcome following transplantation for hepatic secondary tumours is not encouraging,²³⁰ except perhaps for palliation in APUDomas, including carcinoid tumours.²³⁰

Immunosuppression for liver transplantation

Cyclosporin A has been a major factor in the improvement of results of liver transplantation during the past decade.²³¹ However, its nephrotoxicity and other limitations have stimulated a continuing search for alternative agents. The publication in the *Lancet* of the impressive results of the first clinical experience of the potent new immunosuppressant FK-506 in human liver transplantation by T.E. Starzl and his colleagues has aroused much interest.²³²

FK-506 is an 822 kD macrolide antibiotic extracted from the fermentation of the broth of the

soil fungus *Streptomyces tsukubaensis*. Although totally different in structure from the cyclic peptide cyclosporin A, it exhibits similar, if not identical selective anti-lymphocytic activity. The immunosuppressive potency of FK-506, however, is very much greater.^{233,234} In the study reported by Starzl *et al.*, FK506 has the capacity to rescue critically ill patients with both acute and chronic rejection that was resistant to anti-lymphocyte globulin (OKT3) or steroid therapy.²³² In those liver graft recipients receiving FK506, a significant reduction of severe viral infection was observed compared with the control group which received the conventional immunotherapy. In particular, there had been no symptomatic cytomegalovirus (CMV) infection with FK506.

Despite the fact that they are structurally distinct, the combined use of FK506 and cyclosporin should be avoided since an interaction between the two agents leading to marked elevation of cyclosporin A blood levels and exacerbation of renal impairment has been encountered.^{232,235} Multicentre controlled trials are currently being undertaken in the USA and UK to compare these two agents.

Tissue typing and liver transplantation

It is established that matching the graft to the recipients ABO and Rhesus blood groups reduces the risk of hyperacute rejection caused by pre-formed antibodies. Although, transplantation across the blood group is occasionally performed in patients with acute liver failure when an appropriate donor of the same blood group can not be found. The need to match class I and II major histocompatibility antigens is less clear. Presently there is insufficient evidence to suggest that HLA matching would be associated with improved graft survival and there are some indications that selective mismatches may be advantageous in reducing the incidence of chronic rejection.²³⁶

The management of portal hypertension

In addition to the elevated intrahepatic resistance to portal blood flow, portal hypertension is maintained by an increase in portal venous flow. It has been postulated that the hyperdynamic circulation develops as a consequence of enhanced production of endogenous vasodilators²³⁷ and a number of potential agents is given in Table VIII. Many of these vasodilators act via endothelium derived relaxing factor (EDRF)²³⁸ and increased production of EDRF itself has been proposed as the mediator of the circulatory abnormality in cirrhosis.²³⁹ Antagonists of EDRF synthesis are available but their effects in animal models of cirrhosis

Table VIII Vasodilators implicated in the pathophysiology of the hyperdynamic circulation in cirrhosis

Glucagon
Cholecystokinin
Vasoactive intestinal polypeptide
PGE ₂ , LTB ₁
Purines
Adenosine
Enkephalin
Gamma-amino butyric acid
Bradykinin
Endothelium-derived relaxing factor

need to be assessed before clinical studies commence as one of these agents, L-mono methyl arginine, increases mortality in rats given endotoxin.²³⁹

The management of portal hypertension is still centred on the primary and secondary prevention of variceal haemorrhage as the latter continues to contribute significantly to the morbidity and mortality of chronic liver disease. Several randomized controlled trials of the effects of β -blockers in the primary prevention of variceal haemorrhage have been published and a formal meta-analysis performed on their results has shown that β -blockade does reduce the incidence of the first bleed.²⁴⁰ However, the administration of β -blockers does

not improve outcome as the dominant influence on survival is liver function. The current consensus is against the use of sclerotherapy for primary prophylaxis, whereas emergency sclerotherapy is the mainstay in the treatment of patients bleeding from oesophageal varices.²⁴¹

Endoscopic ligation of oesophageal varices, which effects strangulation of varices using elastic 'O' rings, offers a new approach to the treatment of variceal bleeding²⁴² but controlled trials are required to assess its potential. Gastric varices account for up to 20% of variceal bleeding. High lesser-curve varices may be amenable to sclerotherapy and, although bleeding from fundal varices usually requires surgery, some success has been achieved injecting the tissue adhesive bucrylate.²⁴³ When sclerotherapy is not immediately available balloon tamponade with a Sengstaken-Blakemore tube can afford temporary control of bleeding but its use is associated with a high incidence of side effects, such as aspiration, oesophageal ulceration and perforation.²⁴⁴ In this situation we recommend pharmacological therapy with a combination of intravenous vasopressin and transdermal nitroglycerin.²⁴¹ If a patient rebleeds after sclerotherapy has been performed on two occasions already, then surgery is indicated.²⁴⁵ The potential difficulties for subsequent orthotopic liver transplantation should be considered in all patients referred for operative intervention.

References

Hepatitis A Virus

1. Feinstone, S.M., Kapikian, A.Z. & Purcell, R.H. Hepatitis A: detection by immune electron microscopy of a virus like antigen associated with acute illness. *Science* 1973, **182**: 1026–1028.
2. Siegel, G. Virology of hepatitis A. In: Zuckerman, A.J. (ed) *Viral Hepatitis and Liver Disease*. Alan R Liss, New York, 1988, pp. 3–7.
3. Cohen, J.L., Ticehurst, J.R., Purcell, R.H., Buckler-White, A. & Baroudy, B.M. Complete nucleotide sequence of wild-type hepatitis A virus: comparison with different strains of hepatitis A virus and other picornaviruses. *J Virol* 1987, **61**: 50–59.
4. Anderson, D.A., Locarnini, S.A., Ross, B.C., Coulepis, A.G., Anderson, B.N. & Gust, I.D. Single-cycle growth kinetics of hepatitis A virus in BSC-1 cells. In: Britton, W.A., Rueckert, R.R. (eds) *Positive Strand RNA Virus*. Alan R Liss, New York, 1986, pp. 495–507.
5. Weitz, M. & Siegel, G. Variation among hepatitis A virus strains. Genomic variation detected by oligonucleotide mapping. *Virus Res* 1985, **4**: 53–67.
6. Ticehurst, J.R. Hepatitis A virus: clones, culture and vaccines. *Semin Liver Dis* 1986, **6**: 46–55.
7. Huges, J.V., Stanton, L.W., Tomassini, J.E., Long, W.J. & Scolnick, E.M. Neutralising monoclonal antibodies to hepatitis A virus: partial purification of a neutralising antigenic site. *J Virol* 1984, **52**: 465–473.
8. Stapleton, J.T. & Lemon, S.M. Neutralisation escape mutants define a dominant immunogenic neutralisation site on hepatitis A virus. *J Virol* 1987, **61**: 491–498.
9. Provost, P.F. *In vitro* propagation of hepatitis A virus. In: Gerety, R.J. (ed) *Hepatitis A*. Academic Press, New York, 1984, pp. 245–262.
10. Siegel, G. The biochemistry of hepatitis A virus. In: Gerety, R.J. (ed) *Hepatitis A*. Academic Press, New York, 1984, pp. 9–32.
11. Vallbracht, A., Hofmann, L., Wurster, K.G. & Flehming, B. Persistent infection of human fibroblasts by hepatitis A virus. *J Gen Virol* 1984, **65**: 609–615.
12. Gauss-Muller, V. & Dienhardt, F. Effect of hepatitis A virus infection on cell metabolism *in vitro*. *Proc Soc Exp Biol Med* 1984, **175**: 10–15.
13. Venuti, A., Di Russo, C., del Grosso, N. *et al*. Isolation and molecular cloning of a fast growing strain of human hepatitis A virus from its double stranded replicative form. *J Virol* 1985, **56**: 579–588.
14. Divizia, M., Ruggeri, F., Hentschel, J., Flehming, B., Pana, A. & Perez-Bercoff, R. Preliminary characterisation of a fast growing strain of human hepatitis A virus. *Microbiologica* 1986, **9**: 269–278.
15. Anderson, D.A., Cytopathicity, plaque assay and heat inactivation of hepatitis A virus strain HM175. *J Med Virol* 1987, **22**: 35–44.
16. Slusarczyk, J., Hannon, B.G., Nordenfelt, E., Krawczynski, K., Karwowska, S. & Knap, J. Etiopathogenic aspects of hepatitis. Excretion of hepatitis A virus, biochemistry of liver function, and humoral immune response in patients with hepatitis A on admission to hospital. *J Med Virol* 1985, **15**: 105–116.

17. Gabriel, P., Vallbracht, A. & Flehmig, B. Lack of complement dependent cytolytic antibodies in hepatitis A virus. *J Med Virol* 1986, **20**: 23–31.
 18. Vallbracht, A., Gabriel, P., Zahn, J. & Flehmig, B. Hepatitis A virus infection and the interferon system. *J Infect Dis* 1985, **152**: 211–213.
 19. Kurane, I., Binn, L.N., Bankroft, W.H. & Ennis, F.A. Human lymphocyte response to hepatitis A virus-infected cells: interferon production and lysis of infected cells. *J Immunol* 1985, **135**: 2140–2144.
 20. Vallbracht, A., Gabriel, P., Maier, K. *et al.* Cell-mediated cytotoxicity in hepatitis A virus infection. *Hepatology* 1986, **6**: 1308–1314.
 21. Gust, I.D. & Feinstone, S.M. Epidemiology. In: *Hepatitis A*. CRC Press, Florida, 1988, pp. 163–191.
 22. Lemon, S.M. Type A viral hepatitis. New developments in an old disease. *N Engl J Med* 1985, **313**: 1059–1067.
 23. Forbes, A. & Williams, R. Changing epidemiology and clinical aspects of hepatitis A. In: Zuckerman, A.J. (ed) *British Medical Bulletin: Viral Hepatitis*. Churchill Livingstone, London, 1990, pp. 303–318.
 24. Yang, N.Y., Yu, P.H., Mao, Z.X., Chen, N.L., Chai, S.A. & Mai, J.L. Inapparent infection of hepatitis A virus. *Am J Epidemiol* 1988, **127**: 599–604.
 25. Gorden, S.C., Reddy, K.R., Schiff, L. & Schiff, E.R. Prolonged intrahepatic cholestasis secondary to acute hepatitis A. *Ann Intern Med* 1984, **101**: 635–637.
 26. Sjogren, M.H., Tanno, H., Fay, O. *et al.* Hepatitis A virus in stool during clinical relapse. *Ann Intern Med* 1987, **106**: 221–226.
 27. Gust, I.D. & Feinstone, S.M. Clinical features. In: *Hepatitis A*. CRC Press, Florida, 1988, pp. 145–162.
 28. Lemon, S.M., Brown, C.D., Brooks, D.S., Simms, T.E. & Bancroft, W.H. Specific immunoglobulin M response to hepatitis A virus determined by solid-phase radioimmunoassay. *Infect Immunol* 1980, **28**: 927–936.
 29. Gust, I.D. & Feinstone, S.M. General preventive measures. In: *Hepatitis A*. CRC Press, Florida, 1988, pp. 163–191.
 30. Cossar, J.H. & Reid, D. Not all travellers need immunoglobulin for hepatitis A. *Br Med J* 1987, **294**: 1503.
 31. Gust, I.D. Design of hepatitis A vaccines. In: Zuckerman, A.J. (ed) *British Medical Bulletin: Viral Hepatitis*. Churchill Livingstone, London, 1990, pp. 319–328.
 32. Flehmig, B., Heinrich, U. & Pfisterer, M. Immunogenicity of a recombinant hepatitis A vaccine in seronegative volunteers. *Lancet* 1989, **i**: 1039.
 33. Feinstone, S.M., Daemer, R.J., Gust, I.D. *et al.* Liver attenuated vaccine for hepatitis A. *Dev Biol Stand* 1983, **54**: 429–432.
- Hepatitis B Virus**
34. Maynard, J.E., Kane, M.A., Alter, M.J. & Hadler, S.C. Control of hepatitis B by immunization: global prospective. In: Zuckerman, A.J. (ed) *Viral Hepatitis and Liver Diseases*. Alan R Liss, New York, 1988, pp. 967.
 35. *Bull WHO* 1988, **66**: 443.
 36. Michel, M.L. & Tiollais, P. Structure and expression of the hepatitis B virus genome. *Hepatology* 1987, **7**: 61S–63S.
 37. Miller, R.H., Kaneko, S., Chung, C.T., Girones, R. & Purcell, R.H. Compact organisation of the hepatitis B virus genome. *Hepatology* 1989, **9**: 322–327.
 38. Leenders, W.P.J., Glangsbeck, H.J., de Bruin, W.C.C. & Yap, S.H. Binding of the major and large HBsAg to human hepatocytes and liver plasma membranes: putative external and internal receptors for infection and secretion of hepatitis B virus. *Hepatology* 1990, **12**: 141–147.
 39. Imai, M., Yanase, Y., Nojiri, T. *et al.* A receptor for polymerised human and chimpanzee albumins on hepatitis B virus particles co-occurring with HBsAg. *Gastroenterology* 1979, **76**: 242–247.
 40. Machida, A., Kishimoto, S., Ohnuma, H. *et al.* A hepatitis B surface antigen polypeptide (P31) with the receptor for polymerised human as well as chimpanzee albumins. *Gastroenterology* 1983, **85**: 268–274.
 41. Machida, A., Kishimoto, S., Ohnuma, H. *et al.* A polypeptide containing 55 amino acid residues coded by the pre-S region of hepatitis B virus DNA, bears the receptor for polymerised human as well as chimpanzee albumins. *Gastroenterology* 1984, **86**: 910–918.
 42. Thung, S.N. & Gerber, M.A. Polyalbumin receptors: their role in the attachment of hepatitis B virus to hepatocytes. *Semin Liver Dis* 1984, **4**: 69–75.
 43. Thung, S.N. & Gerber, M.A. Albumin binding sites of human hepatocytes. *Liver* 1983, **3**: 290–294.
 44. Trevisan, A., Gudat, F., Guggenheim, R. *et al.* Demonstration of albumin receptors on isolated human hepatocytes by light and scanning electron microscopy. *Hepatology* 1982, **2**: 832–835.
 45. Heermann, K.H., Waldeck, F. & Gerlich, W.H. Interaction between native human serum and the pre-S2 domain of HBsAg. In: Zuckerman, A.J. (ed) *Viral Hepatitis and Liver Disease*. Alan R Liss, New York, 1988, pp. 697.
 46. Yu, M.W., Finlayson, J.S. & Shih, J.W.K. Interaction between various polymerised human albumins and HBsAg. *J Virol* 1985, **55**: 736–743.
 47. Krone, B., Lenz, A., Heermann, K.H., Seifer, M., Lu, X.Y. & Gerlich, W.H. Interaction between hepatitis B surface proteins and monomeric serum albumin. *Hepatology* 1990, **11**: 1050–1056.
 48. Vento, S., Chen, S.H., Giuliani-Piccarri, G. *et al.* Cellular immunity to nucleocapsid and pre-S determinants in asymptomatic carrier of hepatitis B virus. *Immunology* 1987, **62**: 593–798.
 49. Vento, S., Hergarty, J.E., Alberti, A. *et al.* T lymphocyte sensitisation to HBcAg and T cell mediated unresponsiveness to HBsAg in hepatitis B related chronic liver disease. *Hepatology* 1985, **5**: 192–197.
 50. Mondelli, M., Mieli-Vergiani, G., Alberti, A. *et al.* Specificity of T lymphocyte cytotoxicity to autologous hepatocytes in chronic HBV infection: evidence that T cells are directed against HBcAg expressed on hepatocytes. *J Immunol* 1982, **129**: 2773–2778.
 51. Naoumov, N.V., Mondelli, M., Alexander, G.J.M. *et al.* Relationship between expression of hepatitis B virus antigens in isolated hepatocytes and autologous lymphocyte cytotoxicity in patients with chronic hepatitis B virus infection. *Hepatology* 1984, **4**: 63–68.
 52. Mondelli, M., Bortolotti, F., Pontisso, P. *et al.* Definition of HBV specific target antigens recognised by cytotoxic T cells in acute HBV infection. *Clin Exp Immunol* 1987, **68**: 245–250.
 53. Ferrari, C., Penna, A., Giuberti, T. *et al.* Intrahepatic nucleocapsid antigen specific T cells in chronic active hepatitis B. *J Immunol* 1987, **139**: 2050–2058.
 54. Ferrari, C., Mondelli, M.Y., Penna, A. *et al.* Functional characterisation of cloned intrahepatic HBV nucleoprotein specific helper T cell lines. *J Immunol* 1987, **139**: 539–544.
 55. Ou, J., Laub, O. & Rutter, W.J. Hepatitis B virus gene function: the precore region targets the core antigen to cellular membranes and causes the secretion of the e antigen. *Proc Natl Acad Sci USA* 1986, **83**: 1578–1582.
 56. Uy, A., Bruss, V., Gerlich, W.H., Kochel, H.G. & Thomsen, R. Precore sequence of hepatitis B virus inducing e antigen and membrane association of the viral core protein. *Virology* 1986, **155**: 89–96.
 57. Feitelson, M.A. Products of the 'X' gene in hepatitis B and related viruses. *Hepatology* 1986, **6**: 191–198.
 58. Katayama, K., Hayashi, N., Sasaki, Y. *et al.* Detection of hepatitis B virus X gene protein and antibody in type B chronic liver disease. *Gastroenterology* 1989, **97**: 990–998.

59. Feitelson, M.A. & Clayton, M.M. X antigen/antibody markers in hepadnavirus infections. *Gastroenterology* 1990, **99**: 500–507.
60. Ayoola, E.A. Viral hepatitis in Africa. In: Zuckerman, A.J. (ed) *Viral Hepatitis and Liver Disease*. Alan R Liss, New York, 1988, pp. 161–169.
61. Carman, W.F., Jacyna, M.R., Hadziyannis, S. *et al.* Mutation preventing formation of hepatitis B e antigen in patients with chronic hepatitis B infection. *Lancet* 1989, **ii**: 588–591.
62. Brunetto, M.R., Stemler, M., Bonino, F. *et al.* A new hepatitis B virus strain in patients with severe anti-HBe positive chronic hepatitis B. *J Hepatol* 1990, **10**: 258–261.
63. Fagan, E.A., Davison, F., Trowsbridge, R. *et al.* Detection of HBV-DNA sequences in liver in HBsAg sero-negative patients with liver disease with or without anti-HBe antibodies. *Hepatology* 1989, **9** (suppl): abstr 26.
64. Trent-Mills, C., Lee, E. & Perrillo, R. Relationship between histology, aminotransferase levels, and viral replication in chronic hepatitis B. *Gastroenterology* 1990, **99**: 519–524.
65. Chisari, F.V., Filippi, P., Buras, J. *et al.* Structural and pathological effects of synthesis of hepatitis B virus large envelope polypeptide in transgenic mice. *Proc Natl Acad Sci USA* 1987, **84**: 6909–6913.
66. Moriyama, T., Guilhot, S., Klopchin, K. *et al.* Immunobiology and pathogenesis of hepatocellular injury in hepatitis B virus transgenic mice. *Science* 1990, **248**: 361–364.
67. Alexander, G.J.M. Immunology of hepatitis B virus infection. In: Zuckerman, A.J. (ed) *British Medical Bulletin: Viral Hepatitis*. Churchill Livingstone, London, 1990, pp. 354–367.
68. Mieli-Vergiani, G., Vergiani, D., Portmann, B. *et al.* Lymphocyte cytotoxicity to autologous hepatocyte in HBsAg positive chronic liver disease. *Gut* 1982, **23**: 1029–1036.
69. Ikeda, T., Lever, A.M.L. & Thomas, H.C. Evidence for a deficiency of interferon production in patients with chronic hepatitis B virus acquired in adult life. *Hepatology* 1986, **6**: 962.
70. Zachoval, R., Abb, J., Zachoval, V. *et al.* Interferon alpha in hepatitis B and non-A, non-B defective production by peripheral blood mononuclear cells in chronic infection and development of serum interferon in acute disease. *J Hepatol* 1988, **6**: 364–368.
71. Nouri-Aria, K.T., Magrin, S., Alexander, G.J.M., Anderson, M.G., Williams, R. & Eddleston, A.L.W.F. Abnormal T cell activation in chronic hepatitis B virus infection: a consequence of monocyte activation. *Immunology* 1988, **64**: 733–738.
72. Alexander, G.J.M., Nouri-Aria, K.T., Neuberger, J. *et al.* *In vitro* effects of lymphoblastoid interferon on lymphocyte activation and cell-mediated cytotoxicity in patients with chronic hepatitis B virus infection. *J Hepatol* 1986, **3** (Suppl. 2): S269.
73. Twu, J.S., Lee, C.H., Lin, M.P. & Schloemer, R.H. Hepatitis B virus suppresses expression of human interferon. *Proc Natl Acad Sci USA* 1988, **85**: 252–256.
74. Onji, M., Lever, A.M.L., Saito, I. & Thomas, H.C. Defective response to interferons in cells transfected with the hepatitis B virus genome. *Hepatology* 1989, **9**: 92–96.
75. De Martino, M., Rossi, M.E.R., Muccioli, A.T., Resti, M. & Vierucci, A. Interference of hepatitis B virus surface antigen with natural kill cell function. *Clin Exp Immunol* 1985, **61**: 90–95.
76. Zhao, W., Egan, M. & Peters, M. B cell responses in patients with acute viral hepatitis B. In: *Proceedings of the AASLD 1990 Immunology and the Liver meeting*. pp. 29.
77. Anastassakos, Ch., Alexander, G.J.M., Wolstencroft, R.A. *et al.* Interleukin 1 and interleukin 2 activity in chronic hepatitis B virus infection. *Gastroenterology* 1988, **94**: 999–1005.
78. Sheron, N., Lau, J., Daniels, H.M., Goka, J., Eddleston, A.L.W.F., Alexander, G.J.M. & Williams, R. Increased production of tumour necrosis factor-alpha in chronic hepatitis B virus infection. *J Hepatol* 1991, **12**: 241–245.
79. Fattovich, G., Brollo, L., Alberti, A. *et al.* Chronic persistent hepatitis type B can be a progressive disease when associated with sustained virus replication. *J Hepatol* 1990, **11**: 29–33.
80. Bain, V.G., Alexander, G.J.M. & Eddleston, A.L.W.F. Immunopathogenesis of viral hepatitis. In: Targan, S.R. & Shanahan, F. (eds) *Immunology and Immunopathology of the Liver and Gastroenterology Tract*. Igaku-Shoin Press, New York, 1989, pp. 255–280.
81. Alexander, G.J.M., Brahm, J., Fagan, E.A. *et al.* Loss of HBsAg with interferon therapy in chronic hepatitis B virus infection. *Lancet* 1987, **ii**: 66–69.
82. Lok, A.S.F., Novick, D.M., Karayiannis, P. *et al.* A randomised study of the effects of adenine arabinoside 5'-monophosphate (short or long courses) and lymphoblastoid interferon on hepatitis B virus replication. *Hepatology* 1985, **5**: 1132–1138.
83. Dusheiko, G., Dibisceglie, A., Bowyer, S. *et al.* Recombinant leukocyte interferon treatment of chronic hepatitis B. *Hepatology* 1985, **5**: 556–560.
84. Perrillo, R.P., Schiff, E., Davis, G.L. *et al.* A randomised controlled trial of interferon alfa-2b alone and after prednisone withdrawal for the treatment of chronic hepatitis B. *N Engl J Med* 1990, **323**: 295–301.
85. Hoofnagle, J.H., Peters, M., Mullen, K.D. *et al.* Randomized controlled trial of recombinant human α interferon in patients with chronic hepatitis B. *Gastroenterology* 1988, **95**: 1318–1325.
86. Dooley, J.S., Vergalla, J., Hoofnagle, J.H., Zoon, K.C., Munson, P.J. & Jones, E.A. Specific binding of human alpha interferon to high affinity cell surface binding sites on peripheral blood mononuclear cells. *J Lab Clin Med* 1989, **113**: 623–631.
87. Lau, J.Y.N., Sheron, N., Morris, A.G., Bomford, A.B., Alexander, G.J.M. & Williams, R. Interferon-alpha receptor expression and regulation in chronic hepatitis B virus infection. *Hepatology* 1990, **13**: 332–338.
88. Bissett, J., Eisenberg, M., Gregghory, P., Robinson, W.S. & Merigan, T.C. Recombinant fibroblast interferon and immune interferon for treating chronic hepatitis B virus infection: patients' tolerance and the effect on viral markers. *J Infect Dis* 1988, **157**: 1076–1080.
89. Gomez, C., LaBanda, F., Porres, J.C. *et al.* Combined recombinant alpha and gamma interferon treatment of chronic hepatitis B virus infection. In: Zuckerman, A.J. (ed) *Viral Hepatitis and Liver Disease*. Alan R Liss, New York, 1988, pp. 872–874.
90. Di-Bisceglie, A.M., Rustgi, V.K., Kassianides, C. *et al.* Therapy of chronic hepatitis B with recombinant human alpha and gamma interferon. *Hepatology* 1990, **11**: 266–270.
91. Thagarajan, S.P., Subramanian, S., Thirunalasundari, T., Venkateswaran, P.S. & Blumberg, B.S. Effect of *Phyllanthus amarus* on chronic carriers of hepatitis B virus. *Lancet* 1988, **ii**: 764–766.
92. Kassianides, C., Hoofnagle, J.H., Miller, R.H. *et al.* Inhibition of duck hepatitis B virus replication by 2',3'-dideoxycytidine. *Gastroenterology* 1989, **97**: 1275–1280.
93. Zuckerman, A.J. Immunisation against hepatitis B. In: Zuckerman, A.J. (ed) *British Medical Bulletin: Viral Hepatitis*. Churchill Livingstone, London, 1990, pp. 383–398.
94. Stevens, C.E., Taylor, P.E., Tong, M.J. *et al.* Hepatitis B vaccine: An overview. In: Vyas, G.N., Dienstag, J.L., Hoofnagle, J.H. (eds) *Viral Hepatitis and Liver Disease*. Grune & Stratton, Orlando, Florida, 1984, p. 275.
95. Hadler, S.C., Francis, D.P., Maynard, J.E. *et al.* Long term immunogenicity and efficacy of hepatitis B vaccine in homosexual men. *N Engl J Med* 1986, **315**: 209.
96. Carman, W.F., Zanetti, A.R., Karayiannis, P. *et al.* Vaccine-induced escape mutant of hepatitis B virus. *Lancet* 1990, **336**: 325–329.
97. Itoh, Y., Takai, E., Ohnuma, H. *et al.* A synthetic peptide vaccine involving the product of the pre-S2 region of hepatitis B virus DNA: protective efficacy in chimpanzees. *Proc Natl Acad Sci USA* 1986, **83**: 9174–9178.

98. Meuer, S.C., Dumann, H., Meyer zum Brischenfelde, K.H. & Kohnler, H. Low dose interleukin-2 induces systemic immune responses against HBsAg in immunodeficient non-responders to hepatitis B vaccination. *Lancet* 1989, **i**: 15.

Hepatitis C Virus

99. Feinstone, S.M., Kapikian, A.Z., Purcell, R.H., Alter, H.J. & Holland, P.V. Transfusion-associated hepatitis not due to viral hepatitis type A or B. *N Engl J Med* 1975, **292**: 767–770.
100. Knodell, R.G., Conrad, M.E., Dienstag, J.L. & Bell, C.J. Etiological spectrum of post-transfusion hepatitis. *Gastroenterology* 1975, **69**: 1278–1285.
101. Dienstag, J.L., Non-A, non-B hepatitis: recognition, epidemiology and clinical features. *Gastroenterology* 1983, **83**: 439–462.
102. Shih, J.W.K., Esteban, J.I. & Alter, H.J. Non-A, non-B hepatitis: advances and unfulfilled expectations of the first decade. *Prog Liver Dis* 1986, **8**: 433–452.
103. Dienstag, J.L. & Alter, H.J. Non-A, non-B hepatitis: evolving epidemiological and clinical perspectives. *Semin Liver Dis* 1986, **6**: 67–81.
104. Overby, L.R. Serology of liver diseases. *Curr Hepatol* 1987, **7**: 35–68.
105. Hallam, N.F. Non-A, non-B hepatitis: reverse transcriptase activity. *Lancet* 1985, **ii**: 665.
106. Alter, H.J., Purcell, R.H., Holland, P.V. & Popper, H., Transmissible agent in non-A, non-B hepatitis. *Lancet* 1978, **i**: 459–463.
107. Shimizu, Y.K., Feinstone, S.M., Purcell, R.H., Alter, H.J. & London, W.T. Non-A, non-B hepatitis: ultrastructural evidence for two antigens in experimentally infected chimpanzees. *Science* 1979, **205**: 197–200.
108. Bradley, D.W., Maynard, J.E., Popper, H. *et al.* Post-transfusion non-A, non-B hepatitis: Physicochemical properties of two distinct agents. *J Infect Dis* 1983, **148**: 254–265.
109. Bradley, D.W., McCaustland, K.A., Cook, E.H. *et al.* Post-transfusion non-A, non-B hepatitis in chimpanzees: physicochemical evidence that the tubule-forming agent is a small enveloped virus. *Gastroenterology* 1985, **88**: 773–779.
110. He, L.F., Alling, D., Popkin, T., Shapiro, M., Alter, H.J. & Purcell, R.H. Determining the size of non-A, non-B hepatitis virus by filtration. *J Infect Dis* 1987, **156**: 636–640.
111. Hollinger, F.B., Mosley, J.W., Szmunes, W., Aach, R.D., Peters, R.I. & Stevens, C. Transfusion-transmitted viruses study: experimental evidence for two non-A, non-B hepatitis agents. *J Infect Dis* 1980, **142**: 400–407.
112. Bradley, D.W., Maynard, J.E., Cook, E.H. *et al.* Non-A, non-B hepatitis in experimental infected chimpanzees: cross challenge and electron microscopic studies. *J Med Virol* 1980, **6**: 185–201.
113. Choo, Q.L., Kuo, G., Weiner, A.J., Overby, L.R., Bradley, D.W. & Houghton, M. Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science* 1989, **244**: 359.
114. Arima, T., Nagashima, H., Murakami, S., Kaji, C., Fujita, J., Shimomura, H. & Tsuji, T. Cloning of a cDNA associated with acute and chronic hepatitis C infection generated from patients serum RNA. *Gastroenterol-Jpn* 1989, **24**: 540–544.
115. Kuo, G., Choo, Q.L., Alter, H.J. *et al.* An assay for circulating antibodies to a major etiologic virus of non-A, non-B hepatitis. *Science* 1989, **244**: 362.
116. Editorial. Will the real hepatitis C stand up. *Lancet* 1989, **ii**: 307–308.
117. Weiner, A.J., Kuo, G., Bradley, D.W. *et al.* Detection of hepatitis C viral sequences in non-A, non-B hepatitis. *Lancet* 1990, **335**: 1–5.
118. Choo, Q.L., Weiner, A.J., Overby, L.R., Kuo, G., Houghton, M. & Bradley, D.W. Hepatitis C virus: the major causative agent of viral non-A, non-B hepatitis. In: Zuckerman, A.J. (ed) *British Medical Bulletin: Viral Hepatitis*. Churchill Livingstone, London, 1990, pp. 423–441.

119. Alter, H.J., Purcell, R.H., Shih, J.W. *et al.* Detection of antibody to hepatitis C virus in prospectively followed transfusion recipients with acute and chronic non-A, non-B hepatitis. *N Engl J Med* 1989, **321**: 1494–1500.
120. Colombo, M., Kuo, G., Choo, Q.L. *et al.* Prevalence of antibodies to hepatitis C virus in Italian patients with hepatocellular carcinoma. *Lancet* 1989, **ii**: 1006–1009.
121. Sirchia, G., Bellobuono, A., Giovanetti, A. & Marconi, M. Antibodies to hepatitis C virus in Italian blood donors. *Lancet* 1989, **ii**: 797.
122. Esteban, J.I., Esterban, R., Viladomiu, L. *et al.* Hepatitis C virus antibodies among high risks groups in Spain. *Lancet* 1989, **ii**: 294–296.
123. Bruiz, J., Barrera, J.M., Calvet, X. *et al.* Prevalence of antibodies to hepatitis C virus in Spanish patients with hepatocellular carcinoma and hepatic cirrhosis. *Lancet* 1989, **ii**: 1004–1006.
124. Lenzi, M., Ballardini, G., Fusconi, M., *et al.* Type 2 autoimmune hepatitis and hepatitis C virus infection. *Lancet* 1990, **335**: 258–259.
125. Jeffers, L., deMedina, M., Hasan, F. *et al.* Hepatitis C associated idiopathic chronic hepatitis and cryptogenic cirrhosis. *Hepatology* 1989, **9** (suppl): abstract 301.
126. Katkov, W.N., Cody, H., Evans, A.A. *et al.* The role of hepatitis C virus in chronic liver disease. *Hepatology* 1989, **9** Suppl: abstract 302.
127. Evans, A.A., Cody, H., Kuo, G. *et al.* Seroepidemiology of hepatitis C virus in selected populations. *Hepatology* 1989, (suppl): abstract 304.
128. McFarlane, I.G., Smith, H.M., Johnson, P.J., Bray, G., Vergani, D. & Williams, R. Significance of possible anti-HCV antibodies in autoimmune chronic active hepatitis. *Lancet* 1990, **335**: 754–757.
129. Davis, G.L., Balart, L.A., Schiff, E.R. *et al.* Treatment of chronic hepatitis C with recombinant interferon alfa – multicentre randomised controlled trial. *N Engl J Med* 1991, **321**: 1501–1506.
130. Di Bisceglie, A.M., Martin, P., Kassianides, C. *et al.* Recombinant interferon alfa therapy for chronic hepatitis C – a randomised, double-blinded, placebo-controlled trial. *N Engl J Med* 1989, **321**: 1506–1510.
131. Hoofnagle, J.H. & Di Bisceglie, A.M., Therapy of chronic hepatitis: chronic hepatitis D and non-A, non-B hepatitis. In: Zuckerman, A.J. (ed) *Viral Hepatitis and Liver Disease*. Alan R Liss, New York, 1988, pp. 823–830.

Hepatitis D Virus

132. Rizzetto, M., Canese, M.G., Arico, S. *et al.* Immunofluorescence detection of a new antigen-antibody system (delta-antidelta) associated with the hepatitis B virus in the liver and in the serum of HBsAg carriers. *Gut* 1977, **18**: 997–1003.
133. Monjardino, J.P., Saldanha, J.A., Delta hepatitis – the disease and the virus. In: Zuckerman, A.J. (ed) *British Medical Bulletin: Viral Hepatitis*. Churchill Livingstone, London, 1990, pp. 399–407.
134. Rizzetto, M., Gerlin, J.L. & Purcell, R.H. The hepatitis delta virus and its infection. *Prog Clin Biol Res* 1987, **234**: 361–534.
135. Hanson, B.G., Moestrup, T., Widell, A. *et al.* Infection with delta agent in Sweden: introduction of a new hepatitis agent. *J Infect Dis* 1982, **146**: 472–478.
136. Hoofnagle, J.H. Type D (delta) hepatitis. *JAMA* 1989, **261**: 1321.
137. Rizzetto, M. & Verme, G. Delta hepatitis. *J Hepatol* 1985, **1**: 187–193.
138. Rizzetto, M., Hoyer, B., Canese, M.G. *et al.* Delta agent: association delta antigen with hepatitis B surface antigen and RNA in serum of delta-infected chimpanzees. *Proc Natl Acad Sci USA* 1980, **77**: 6124–6128.
139. Wand, K.S., Choo, Q.L., Weiner, A.J. *et al.* Structure, sequence and expression of the hepatitis delta (δ) viral genome. *Nature* 1986, **323**: 508–514.

140. Kos, A., Dijkema, R., Arnberg, A.C. *et al.* The hepatitis delta virus possesses a circular RNA. *Nature* 1986, **323**: 558–560.
 141. Chen, P.J., Kalpana, G., Goldberg, J. *et al.* Structure and replication of the genome of the hepatitis delta virus. *Proc Natl Acad Sci USA* 1986, **83**: 8774–8778.
 142. Sharmeen, L., Kuo, M.Y.P., Dinter-Gottlieb, G. & Taylor, J. Antigenomic RNA of human hepatitis delta virus can undergo self-cleavage. *J Virol* 1988, **62**: 2674–2679.
 143. Wu, H.N., Lin, Y.J., Lin, F.P. *et al.* Human hepatitis delta virus RNA subfragments contain an autocleavage activity. *Proc Natl Acad Sci USA* 1989, **86**: 1831–1835.
 144. Sharmeen, L., Kuo, M.Y.P. & Taylor, J. Self-ligating RNA sequences on the antigenome of human hepatitis delta virus. *J Virol* 1989, **63**: 1428–1430.
 145. Wu, H.N. & Lai, M.M.C. Reversible cleavage and ligation of hepatitis delta RNA. *Science* 1989, **243**: 652–654.
 146. Taylor, J., Mason, W., Summers, J. *et al.* Replication of human hepatitis delta virus in primary cultures of woodchuck hepatocytes. *J Virol* 1987, **61**: 2891–2895.
 147. Kuo, M.Y.P., Chao, M. & Taylor, J. Initiation of replication of the human hepatitis delta virus genome from cloned DNA: role of delta antigen. *J Virol* 1989, **63**: 1945–1950.
 148. Negro, F., Baldi, M., Bonino, F. *et al.* Chronic HDV (hepatitis delta virus) hepatitis: intrahepatic expression of delta antigen, histologic activity and outcome of liver disease. *J Hepatol* 1988, **6**: 8–14.
 149. Chu, C.M. & Liaw, Y.F. Studies on the composition of the mononuclear cell infiltrates in liver from patients with chronic active delta hepatitis. *Hepatology* 1989, **10**: 911–915.
 150. Lenzi, M., Fusconi, M., Cassani, F. *et al.* Characterisation of LKM1 and LKM2 antigens. *Prog Clin Biol Res* 1987, **234**: 343–345.
 151. Krogsgaard, K., Aldershvile, J., Kryger, P. *et al.* Hepatitis B virus DNA, HBeAg and delta infection during the course from acute to chronic hepatitis B virus infection. *Hepatology* 1985, **5**: 778–782.
 152. Krogsgaard, K., Kryger, P., Aldershvile, J. *et al.* Delta-infection and suppression of hepatitis B virus replication in chronic HBsAg carriers. *Hepatology* 1987, **7**: 42–45.
 153. Rizzetto, M., Ponzetto, A., Bonino, F. & Smedile, A. Hepatitis delta virus infection: clinical and epidemiological aspects. In: Zuckerman, A.J. (ed.) *Viral Hepatitis and Liver Disease*. Alan R. Liss, New York, 1988, pp. 389–394.
 154. Smedile, A., Lavarini, C., Crivelli, O., Raimondo, G., Fassone, M., Rizzetto, M. Radioimmunoassay detection of IgM antibodies to HBV associated delta antigen: clinical significance in delta infection. *J Med Virol* 1982, **9**: 131–138.
 155. Smedile, A., Rizzetto, M., Denniston, K. *et al.* Type D hepatitis: the clinical significance of hepatitis D virus RNA in serum as detected by a hybridisation-based assay. *Hepatology* 1986, **6**: 1297–1302.
 156. Govindarajan, S., Gupta, S., Valinluck, B. & Redeker, A.G. Correlation of IgM anti-hepatitis D virus (HDV) to HDV RNA in sera of chronic HDV. *Hepatology* 1989, **10**: 34–35.
 157. Negro, F., Bonino, F., Di Bisceglie, A., Hoofnagle, J.H. & Gerin, J.L. Intrahepatic markers of hepatitis delta virus infection: a study by *in situ* hybridisation. *Hepatology* 1989, **10**: 916–920.
 158. Smedile, A., Bergmann, F., Baroudy, B.M. *et al.* Riboprobe assay for HDV RNA: a sensitive method for the detection of the HDV genome in clinical serum samples. *J Med Virol* 1990, **30**: 20–24.
 159. Hoofnagle, J.H., Mullen, K., Peters, M. *et al.* Treatment of chronic delta hepatitis with recombinant alpha interferon. In: Rizzetto, M., Gerin, J.L., Purcell, R.H. (eds) *The Hepatitis Delta Virus and its Infection*. Alan R. Liss, New York, 1987, pp. 291–298.
 160. Rizzetto, M., Rosina, F., Saracco, G. *et al.* Treatment of chronic delta hepatitis with alpha-2 recombinant interferon. *J Hepatol* 1986, **3**: S229–S233.
 161. Thomas, H.C., Farci, P., Shein, R. *et al.* Inhibition of hepatitis delta virus (HDV) replication with lymphoblastoid human alpha interferon. In: Rizzetto, M., Gerin, J.L., Purcell, R.H. (eds) *The Hepatitis Delta Virus and Its Infections*. Alan R. Liss, New York, 1987, pp. 277–290.
 162. Hedin, G., Weiland, O., Ljunggren, K. *et al.* Treatment of fulminant hepatitis B and fulminant hepatitis B and D coinfection with foscarnet. In: Rizzetto, M., Gerin, J.L., Purcell, R.H. (eds) *The Hepatitis Delta Virus and Its Infections*. Alan R. Liss, New York, 1987, pp. 309–320.
- ### Hepatitis E Virus
163. Viewanath, R. Epidemiology: infectious hepatitis in Delhi 1955–6. *Indian J Med Res* 1957, **45** (Suppl): 1–29.
 164. Naidu, S.S. & Viewanath, R. Infectious hepatitis in Delhi (1955–6): a critical study; observations in pregnant women. *Indian J Med Res* 1957, **45** (Suppl 1): 71–76.
 165. Gupta, D.N. & Smetana, H.F. The histopathology of viral hepatitis is seen in the Delhi epidemic. *Indian J Med Res* 1957, **45** (Suppl 1): 101–103.
 166. Wong, D.C., Purcell, R.H., Sreenivasan, M.A. *et al.* Epidemic and endemic hepatitis in India: evidence for a non-A, non-B hepatitis virus aetiology. *Lancet* 1980, **ii**: 876–879.
 167. Ramalingaswami, V. & Purcell, R.H. Waterborne non-A, non-B hepatitis. *Lancet* 1988, **i**: 571–573.
 168. Bradley, D.W. Enterically-transmitted non-A, non-B hepatitis. In: Zuckerman, A.J. (ed.) *British Medical Bulletin: Viral Hepatitis*. Churchill Livingstone, London, 1990, pp. 442–461.
 169. De Cock, K.M., Bradley, D.W., Sanford, N.L., Govindarajan, S., Maynard, J.E. & Refeker, A.G. Epidemic non-A, non-B hepatitis in patients from Pakistan. *Ann Intern Med* 1987, **106**: 227–230.
 170. Balayan, M.S., Andzhaparidze, A.G., Savinskaya, S.S. *et al.* Evidence for a virus in non-A, non-B hepatitis transmitted via the fecal oral route. *Intervirology* 1983, **20**: 23.
 171. Arankalle, V.A., Ticehurst, J., Sreenivasan, M.A. *et al.* Aetiological association of a virus-like particle with enterically transmitted non-A, non-B hepatitis. *Lancet* 1988, **i**: 550–554.
 172. Abe, K., Shikata, T., Fujisawa, K. *et al.* Transmission of non-A, non-B hepatitis agent to chimpanzees from patients of epidemic hepatitis. *Jpn J Med Sci Biol* 1986, **39**: 119–127.
 173. Bradley, D.W., Krawczynski, K., Cook, E.H. *et al.* Enterically transmitted non-A, non-B hepatitis: serial passage of disease in cynomolgus macaques and tamarins and recovery of disease-associated 27 to 34 nm virus like particles. *Proc Natl Acad Sci USA* 1987, **84**: 6277–6281.
 174. Krawczynski, K. & Bradley, D.W. Enterally transmitted non-A, non-B hepatitis: identification of virus associated antigen in experimentally infected cynomolgus macaques. *J Infect Dis* 1989, **159**: 1042.
 175. Kane, M.A., Bradley, D.W., Shrestha, S.M. *et al.* Epidemic non-A, non-B hepatitis in Nepal. Recovery of a possible etiologic agent and transmission studies in possible etiologic agent and transmission studies in marmosets. *JAMA* 1984, **252**: 3140–3145.
 176. Panda, S.K., Datta, R., Kaur, J., Zuckerman, A.J. & Nayak, N.C. Enterically transmitted non-A, non-B hepatitis: recovery of virus like particles from an epidemic in South Delhi and transmission studies in rhesus monkeys. *Hepatology* 1989, **10**: 466–472.
 177. Reyes, G.R., Purdy, M.A., Kim, J.P. *et al.* Isolation of a cDNA from the virus responsible for enterically transmitted non-A, non-B hepatitis. *Science* 1990, **247**: 1335–1339.
 178. Zuckerman, A.J. Hepatitis E virus. *Br Med J* 1990, **300**: 1475–1476.
 179. Khuroo, M.S., Teli, M.R., Skidmore, S. *et al.* Incidence and severity of viral hepatitis in pregnancy. *Am J Med* 1981, **70**: 252–255.

180. Gupta, D.N. & Smetana, H.F. The histopathology of viral hepatitis as seen in the Delhi epidemic. *Indian J Med Res* 1957, **45**: 101–113.

Fulminant hepatic failure

181. Trey, C. & Davidson, C.S. *The Management of Fulminant Hepatic Failure*. Grune & Stratton, New York, 1970, 282–298.
182. Minton, N.A., Henry, J.A. & Frankel, R.J. Fatal paracetamol poisoning in an epileptic. *Hum Toxicol* 1988, **33–4**: 33–34.
183. Lauterburg, B.H. & Velez, M.E. Glutathione deficiency in alcoholics: risk factor for paracetamol hepatotoxicity. *Gut* 1988, **29**: 1153–1157.
184. Prescott, L.F., Illingworth, R.N., Critchley, J.A.J.H., Stewart, M.J., Adam, R.D. & Proudfoot, A.T. Intravenous N-acetylcysteine: the treatment of choice for paracetamol poisoning. *Br Med J* 1979, **2**: 1097–1100.
185. Smilkstein, M.J., Knapp, G.L., Kulig, K.W. & Rumack, B.H. Efficacy of oral N-acetylcysteine in the treatment of acetaminophen overdose. *N Engl J Med* 1988, **319**: 1557–1562.
186. Harrison, P., Keays, R., Bray, G., Alexander, G. & Williams, R. Improved outcome in paracetamol-induced fulminant hepatic failure following late administration of N-acetylcysteine. *Lancet* 1990, **335**: 1572–1573.
187. Keays, R., Forbes, A., Davies, S. *et al.* N-acetylcysteine improves outcome in paracetamol-induced fulminant hepatic failure. *Gut* 1989, **30**: A1512.
- 187a. Harrison, P.M., Wendon, J.A., Grimson, A.E.S., Alexander, G.J.M. & Williams, R. Improvement by acetylcysteine of systemic hemodynamics and oxygen transport in fulminant hepatic failure. *N Engl J Med* 1991, in press.
188. Shoemaker, W., Appel, P., Kram, H., Waxman, K. & Lee, T.-S. Prospective trial of supranormal values as therapeutic goals in high-risk surgical patients. *Chest* 1988, **94**: 1176–1186.
189. Gammal, S.H. & Jones, E.A. Hepatic encephalopathy. *Med Clin North Am* 1989, **73**: 793–813.
190. Basile, A.S., Pannell, L., Jaouni, T. *et al.* Brain concentrations of benzodiazepines are elevated in an animal model of hepatic encephalopathy. *Proc Natl Acad Sci USA* 1990, **87**: 5263–5267.
191. Ganz, R., Swain, M., Traber, P., DalCanto, M., Butterworth, R.F. & Blei, A.T. Ammonia-induced swelling of rat cerebral cortical slices: implications for the pathogenesis of brain edema in acute hepatic failure. *Metab Brain Dis* 1989, **4**: 213–223.
192. Almdal, T., Schroeder, T. & Ranek, L. Cerebral blood flow and liver function in patients with encephalopathy due to acute and chronic liver diseases. *Scand J Gastroenterol* 1989, **24**: 299–303.
193. Davenport, A., Will, E.J. & Davison, A.M. Effect of posture on intracranial pressure and cerebral perfusion pressure in patients with fulminant hepatic and renal failure after acetaminophen self-poisoning. *Crit Care Med* 1990, **18**: 286–289.
194. Sari, A., Yamashita, S., Ohosita, S. *et al.* Cerebrovascular reactivity to CO₂ in patients with hepatic or septic encephalopathy. *Resuscitation* 1990, **19**: 125–134.
195. Canalese, J., Gimson, A.E.S., Davis, C. *et al.* Controlled trial of dexamethasone and mannitol for the cerebral oedema of fulminant hepatic failure. *Gut* 1982, **23**: 625–629.
196. Forbes, A., Alexander, G.J., O'Grady, J.G. *et al.* Thiopental infusion in the treatment of intracranial hypertension complicating fulminant hepatic failure. *Hepatology* 1989, **10**: 306–310.
197. Davenport, A., Will, E.J., Davison, A.M. *et al.* Changes in intracranial pressure during machine and continuous haemofiltration. *Int J Artif Organs* 1989, **12**: 439–444.
198. Davenport, A., Will, E.J., Davison, A.M. *et al.* Changes in intracranial pressure during haemofiltration in oliguric patients with grade IV hepatic encephalopathy. *Nephron* 1989, **53**: 142–146.
199. Rolando, N., Harvey, F., Brahm, J. *et al.* Prospective study of bacterial infection in acute liver failure: an analysis of fifty patients. *Hepatology* 1990, **11**: 49–53.
200. Rolando, N., Harvey, F., Brahm, J. *et al.* Fungal infection: a common, unrecognised complication of acute liver failure. *J Hepatol* 1991, **12**: 1–9.

Liver regeneration

201. Rakela, J., Lange, S.M., Ludwig, J. & Baldus, W. Fulminant hepatitis: Mayo clinic experience with 34 cases. *Mayo Clin Proc* 1985, **60**: 289–292.
202. McGowan, J.A., Strain, A.J. & Bucher, N.L.R. DNA synthesis in primary cultures of adult rat hepatocytes in a defined medium: effects of epidermal growth factor, insulin, glucagon and cyclic-AMP. *J Cell Physiol* 1981, **180**: 353–363.
203. Mead, J.E. & Fausto, N. Transforming growth factor TGF α may be a physiological regulator of liver regeneration by means of an autocrine mechanism. *Proc Natl Acad Sci USA* 1989, **86**: 1558–1562.
204. Tsubouchi, H., Hirono, S., Gohda, E. *et al.* Clinical significance of human hepatocyte growth factor in blood from patients with fulminant hepatic failure. *Hepatology* 1989, **9**: 875–881.
205. Harrison, P.M., Hughes, R.D., Forbes, A., Potmann, B., Alexander, G.J. & Williams, R. Failure of insulin and glucagon infusion to stimulate liver regeneration in fulminant hepatic failure. *J Hepatol* 1990, **10**: 332–336.
206. Nakamura, T., Arakaki, R. & Ichihara, A. Interleukin-1 is a potent growth inhibitor of adult rat hepatocytes in primary culture. *Exp Cell Res* 1988, **179**: 488–497.

Liver transplantation

207. Emond, J.C., Aran, P.P., Whittington, P.F., Broelsch, C.E. & Baker, A.L. Liver transplantation in the management of fulminant hepatic failure. *Gastroenterol* 1989, **96**: 1583–1588.
208. Iwatsuki, S., Stieber, A.C., Marsh, J.W. *et al.* Liver transplantation for fulminant hepatic failure. *Transplant Proc* 1989, **21**: 2431–2434.
209. Klein, A.S., Hart, J., Brems, J.J., Goldstein, L., Lewin, K. & Busuttil, R.W. Amanita poisoning: treatment and the role of liver transplantation. *Am J Med* 1989, **86**: 187–193.
210. O'Grady, J.G., Alexander, G.J., Thick, M., Potter, D., Calne, R.Y. & Williams, R. Outcome of orthotopic liver transplantation in the aetiological and clinical variants of acute liver failure. *Q J Med* 1988, **68**: 817–824.
211. Rakela, J., Perkins, J.D., Gross, J.J. *et al.* Acute hepatic failure: the emerging role of orthotopic liver transplantation. *Mayo Clin Proc* 1989, **64**: 424–428.
212. Schafer, D.F. & Shaw, B.J. Fulminant hepatic failure and orthotopic liver transplantation. *Semin Liver Dis* 1989, **9**: 189–194.
213. Van S.J., Gelin, M., Bourgeois, N., De, P.C., Adler, M. & Lambilliotte, J.P. Liver transplantation for fulminant and subacute viral hepatic failure in adults. *Transplant Proc* 1990, **22**: 1505–1508.
214. O'Grady, J.G., Alexander, G.J., Hayllar, K.M. & Williams, R. Early indicators of prognosis in fulminant hepatic failure. *Gastroenterology* 1989, **97**: 439–445.
215. Harrison, P.M., O'Grady, J.G., Keays, R., Alexander, G.J.M. & Williams, R. Serial prothrombin time as prognostic indicator in paracetamol-induced fulminant hepatic failure. *Br Med J* 1990, **301**: 964–966.

216. Moritz, M.J., Jarrell, B.E., Armenti, V. *et al.* Heterotopic liver transplantation for fulminant hepatic failure—a bridge to recovery. *Transplantation* 1990, **50**: 524–526.
217. Sundberg, R., Alumets, J., Ar'Rajab, A., Ahren, B. & Bengmark, S. Ultrastructure in liver preservation with Eurocollins and UW solution. *Transplant Proc* 1990, **22**: 176–179.
218. Iwatsuki, S., Starzl, T.E., Todo, S. *et al.* Experience in 1,000 liver transplants under cyclosporine-steroid therapy: a survival report. *Transplant Proc* 1988, **20**: 498–504.
219. O'Grady, J.G., Smith, H.M., Davis, S.E. *et al.* Hepatitis B virus reinfection after orthotopic liver transplantation: serological and clinical implications. *J Hepatol* (in press).
220. Pontisso, P., Poon, M.C., Tiollais, P. & Brechot, C. Detection of hepatitis B virus DNA in mononuclear blood cells. *Br Med J* 1984, **288**: 1563–1566.
221. Davison, F., Alexander, G.J.M., Trowbridge, R., Fagan, E.A. & Williams, R. Detection of hepatitis B virus DNA in spermatozoa, urine, saliva and leukocytes, of chronic HBsAg carriers: a lack of relationship with serum markers of replication. *J Hepatol* 1987, **4**: 37–44.
222. Davies, S.E., Portmann, B., O'Grady, J.G. *et al.* Hepatic histology following transplantation for chronic hepatitis B virus infection including a unique pattern of fibrosing cholestatic hepatitis. *Hepatology* 1991, **13**: 150–157.
223. Tur-Kaspa, R., Burk, R.D., Shaul, Y. & Shafritz, D.A. Hepatitis B virus DNA contains a glucocorticoid responsive element. *Proc Natl Acad Sci USA* 1986, **83**: 1627–1631.
224. Tur-Kaspa, R. & Laub, O. Corticosteroids stimulate hepatitis B virus DNA, mRNA and protein production in a stable expression system. *J Hepatol* 1990, **11**: 34–36.
225. Lau, J.Y.N., Lai, C.L., Lin, H.J. *et al.* Fatal reactivation of chronic hepatitis B virus infection following chemotherapy withdrawal in lymphoma patients. *Q J Med* 1989, **73**: 911–917.
226. Bird, G.L.A., Smith, H.M., Portman, B., Alexander, G.J.M. & Williams, R. Acute liver decompensation on withdrawal of cytotoxic chemotherapy and immunosuppressive therapy in hepatitis B carriers. *Q J Med* 1989, **73**: 895–902.
227. Reynes, M., Zigneg, L., Samuel, D. *et al.* Graft hepatitis delta virus reinfection after orthotopic liver transplantation in HDV cirrhosis. *Transplant Proc* 1989, **21**: 2424–2425.
228. Lauchart, W., Muller, R. & Pichlmayr, R. Immunoprophylaxis of hepatitis B virus reinfection in recipients of human allografts. *Transplant Proc* 1987, **19**: 2387–2389.
229. O'Grady, J.G., Polson, R.J., Rolles, K., Calne, R.Y. & Williams, R. Liver transplantation in malignant disease: results in 93 consecutive patients. *Ann Surg* 1988, **207**: 373–379.
230. Arnold, J.C. O'Grady, J.G., Bird, G., Calne, R.Y. & Williams, R. Liver transplantation for primary and secondary hepatic APUDomas. *Br J Surg* 1989, **76**: 248–249.
231. Calne, R.Y., Rolles, K., White, D.J.G. *et al.* Cyclosporin A initially as the only immunosuppressant in 34 recipients of cadaveric organs: 32 kidneys, 2 pancreas, 2 liver. *Lancet* 1979, **ii**: 1033–1036.
232. Starzl, T.E., Todo, S., Fung, J., Demetris, A.J. Venkataraman, R. & Jain, A. FK506 for liver, kidney, and pancreas transplantation. *Lancet* 1989, **ii**: 1000–1004.
233. Kino, T., Hatanaka, H., Miyata, S. *et al.* FK506, a novel immunosuppressant isolated from streptomyces. Immunosuppressive effects of FK506 *in vitro*. *J Antibiot (Tokyo)* 1987, **40**: 1256–1265.
234. Ochial, T., Nakajima, K., Nagata, M. *et al.* Effect of a new immunosuppressive agent, FK506, on heterotopic allotransplantation in the rat. *Transplant Proc* 1987, **19**: 1284–1286.
235. Starzl, T.E., Makowka, L., Todo, S. *et al.* FK506: a potential breakthrough in immunosuppression. *Transplant Proc* 1987, **19** (Suppl 6): 103–104.
236. O'Grady, J.G., Alexander, G.J.M., Sutherland, S. *et al.* Cytomegalovirus infection and donor/recipient HLA antigens: interdependent co-factors in pathogenesis of vanishing bile duct syndrome after liver transplantation. *Lancet* 1988, **ii**: 302–305.

Portal hypertension

237. Schrier, R.W., Arroyo, V., Bernardi, M. *et al.* Peripheral arterial vasodilation hypothesis: A proposal for the initiation of renal sodium and water retention in cirrhosis. *Hepatology* 1988, **8**: 1151–1157.
238. Palmer, R.M., Ashton, D.S. & Moncada, S. Vascular endothelial cells synthesize nitric oxide from L-arginine. *Nature* 1988, **333**: 664–666.
239. Vallance, P.J.T. & Moncada, S. Hyperdynamic circulation in cirrhosis: a role for nitric oxide. *Lancet* 1991, **337**: 776–778.
240. Pagliaro, L., Burroughs, A.K., Sorensen, T.I.A. *et al.* Therapeutic controversies and randomised controlled trials (RCTs): prevention of bleeding and rebleeding in cirrhosis. *Gastroenterol Int* 1989, **2**: 71–84.
241. MacMathuna, P., Westaby, D. & Williams, R. Taking the tension out of the portal system: an approach to the management of portal hypertension in the 1990's. *Scand J Gastroenterol* 1990, **25** (Suppl 175): 131–145.
242. Stiegmann, G.V., Goff, J.S., Sun, J.H. & Hruza, D. Endoscopic ligation of oesophageal varices. *Am J Surg* 1990, **159**: 21–26.
243. Gimson, A.E.S., Westaby, D. & Williams, R. Endoscopic sclerotherapy and management of gastric variceal haemorrhage. *Gut* 1989, **30**: A1497.
244. Vlavianos, P., Gimson, A.E., Westaby, D. & Williams, R. Balloon tamponade in variceal bleeding: use and misuse. *Br Med J* 1989, **298**: 1158.
245. Burroughs, A.K., Hamilton, G., Phillips, A., Mezzanotte, G., McIntyre, N. & Hobbs, K.E. A comparison of sclerotherapy with staple transection of the esophagus for the emergency control of bleeding from esophageal varices. *N Engl J Med* 1989, **321**: 857–862.