Reviews in Medicine

General surgery

Irving Taylor

University Surgical Unit, F Level Centre Block, Southampton General Hospital, Tremona Road, Southampton SO9 4XY, UK

This review of general surgical 'highlights' will once again concentrate on the following topics; breast, gastro-oesophageal, colorectal, hepato-pancreaticobiliary disease and the acute abdomen. I apologize in advance for inevitable omissions but I hope I have incorporated the major advances and developments.

Breast cancer

Breast cancer is common and may be increasing in incidence. A Swedish study has suggested that this is particularly so in younger women with an average annual increase of 3.2% in the age group 25–29 and 3.4% in the 30–34s.¹ Speculation on possible causes include delayed child bearing and the question of oral contraception. Two recent studies have addressed this problem. The WHO collaborative study² compared 2,116 women with newly-diagnosed breast cancer and 12,077 controls. The relative risk of breast cancer in women who ever used oral contraceptives was estimated to be 1.15. Estimates for women under and over 35 were not statistically different. Risk was highest in recent and current users and declined with time since last use. A relative risk of 1.5 (of borderline significance) was observed in women who used oral contraceptives for more than 2 years before age 25. These results could be due to chance or represent a weak causal relationship. The UK National Case-Control Study Group³ previously reported an increased risk of breast cancer associated with oral contraceptive use in women in whom the disease was diagnosed before the age of 36. A further subgroup analysis of this data has shown little evidence of altered risk in relation to subgroups. Women with a family history of breast cancer had a slightly greater risk associated with oral contraceptives than those without, but the difference in trends was not significant. Chilvers *et al.*⁴ have

reviewed all the evidence for the relationship between oral contraceptives and breast cancer. In essence the cumulative risk of breast cancer up to age 36 in the UK is 1 in 500. Even after 8 years of oral contraceptive use this risk would increase to only about 1 in 300. Williams *et al.*,⁵ in a case-control study, reported an increased risk of breast cancer associated with decreasing interval since last term birth, increasing age at first term birth and decreasing parity. The results also suggested a transient increase in the risks of breast cancer after full-term pregnancy.

In selecting optimum treatment for breast cancer it is important to recognize sensitive prognostic indicators; this is particularly so for lymph node negative patients. Two interesting studies have reported on this aspect. The International (Ludwig) Breast Cancer Study Group⁶ looked at serial sectioning of ipsilateral axillary lymph nodes reported as negative and found micrometastases in 9% of 921 patients. These patients had a poorer disease-free and overall survival at 5 years compared to those who remained lymph node negative. Hence detection of micrometastases may identify a high risk 'node-negative' group of patients. Other groups have derived prognostic scores based on various parameters. For example, Chevallier *et al.*⁷ utilized oestrogen receptor, progesterone receptor, tumour size and grade to divide patients into 3 prognostic groups (good, intermediate and bad). Studies such as these, however, require prospective validation and should be tested with appropriate adjuvant regimens. Locker *et al.*⁸ have demonstrated that the cribriform subtypes of symptomatic ductal carcinoma-in-situ (DCIS) is less aggressive than other forms, which may affect management.

The prediction of early relapse and the diagnosis of recurrence is clearly of importance and two new tests have addressed this problem. CA 15-3 is an accurate indicator of synchronous distant metastases in patients with locoregional recurrence⁹ and urokinase-type plasminogen activator (u-PA) may be an independent prognostic factor in predicting early relapse¹⁰ as indeed may urinary tissue factor activity.¹¹

Surgical treatment

The optimum treatment for early breast cancer (T1-2 N 0-1) continues to remain controversial. Limbergen *et al.*¹² reported the results of radiotherapy alone in 221 patients with operable breast cancer. The mean follow-up was 15.5 years. An actuarial local control rate of 75.4% after 15 years was achieved. The dose was dependent on tumour size. An overall disease-free survival of 60.6%, 46.8% and 38% was noted at 5, 10 and 15 years respectively. These data compare favourably with surgical patients but are a selected group and high dose radiotherapy was administered.

Two reports from Milan on breast conservation treatment are reassuring. No differences in disease-free or overall survival were seen in 701 women with a tumour diameter under 2 cm randomized either to radical mastectomy or to quadrantectomy, axillary clearance and breast radiotherapy (QUART). Overall 13 year survivals were 69% and 71% respectively.¹³ Prior to this report, Fisher and colleagues had confirmed equivalent overall survival between mastectomy or breast conservation in women with tumours of 4 cm or less.¹⁴ In the second Milan study,¹⁵ 705 women with cancers less than 2.5 cm were randomized into QUART or more limited treatment consisting of tumorectomy, external radiotherapy and a boost with ¹⁹²Ir implantation (TART). No differences in distant metastases and survival were observed in the two groups although lumpectomy patients had a higher frequency of local recurrences (7.0% vs 2.2%).

The incidence of local recurrence may be related to the permeation of tumour into the breast duct system. Two recent studies have shown that patients with extensive intraduct carcinoma (EIC) in their invasive cancers are twice as likely to have bulky, residual, predominantly intraduct disease in the tumour bed at re-excision.^{16,17} Seventy-one per cent of EIC + ve patients had residual intraduct carcinoma compared to 28% of EIC - ve patients. This may help to explain the association between EIC and subsequent breast recurrence in patients treated with limited excision before radiotherapy.

Additional controversy has recently emerged with regard to the surgical management of the ipsilateral axilla in women with breast cancer. Full axillary clearance is associated with reduced locoregional recurrence and overall survival in some series¹⁸ and has the advantage of avoiding axillary radiotherapy. In addition, important prognostic information is obtained which may be incomplete when only axillary sampling is performed.^{19,20} Du Toit *et al.*,²¹ however, have advocated the triple node biopsy technique, which samples both lateral and medial lymph node chains and the low axillary node. The prognostic inform-

ation is regarded as useful and sufficient to allow better planning of adjuvant systemic treatment.

Andre *et al.*²² have reported locoregional recurrence rates after 649 modified radical mastectomies. The actuarial cumulative locoregional incidence rate was 6%, 14% and 19% after 1, 5 and 7 years respectively. In this series local recurrence was not thought to be a significant sign of imminent generalized disease. Actuarial 5-year survival after local recurrence was 26.2%. These results are similar to those obtained following selective conservation therapy. Barr *et al.*²³ have reviewed the indications and complications of subsequent salvage mastectomy in 25 patients who developed local recurrence after breast-conserving treatment. Two patients required myocutaneous flaps and, in total, 5 patients developed subsequent local relapse and 2 of these patients died with uncontrolled chest wall nodules. Clearly this is associated with a poor overall prognosis.

Systemic therapy

Although adjuvant systemic therapy is now established for post-menopausal women and premenopausal women with lymph node positive tumours, the role of systemic therapy in patients with node - ve breast cancer is still controversial. Two recent studies from the National Surgical Adjuvant Breast Project (NSABP) group have suggested that the oestrogen receptor status of a tumour is not sufficiently sensitive as a prognostic indicator to determine whether adjuvant systemic chemotherapy should be given to node - ve women.²⁴ A longer follow-up is required for these studies to determine whether a true survival advantage exists. A recent review, however, adds an element of caution with regard to the role of adjuvant cytotoxic chemotherapy in node - ve women, and suggests that only women at high risk should be offered this form of treatment. This is probably the view of the majority of UK surgeons but further work and studies are required to establish this contentious point. Two recent reviews both indicate a future for adjuvant chemotherapy^{25,26} but will depend upon further trials and particularly ones which are designed so that biological sub-set interactions can be detected more reliably and variables, such as nuclear pleomorphism, grade, receptor status, S phase fraction, ploidy, and oncogene amplifications, can be taken into consideration. In this way it is likely that subgroups of women who are most likely to respond will be properly treated.

As far as systemic treatment for advanced disease is concerned, two interesting studies have been reported. Harris *et al.*²⁷ have demonstrated, in a randomized trial, that the use of short term or continuous chemotherapy (mitozantrone) is

similar in time to the disease progression, response duration, or survival. Toxicity is milder, however, during a short chemotherapy course. Stopping chemotherapy early in responders did not cause rapid relapse in this study.

Gazet *et al.*²⁸ have performed a randomized trial comparing chemotherapy with endocrine therapy in 60 patients presenting with significantly locally advanced breast cancer. In this small study, chemotherapy was more effective in rapidly reducing the size of the primary breast cancer than endocrine therapy and altered the future management of these patients. However, on completion of follow-up there was no significant difference in disease-free recurrence, or survival.

Breast screening

Several recent studies on breast screening have been reported. The Edinburgh Trial²⁹ reported the results of 45,130 women aged 45–64 randomized between 1979 and 1981. After 7 years of follow-up a mortality reduction of 17% was achieved which was not statistically significant even when corrected for socio-economic status. Women aged 50 years and over had a mortality reduction of 20%. Hobbs *et al.*³⁰ demonstrated that response rate and compliance to treatment were similar in the older and younger women and this should not necessarily be a bar to offering this service.

The management of ductal carcinoma-in-situ (DCIS) is still controversial. Ciatto *et al.*³¹ analysed the presentation and outcome of 156 consecutive cases of DCIS. Different surgical options were used but a temporal trend in favour of conservative treatment was evident. The prognosis for this condition is excellent providing local control is achieved. Subsequent ipsilateral or contra-lateral breast cancer was recorded in only 7 patients. In this study death from breast cancer in DCIS patients could always be attributed to contra-lateral or subsequent ipsilateral infiltrating cancer. This finding confirms the view that local control of DCIS corresponds to cure. In some patients, however, DCIS is extensive and covers a large area. Holland *et al.*³² showed that 66% of tumours involved one breast quadrant, whereas 23% extended over more than one quadrant and 11% were centrally located. Clearly, adequate excision of DCIS may require a wide excision involving up to a whole quadrant.

Finally, two studies have suggested a benefit from cytotoxic wash-out of the tumour bed following excision of the carcinoma. Lucarotti³³ recognized malignant cells on scrape cytology of the excision cavity in 20% of patients following excision. Topical use of antiseptics and cytotoxic agents should destroy these cells. Similarly Park³⁴ compared cytotoxicity of different solutions used

for operative lavage and found that Savlon, noxythioline and povidone iodine were the most effective in reducing the number of recoverable viable malignant cells.

Upper gastro-intestinal surgery

Oesophagus

Reflux oesophagitis continues to be a troublesome problem. Recent randomized trials have compared different pharmacological treatments. Maxton *et al.*³⁵ in a study of 80 patients showed that Pyrogastron and cimetidine are equally effective in healing of oesophagitis and Pyrogastron was marginally better in terms of time to relapse. For patients with refractory oesophagitis Bardhan *et al.*³⁶ reported that omeprazole 40 mg daily for up to 8 weeks was effective in healing and relieving symptoms of erosive oesophagitis refractory to treatment with either cimetidine or ranitidine. Similarly, Dehn *et al.*³⁷ in a randomized double-blind trial demonstrated that omeprazole achieved greater and more rapid symptom relief as well as more rapid endoscopic healing than cimetidine. It would appear, therefore, that omeprazole is establishing itself as the treatment of choice for severe oesophagitis.

However, some patients require an anti-reflux surgical procedure for their symptoms.³⁸ An appraisal of the techniques suggests that surgical correction remains the only feasible long-term solution in selected patients. Of the procedures available the Angelchik anti-reflux prosthesis is becoming more popular. Timoney *et al.*,³⁹ in a review of 44 patients treated over a 5-year period, describe a 74% excellent outcome with 92% of patients having relief of reflux symptoms. However, the results in patients presenting with dysphagia are less good. Mughal *et al.*⁴⁰ studied 126 patients in a prospective study, with preoperative oesophageal manometry. Patients undergoing Nissen fundoplication, obtained satisfactory clinical results in 83.3%. The poor results are due to recurrent reflux or technical failure and unfortunately pre-operative oesophageal studies do not determine the suitability of individual patients for surgical correction.

The treatment of carcinoma of the oesophagus is still highly controversial. The mainstay of therapy for resectable lesions is surgical excision. However, resection is still risky although operative mortality has decreased in incidence from 30% in 1980 to 10–15% in recent years. In many studies the average 5-year survival for patients surviving resection is approximately 12%. Patients with adenocarcinomas fare less well than patients with squamous carcinomas.

An excellent review by Muller *et al.*⁴¹ discusses all

surgical publications on oesophageal carcinoma in the past 10 years; 133 papers were reviewed and an overall postoperative mortality rate of 8% calculated but only 21% of patients were suitable for resection in the total series. In these patients a 5-year survival rate for resectable patients of 11% was found. In addition, the results of adjuvant radiotherapy and chemotherapy have been disappointing. A separate series from South Africa⁴² of 127 patients demonstrated possible benefit with radiotherapy in patients with probable residual tumour. Earlam and Johnson⁴³ have advocated radical radiotherapy not only for inoperable squamous cell carcinoma but also for patients with operable disease. In a small series of 22 patients, a 46% one year and 14% 5-year survival was achieved, which is comparable to surgical reports. In conclusion, it would appear that the operative mortality for surgery in oesophageal carcinoma has improved over the last 10 years and now should be less than 10% in specialist centres. Subsequent survival rates have been remarkably static⁴⁴ and, sadly, trials to compare radiotherapy with surgery for operable squamous cell carcinoma have failed to attract sufficient patient recruitment.⁴⁵

Many patients with oesophageal cancer are treated by palliation only, in an attempt to reduce dysphagia. Oliver *et al.*⁴⁶ demonstrated that radiotherapy adds very little to intubation in terms of improvement in survival but patients receiving radiotherapy have a much longer hospital stay (46 days vs 23 days for intubation alone). Laser therapy has also been used in an attempt to deal with severe dysphagia. A prospective randomized trial in 46 patients showed no significant difference in the quality of life between laser therapy alone and laser therapy followed by endoscopic intubation.⁴⁷ However, the complication rate was significantly higher in intubated patients (10% vs 40%). Clearly, both techniques are effective in dealing with severe dysphagia and the one chosen will depend very much upon local expertise.

Gastro-duodenal

There is increasing evidence to suggest that total eradication of *Helicobacter pylori* is associated with cure of duodenal ulcer. In one recent trial⁴⁸ 4 weeks' treatment of bismuth subcitrate with amoxycillin and metronidazole was associated with healing and no relapse within one year. A further study comparing omeprazole and ranitidine demonstrated an improvement with omeprazole.⁴⁹ Overall, the results tend to suggest that both duodenal and gastric ulcers heal more rapidly and this is explained by a more effective control of gastric acid secretion.

Although it may be thought that surgery now has very little role to play in the management of peptic

ulcer this view may well be premature. Three recent reports have provided data on the results of highly selective vagotomy with a 5 to 20 year follow up. Johnston *et al.*⁵⁰ described the outcome of 600 patients with a recurrent ulceration rate of 15%, 80% of whom had developed symptoms within 5 years and no patient has had recurrence after 13 years. The patient satisfaction rate for Visick grades 1 and 2 was 92%. There was one post-operative death from cerebral haemorrhage. Macintyre *et al.*⁵¹ have similar results in 307 patients without operative mortality. The recurrent ulceration rate was 17.3% but appeared to be surgeon-dependent. Ninety-one per cent of patients felt their symptoms had been cured or greatly improved and 84.9% considered the operation a success. Post-vagotomy symptoms were found in 5% of patients. Finally, Koruth *et al.*⁵² have provided the long term results (8–15 years) of a randomized trial comparing highly selective vagotomy with truncal vagotomy and pyloroplasty. Visick grading was similar in the two groups. The endoscopically proven recurrence rate was 7% after truncal vagotomy and pyloroplasty and 5% after highly selective vagotomy. These are excellent results and support the optimism related to highly selective vagotomy in the management of duodenal ulceration.

Another technique which has recently been submitted to a randomized controlled trial is the operation of anterior lesser curve seromyotomy and posterior truncal vagotomy.⁵³ In a randomized study of 146 patients with a mean follow-up of 4.5 years dumping and diarrhoea were significantly commoner after truncal vagotomy (31 vs 8). The overall Visick grading was better with this procedure.

The management of patients presenting with upper gastrointestinal haemorrhage from peptic ulceration is controversial. Evidence is accumulating that such patients should be admitted to a joint medical/surgical gastrointestinal unit for treatment. Two studies have reported the benefit of such an approach. Holman *et al.*⁵⁴ described such a policy in 430 patients. An overall mortality of 3.7% was reported and all had serious concomitant pathology. The operation rate was 21.6%. Sanderson *et al.*⁵⁵ described very similar data in 292 patients with a mortality of 4.8%. Few other studies have reported such a low incidence. Patients over the age of 60 with gastrointestinal haemorrhage should be operated upon early if mortality is to be reduced.⁵⁶ Branicki *et al.*⁵⁷ have investigated prospectively the factors which determined re-bleeding and mortality in 701 patients. Age over 60, previous medical illness, shock on admission, large ulcer signs and endoscopic stigmata of recent haemorrhage are the most important. These factors must be taken into account in

determining whether individual patients require urgent surgical intervention.

For patients with bleeding gastric ulceration undersewing of the ulcer with biopsy is an effective form of treatment when subsequently placed on H2 receptor antagonist therapy. This is associated with a satisfactory level of mortality and ulcer recurrence.⁵⁸ Hunt and McIntyre,⁵⁹ in a consecutive series of 201 emergency operations for bleeding duodenal ulcer, found that the size of the ulcer was a major factor which correlated with the operative procedure performed. Truncal vagotomy and pyloroplasty combined with under-running of the bleeding point is an acceptable treatment but gastrectomy may be a better procedure if an ulcer of more than 2 cm is present. However, non-surgical procedures should now be considered for patients with massive peptic ulcer haemorrhage. Lin *et al.*⁶⁰ reported the results of a prospective randomized trial of 137 patients with massive peptic ulcer treated by either endoscopic heat probe thermocoagulation or pure alcohol injection. The initial haemostatic effect of the heat probe was better than that of pure alcohol and the duration of hospital stay was shorter. In addition, the incidence of emergency surgery and the subsequent mortality rate was less in the heat-probe group. They concluded that heat probe thermocoagulation should probably be the initial treatment of choice for arrest of massive peptic ulcer haemorrhage.

Gastric cancer

Prognosis in this condition can be improved by the diagnosis of early gastric cancer with subsequent treatment. De Dombal *et al.*⁶¹ reported 90% 5-year survival for this group. Hallissey *et al.*,⁶² in a prospective study, found that 57 patients investigated for 'dyspepsia' had gastric cancer; 15 of these had early cancer. They concluded that investigation of dyspepsia in patients over 40 might increase the proportion of early gastric cancers detected and hence the proportion of operable cases.

Unfortunately, at the time of presentation, the majority have advanced disease. The results of radical gastrectomy (R3) for these patients suggest some benefit. Bozzetti *et al.*⁶³ have described their experience with extended surgery for gastric carcinoma in 143 patients. The operative mortality rate was 15.4% but decreased during the 8 years of the study. The overall 5-year survival was 19%. Clearly, tumour penetration of adjacent structures and nodal involvement were associated with a poor prognosis.

There is undoubted controversy regarding the role of thoraco-abdominal radical gastrectomy for carcinoma of the cardia. Griffin *et al.*⁶⁴ in a series of 38 patients found that such treatment was indicated since the peri-operative mortality rate was

5% (although subphrenic abscesses occurred in 21%) and 23% of patients survived 3 years. This is in contrast to the experience of Wilson *et al.*⁶⁵ in whom a dismal prognosis was achieved in a series of 47 patients. Only 20% of their series survived one year with a mean duration of 5.4 months. In view of such poor outcome alternative forms of treatment have been suggested. Gill *et al.*⁶⁶ reported their experience in 29 patients with adenocarcinoma of the cardia of the stomach treated by synchronous chemotherapy (5-fluorouracil and cisplatin) and radiotherapy. Subsequently 14 patients had the tumour resected and in 6 no microscopic tumour was found in the specimen. The median survival of patients who responded was 15 months. Such synchronous chemotherapy and radiotherapy has valuable palliative effects and perhaps should be considered more frequently in this condition. The importance of quality of life following surgery for gastric carcinoma should not be forgotten. Buhl *et al.*⁶⁷ have evaluated this aspect in 89 patients treated either by total gastrectomy, distal gastrectomy or proximal gastrectomy. Proximal gastrectomy was associated with most troublesome sequelae and had adverse effects on quality of life. Jejunal pouch construction, however, may be a more satisfactory procedure to alleviate disabling symptoms.⁶⁸

Biliary disease

A major revolution has occurred in the management of patients with gallstones. A recent review on this topic⁶⁹ has highlighted the pros and cons of the different techniques. Undoubtedly laparoscopic cholecystectomy is becoming increasingly popular and widespread. Two editorials^{70,71} have emphasized the advantages in terms of reduced hospital stay and low morbidity. It would be difficult to justify a randomized trial of conventional versus laparoscopic cholecystectomy, and attempts to initiate this have failed.⁷² Nevertheless, comparative studies and careful audit are required and two recent reports have followed this approach. Nathanson *et al.*⁷³ have reported experience in Dundee on 61 patients. Laparoscopic cholecystectomy was completed in 60 patients with a low morbidity rate and a median duration of surgery, including cholangiogram, of 135 minutes (range 65–270 minutes). The median hospital stay was 2.9 days and the median time to resume full work after discharge was 11 days. Grace *et al.*⁷⁴ produced very similar data in 50 consecutive patients. The median operating time was 155 minutes, but the median postoperative hospital stay was 3.5 days compared to 8.8 days following conventional surgery. They conclude that the technique is associated with financial benefits and is cosmetically satisfactory. I

think few individuals would disagree with this.

Piezoelectric lithotripsy has been advocated but only 28% of patients with gallstone disease appear to fulfill the criteria for treatment by this technique.⁷⁵ Another disadvantage relates to the need for subsequent dissolution therapy over a long period of time. Complications from this technique are few and indeed a recent study has happily demonstrated that soft tissue injuries are minimal.⁷⁶ Comparison has been performed in extracorporeal shock wave lithotripsy between an electrohydraulic and a piezoceramic device *in vitro*.⁷⁷ All stones were successfully disintegrated and the actual volume of the stone seemed to be the only determinant which affected ease of fragmentation.

Chiverton *et al.*⁷⁸ have assessed percutaneous cholecystolithotomy in 60 patients as an alternative method for removing gallstones. Four patients required subsequent cholecystectomy. Seven patients had postoperative complications and 2 had recurrence of biliary calculi. This is a technique which might be suitable for elderly patients who have empyema and significant septic problems that are unfit for major abdominal surgery. A further technique which has been described is the use of methyl-tert-butyl-ether (MTBE) infused via a naso-biliary catheter into the common bile duct to treat bile duct stones. A recent report⁷⁹ has described this technique in 33 patients. It is a technique with a significant complication rate and is unlikely to be used widely.

The question of visualization of the common bile duct at operation is always a subject of dispute. Motson *et al.*⁸⁰ advocate the use of operative choledochoscopy, either rigid or flexible, in patients following gallstone removal and regard this as a mandatory part of common bile duct exploration. Many surgeons now regard operative cholangiography as a procedure which should be carried out selectively in patients who fulfill certain criteria, for example, jaundice, cholangitis, pancreatitis and dilated common bile duct. A recent study by Pernthaler *et al.*⁸¹ tends to confirm this view, but also recommends that small stones should be accepted as an indicator for cholangiography.

There is continuing controversy also with regard to endoscopic removal of common bile duct stones. This particularly relates to common bile duct stones in young patients with a gallbladder *in situ*. The current indications and controversies have been reviewed by Horton *et al.*⁸² The major indications remain acute obstructive suppurative cholangitis, acute cholangitis and duct stones after cholecystectomy. Worthley *et al.*⁸³ have compared common duct exploration in 102 patients with 50 patients who had endoscopic sphincterotomy. They conclude that the optimum treatment for post-cholecystectomy patients with bile duct stones

is endoscopic sphincterotomy and few would disagree with this. In elderly patients with an intact gallbladder the bile ducts can also be treated by endoscopic sphincterotomy.

Pancreatic disease

There are several selective enzyme and blood tests available both for the diagnosis and subsequent prognosis of acute pancreatitis. Nevertheless, serum amylase and lipase measurements still represent the best available diagnostic methods.⁸⁴ The other newer techniques require careful evaluation. One of these has been described by Gudgeon *et al.*⁸⁵ and is a trypsinogen activation peptide assay. It has been carried out in 55 patients with acute pancreatitis and its sensitivity and specificity were 80% and 90%. It is also regarded as an important indicator of severe acute disease.

The majority of patients with acute pancreatitis settle with conservative treatment but undoubtedly a number require more aggressive surgical management, often associated with a poor postoperative course. One of the most serious situations relates to pancreatic abscess formation. Bassi *et al.*⁸⁶ reviewed 108 patients with pancreatic abscess formation treated between 1970 and 1987. The overall mortality rate was 15.7% but this was particularly marked (23.6%) in patients with spreading abscess formation rather than a localized disease. The management of such patients can be extremely difficult and requires resection or necrosectomy with extensive local lavage and drainage, often utilising 'open' management. Two reviews have recently focussed on the different techniques.^{87,88} Some patients require retroperitoneal drainage to avoid contamination of the peritoneal cavity and enable adequate debridement to occur.⁸⁹

Total parenteral nutrition has a role to play in the management of acute pancreatitis. Robin *et al.*⁹⁰ reviewed their experience in 156 patients who required total parenteral nutrition. The usual indications are complex local disease, such as abscess or pseudocyst, and those patients who have prolonged ileus without oral intake for many days. Total parenteral nutrition can be administered safely but careful monitoring is necessary.

The management of chronic pancreatitis remains extremely problematical. This is particularly so in patients with severe pancreatic pain. The relatively large number of procedures probably reflects a dissatisfaction with the effects of each of them. Percutaneous blockage of the coeliac plexus with alcohol and phenol are commonly used to provide short-term pain relief. In patients with dilated ducts pancreatico-jejunostomy is regarded as the method of choice with prolonged pain relief in about 60%.⁹⁰ Undoubtedly, careful assessment of each

patient is required and if it is certain that they have abstained from alcohol then operative procedures, depending upon the results of pancreatography, are of value in selected cases.⁹²

Carcinoma of the pancreas remains a depressing condition. Hopefully, improvements in imaging, tissue sampling and better tumour associated antigens will enable an earlier diagnosis to be made. Improvements in surgical technique have decreased postoperative morbidity. Morel *et al.*⁹³ have described the results of 33 pylorus-preserving duodenopancreatectomies with no postoperative mortality. There was a 24% morbidity rate associated with anastomotic leaks, surgical bleeding and anastomotic ulceration. Sixty-two percent of patients were alive with a mean survival time of 20 months. Hall *et al.*⁹⁴ have described a sutureless pancreatico-jejunostomy which is performed during pancreatico-duodenectomy. There was an acceptably low leakage rate and the majority of patients required no enzyme supplements.

Nevertheless, the majority of patients still undergo surgical palliation only. Whether this should be by surgery or stent is still controversial and has been reviewed by Hatfield.⁹⁵ It is suggested that younger, fitter patients who are likely to survive more than 6 months are probably better treated by surgical bypass, leaving endoscopic stenting for the elderly and frail with advanced disease. Two separate studies have suggested that biliary bypass alone is sufficient and the addition of gastroenterostomy almost doubles mortality without any obvious advantage in terms of survival. Clearly, if there is an obstructive gastric outlet element, gastroenterostomy is indicated but not in all patients.^{96, 97} Bickerstaff *et al.*⁹⁸ have advocated palliative endoscopic sphincterotomy in 17 patients with carcinoma of the ampulla of Vater. An endoprosthesis was inserted immediately afterwards. The median survival was 12 months and thus this represents reasonable palliation, which can be carried out safely. Kahn *et al.*⁹⁹ have reported their experience with a combination of surgical palliation and synchronous treatment consisting of intravenous 5-fluorouracil and radiotherapy. This group of patients had a mean survival of 13.5 months compared to 8.9 months in patients who received bypass only. This was not a randomized trial but tends to suggest that in selected patients synchronous treatment might be of some benefit. Finally, the results of a randomized trial of 176 patients with unresectable pancreatic carcinoma who were randomized either to tamoxifen or placebo have been reported.¹⁰⁰ There was no significant difference in either mean or median survival between the two groups of patients. Three women in the tamoxifen group were alive compared to no survivors in the placebo group after 2.5 years. This is an interesting finding but does not

detract from the overall negative results obtained in this study.

Liver

Portal hypertension

The management of patients with bleeding varices in association with portal hypertension continues to be a major problem. Haemodynamic aspects are of importance in management and the correlation with physiology and pathology has been reviewed by Karran.¹⁰¹ Non-surgical treatments have attractions. It is still controversial, however, whether beta adrenoreceptor blocking drugs such as propranolol have a role in this condition. A recent meta-analysis of all controlled clinical trials on the use of propranolol indicate possible benefits in reducing death from haemorrhage, although further trials and comparisons will be required before this problem is solved. Nevertheless, this is a safe treatment and may reduce the risk of bleeding or re-bleeding by up to 40%.¹⁰²

Injection sclerotherapy is still very popular and remains a major therapeutic modality. Such treatment is valuable not only in the acute situation but as long-term control of oesophageal varices. Kahn *et al.*¹⁰³ have reviewed the experience of injection in young patients with extra-hepatic portal venous obstruction. This is a most difficult condition to treat adequately but over a mean period of 14.5 months and 7 injections varices were eradicated in 33 patients. Complications were minor and therefore this therapy should be regarded as a treatment of choice for this condition.

Gastric varices can also be extremely difficult to deal with. Mathur *et al.*¹⁰⁴ reviewed their endoscopic and radiological appraisal of varices in 81 patients. They were more commonly seen along the lesser curvature and bleeding occurred in 9 of these patients more significantly from the fundus than from the lesser curve. The optimum management for such patients is unknown. Greig *et al.*¹⁰⁵ have reported their experience in 9 patients, 6 of whom had been followed for 2 years or longer. Under-running of the gastric varices alone seemed unsatisfactory and left gastric vein ligation was indicated in addition.

Undoubtedly, liver transplantation is being carried out more commonly in patients with cirrhosis and associated portal hypertension. Bird *et al.*¹⁰⁶ have reviewed the selection criteria, survival and relapse in patients undergoing transplantation for alcoholic cirrhosis. Survival and rehabilitation in these patients compares favourably with patients suffering from other forms of cirrhosis. Patients selected are those with recurrent complications despite medical treatments, but in addition, there

should be a minimum period of 6 months of abstinence before transplantation is considered. As far as primary biliary cirrhosis is concerned, Neuberger *et al.*¹⁰⁷ have reviewed the referral criteria for this group. Their recommendation appears to be that potential candidates should be referred before the serum bilirubin concentration approaches 150 $\mu\text{mol/l}$.

Liver tumours

What is the most effective imaging modality in patients with liver tumours? A prospective comparison of ultrasonography, CT scan and angiography was performed in 54 patients and the results compared to intraoperative ultrasound.¹⁰⁸ Intraoperative ultrasound demonstrated 25% to 35% additional lesions, 40% of which were neither visible nor palpable at surgery. These authors therefore recommend intraoperative ultrasound in all patients in whom resection is planned. Increasing interest, however, has been shown in Lipiodol computed tomography for very small hepatocellular carcinomas and undoubtedly this is a technique which will gain in importance.¹⁰⁹

Several recent reports have reviewed the optimum surgical management for primary hepatocellular carcinoma, particularly those complicating cirrhosis. In general the long-term results of resection of hepatocellular carcinoma have been disappointing, principally because of the high rate of intrahepatic recurrence. Belghiti *et al.* have reported a cumulative recurrence rate of 100% at 5 years.¹¹⁰ Patients presenting with advanced disease unsuitable for resection and confined to the liver and in whom conventional hepatic resection is not possible, might be suitable for liver transplantation. Ismail *et al.*¹¹¹ have reported the Birmingham series of 29 grafts performed in 28 patients. Twenty patients have survived more than 30 days with a median survival of 11.5 months. Ringe *et al.*¹¹² and Otto *et al.*¹¹³ have provided similar data from Hanover and Heidelberg respectively. A one year survival rate of 54% in selective patients has been reported. In general these results are not as satisfactory as those obtained in patients who are suitable for resection. Al-Hadeedi *et al.*¹¹⁴ have reported a 33% 2-year survival and 22% 3-year survival following extended hepatectomy for hepatocellular carcinoma. Even better results from a selective policy have been reported by Paquet *et al.*¹¹⁵ and Hasegawa *et al.*¹¹⁶ This latter author has described central bisegmentectomy in patients with advanced disease.

Mortality in patients with bleeding liver tumours is very high and perhaps this condition is unsuitable for any form of therapy. Dewar *et al.*¹¹⁷ however, have reported the experience in 41 patients with spontaneous liver rupture over a 4-year period.

Various techniques were utilized, including angiographic embolization and intratumoral injection of absolute alcohol. Some benefits are reported, but the ultimate prognosis is very poor.

Colorectal liver metastases are very common but the optimal management of such patients is still debated. Karanjia *et al.*¹¹⁸ have highlighted this problem by reporting the results of a questionnaire sent out to consultant surgeons in the United Kingdom. No consensus was achieved as to which patients would benefit from resection, and only 62% said they would consider surgery even in a patient with a solitary metastasis. The major source of differing opinion relates to the natural history of this disorder and the accurate assessment of outcome following surgical resection compared to no treatment. Scheele *et al.*¹¹⁹ have discussed this problem based on an extensive review of almost 1,000 patients. Radical resection of disease confined to one lobe irrespective of the number of metastases in the lobe appeared to prolong survival. They suggest that the outcome can be altered by such an approach but this is not agreed universally.¹²⁰ The debate continues. It would appear that survival rates are not significantly different in some series after resection for solitary compared to multiple liver metastases (26 vs 28 months). Operative mortality from such procedures should be less than 10%.^{121,122}

However, over 90% of patients have multiple metastases involving both lobes of the liver which makes resection impossible. Hunt *et al.*¹²³ have carried out a prospective randomized trial in such patients and demonstrated that long-term infusion chemotherapy with 5-fluorouracil and degradable starch microspheres may be of benefit in enhancing median survival. In patients with less than 50% involvement a median survival of almost 2 years was achieved. Other techniques involve targeting liver metastases with intrahepatic chemotherapy by the use of vasoactive agents to redistribute arterial blood towards tumours. Such treatment consists of angiotension II and albumin microspheres.¹²⁴

Trauma

Major blunt liver trauma is still associated with a high mortality rate. Andersson *et al.*¹²⁵ have reviewed the conservative approach to liver trauma and suggest that up to 80% of patients with liver injuries can be managed by simple measures. However, selection of patients and monitoring by CT scan is essential. In addition, patients should be haemodynamically stable and selective arterial embolization may be required. Nevertheless, some patients will require major hepatic resection and have more serious injuries with a higher mortality. Hollands and Little¹²⁶ have reported experience in 42 patients with an overall operative mortality of

36%. The common complications are respiratory infections, coagulopathy and sepsis.

The most serious injury is damage to the hepatic vein and retrohepatic vena cava. These injuries are extremely difficult to deal with and the majority of patients die. Ciresi *et al.*¹²⁷ however, have reported successful techniques using vascular isolation of the liver in 27 patients, 12 of whom were successfully resuscitated.

Colorectal disease

Coloproctology is an ever expanding specialty. This brief review will deal with two specific aspects; cancer and inflammatory bowel disease.

Colorectal cancer

The aetiology of this very common cancer remains obscure. Nonetheless, increasing understanding of its growth and spread are becoming apparent. Rew *et al.*¹²⁸ have investigated the *in vivo* proliferative characteristics of human colorectal cancer using bromodeoxyuridine. Kinetic relationships to extent of spread have been demonstrated. It has been suggested that differential expression of oncogenes may reflect the way in which colorectal tumours respond to other growth factors. Studies supporting the concept of an enhanced expression include p62 c-myc oncogene¹²⁹ and p53 as prognostic indicators.¹³⁰ Interestingly, there may be a correlation between the chromosome deletions which have been described and the different mechanisms of c-myc overexpression.¹³¹ Growth factors may well have a role and epidermal growth factor receptor (EGFr) expression has been measured in invasive human colorectal cancer using immunohistochemistry. Whereas all tumours stain for EGFr, Dukes C tumours have demonstrated higher levels of receptors than either Dukes A or B and may therefore be an indication of poor prognosis.^{132,133}

It has been recognized for many years that an association exists between malignancy and alterations in haemostasis. Tumours have procoagulant activities and this can be clearly demonstrated in colorectal cancer and may be important not only in local spread but in determining the presence of metastases.¹³⁴ Normal colonic epithelial cells have cell associated urokinase-type plasminogen activity which may be enhanced in malignancy.¹³⁵ Similarly, urinary tissue factor activity is enhanced in colorectal malignancy.¹³⁶

Diet has always been thought to be of importance in the aetiology of large bowel cancer and further evidence supports this.¹³⁷ It would appear that fatty acids are implicated in some way and certainly arachidonic acid and docosahexaenoic

acid are elevated in human colorectal cancer.^{138,139} These studies suggest that high saturated fat promotes the malignant transformation of colorectal adenomas into carcinomas. Bile acids may also have a role and are increased following truncal vagotomy which might be associated with an increased incidence in colorectal malignancy.¹⁴⁰ In addition, bile acids which influence cellular proliferation are abnormal in patients with familial adenomatous polyposis.¹⁴¹

Another aspect which is gaining attention is the study of the intercellular connections between cells. Collins *et al.*¹⁴² have studied the pattern of desmosomes in colorectal cancer and noted alterations in the presence of low calcium.

The prognosis for colorectal cancer could be improved if early diagnosis was made before the cancer had spread. Numerous studies on screening for colorectal cancer using faecal occult blood are continuing and the results of these studies are awaited with interest. In order to increase specificity Kewenter *et al.*¹⁴³ have demonstrated that a rehydrated haemoccult test should be carried out only in those patients with a possible result. This procedure might significantly reduce the work load without necessarily reducing sensitivity. Similarly, an increase in the yield of neoplasia can be obtained if the haemoccult testing is carried out over 6 days rather than 3 even though compliance may be reduced and the number of colonoscopies increased.¹⁴⁴ The most effective method of screening has yet to be established and continuing studies comparing different methods in an attempt to improve specificity and sensitivity are continuing.¹⁴⁵ In spite of this, the controversy as to the potential benefits of colorectal screening remain. Recent reports from Japan suggest significant benefits,¹⁴⁶ whereas other studies have questioned the concept in making any major impact on prognosis.^{147,148} Whether patients should be screened after resection for colorectal cancer to determine early recurrence is equally debatable. Juhl *et al.*¹⁴⁹ suggest annual colonoscopy for at least 6 years post-resection, not only to detect new primary tumours, but also predisposing lesions. Various tumour markers have also been suggested in this role and these include CEA, TPA and CA 19-9.¹⁵⁰

The role of immunoscintigraphy in the diagnosis of colorectal cancer, particularly in the recognition of early recurrence, continues to attract interest. A successful technique would indeed lend itself to therapeutic manipulation. Various monoclonal antibodies have been used for this and include PR1A3,¹⁵¹ carcinoembryonic antigen (CEA)¹⁵² and combinations such as fragments of monoclonal antibodies against CEA and cancer antigen 19-9 (CA 19-9).¹⁵³ The accuracy of all these techniques is doubtful but improvements and technical modifications may provide this technique with

therapeutic benefits. A further diagnostic modality is endorectal ultrasonography to determine whether lymph nodes are involved and the extent of penetration of tumour.¹⁵⁴ Glaser *et al.*¹⁵⁵ reported accurate identification of lymph nodes in approximately 80% and a sensitivity for detection of peri-rectal infiltration of 97%.

The results of surgical treatment alone remain fairly static. This was illustrated by McArdle *et al.*¹⁵⁶ who looked at the 10-year follow-up of colorectal cancer in Scotland. At the time of presentation 50% of patients had clinical evidence of tumour fixity, and 25% had distal metastases. The overall resectability rate was 70.8%. Operative mortality was 13.8% and approximately 70% of patients undergoing curative resection survived for 2 years and 50% for 5 years. This study illustrates once again the advanced nature of the disease at presentation and the high incidence of emergency presentations. Specialist centres, however, do obtain more favourable results as illustrated particularly for rectal cancer. Karanjia *et al.*¹⁵⁷ have demonstrated the importance of total mesorectal excision which may be more important than a reduction in resection margins. Other authors have also emphasized the importance of the extent of mesorectal spread in prognosis with involvement of the lateral resection margin. Cawthorn *et al.*¹⁵⁸ measured mesorectal spread in 167 consecutive patients. There were significant differences in survival between patients with slight and extensive mesorectal spread in Dukes B tumours (66% vs 37%). Indeed, an editorial in the *Lancet* has recently stressed the importance of mesorectal excision in rectal cancer.¹⁵⁹

Other points of technical interest which may help to improve the overall prognosis include a demonstration that, following Hartmann's procedure, early colorectal re-anastomosis should be embarked upon and may be an easier procedure to perform;¹⁶⁰ peroperative observation of marginal artery bleeding may be a useful predictor of anastomotic leakage;¹⁶¹ and finally the role of endoscopic transanal resection of rectal tumours in patients who are either elderly and/or undergoing palliative procedures may well be of great value.¹⁶²

Left-sided malignancy frequently presents with large bowel obstruction. Surgical management of such patients varies but increasing evidence suggests that primary restorative colectomy is justified with acceptable morbidity. Two options are available for this. There are advocates of primary anastomosis following intraoperative colonic washout and acceptable results are reported.¹⁶³ Alternatively, and particularly if the tumour is in a proximal descending colon, sub-total colectomy and primary anastomosis without the need for intraoperative washout can be used with a low morbidity, mortality and an acceptable hospital

stay.¹⁶⁴ Other techniques include a 3 stage operation or a 2 stage procedure, for example, Hartmann's operation. Runkel *et al.*¹⁶⁵ have reviewed the outcome after emergency surgery in Germany. Various procedures have been undertaken and primary resection was performed in 38% of left-sided tumours. An overall mortality in their series of 23% was reported.

There has been increasing interest also in the use of laser for the primary treatment of colorectal cancer and particularly in palliation. Loizou *et al.*¹⁶⁶ presented data on 49 patients with rectosigmoid carcinoma who were unsuitable for surgery but who underwent endoscopic laser therapy for palliation of symptoms and tumour eradication. The results were pleasing in that symptomatic improvement was achieved in all patients with tumours less than 3 cm in diameter and in patients with larger tumours symptomatic improvement was achieved in 74%. Mean hospital stay was 9 days. Other encouraging results have been achieved by von Dittfurth *et al.*¹⁶⁷ in patients with obstruction and bleeding. It may well be that endoscopic photodynamic therapy will be even more effective in selected patients.¹⁶⁸ Early results suggest that the technique is feasible and patients with extensive recurrence can avoid further surgical intervention.

There has been increasing interest in adjuvant chemotherapy in colorectal cancer. Indeed, in the UK a national trial, known as the AXIS Study, has been undertaken to review the effect of adjuvant portal vein infusion and radiotherapy in colon and rectal cancer. A large accrual into this study is required if accurate information is eventually to be obtained.¹⁶⁹ The combination of levamisole and 5-fluorouracil has been associated with an improvement of survival in patients with Dukes C colorectal cancer in two studies.^{170,171} This latter study is particularly important and was associated with a reduction in the risk of recurrence by up to 40%. Indeed, the overall death rate was reduced by 33%. Toxic side effects were limited. Treatment, however, continues for up to 1 year. The results of a prospective randomized trial of systemic 5-fluorouracil, vincristine and CCNU for Dukes C colorectal cancer did not provide any worthwhile improvement in recurrence rates or survival.¹⁷²

Inflammatory bowel disease

Two interesting topics associated with this condition have been reported. Restorative proctocolectomy and ileoanal anastomosis is now becoming accepted as a standard treatment for ulcerative colitis. Several groups have reported experience with this particular procedure. Jarvinen *et al.*¹⁷³ have demonstrated that, in many patients, a covering ileostomy is unnecessary for this procedure, with obvious advantages. A continuing problem of

pouch surgery is the development of inflammation within the pouch itself – ‘pouchitis’. The aetiology of this is still the subject of speculation but is likely to be inflammatory in nature. The various theories on aetiology have been reviewed by Madden *et al.*¹⁷⁴ Several factors will determine whether or not function within an ileoanal pouch is satisfactory.¹⁷⁵ These are both morphological and physiological and require consideration before the procedure is undertaken.

A continuing concern in ulcerative colitis is the subsequent development of malignancy. There appears to be a relationship between the age of onset of ulcerative colitis and age of diagnosis of cancer. Patients with left-sided colitis develop cancer about 10 years later than those with extensive disease.¹⁷⁶ Other studies suggest that patients with extensive colitis have clinically important cancer risk after the disease has been present for about 10 years. The cumulative probability of developing cancer was 3% at 15 years and 9% at 25 years.¹⁷⁷ Screening for colorectal cancer has been advocated but the benefits are dubious and the costs are high. It is likely that only by means of a randomized trial will this question be answered.¹⁷⁸

It should also be remembered that patients with Crohn's disease involving the colon also have an increased risk of large bowel cancer. Ekbohm *et al.*¹⁷⁹ have reviewed 1,655 patients with Crohn's disease and showed an increased overall risk of malignancy of 2.5. Patients with Crohn's disease who were diagnosed before the age of 30 with colonic involvement at diagnosis had a higher relative risk (20.9%) compared to those diagnosed at older ages (2.2%).

Sadly, Crohn's disease is associated with a high incidence of recurrence and radical resection does not appear to reduce the incidence.¹⁸⁰ With improving treatment there has been a significant decrease in the mortality over the last 20 years.¹⁸¹

The acute abdomen

It is becoming increasingly apparent that general surgery will be subdivided into specialties. However, one aspect which will continue to be the province of the ‘general surgeon’ is the diagnosis and subsequent management of the patient admitted as an emergency with an ‘acute abdomen’. Diagnosis depends predominantly on careful history taking and examination performed by experienced doctors who are able to make accurate interpretations of the physical signs. Nevertheless,

the decision to operate or to treat conservatively can be extremely difficult and errors are made. Paterson-Brown and Vipond¹⁸² have reviewed the modern aids in clinical decision-making for patients with an acute abdomen. These include computer-aided diagnosis, laparoscopy, peritoneal cytology and ultrasonography. Under certain circumstances white cell scanning can also help. However, operator variation and difficulties in providing a 24-hour service prevent them from becoming first-line investigations in most hospitals.

Ultrasonography has been assessed in recent studies and particularly in a large review by Ooms *et al.*¹⁸³ Ultrasonography with graded compression was performed in 525 patients with clinical signs of acute appendicitis. As experience improved during the last 3 years of the study the negative appendectomy rate dropped to 7% and delay beyond 6 hours after admission occurred in only 2% of patients with surgically proven appendicitis. It is thought that this test may be used to complement clinical accuracy. Vipond *et al.*¹⁸⁴ have evaluated fine-catheter aspiration cytology of the peritoneal cavity in 61 patients who were admitted with acute abdominal pain. Aspirates were examined microscopically. In patients in whom the clinical need for operation was uncertain (36), 19 required operation. The peritoneal neutrophil count was more than 50% in 18 and in one patient with an ectopic pregnancy fresh blood was aspirated. Again, this study confirms that peritoneal cytology is a useful adjunct in cases when an acute abdomen is difficult and the decision to operate is in doubt.

Non-specific abdominal pain, however, is still the commonest cause of acute abdominal pain in Western countries and positive diagnosis in such patients can be extremely difficult. Raheja *et al.*¹⁸⁵ prospectively studied 80 such patients. Viral studies failed to show any abnormality and psychological studies demonstrated that this group of patients had similar levels of anxiety and depression to a control group. There is no doubt that more serious conditions need to be excluded before this diagnosis is made. It is still responsible for a high proportion of negative laparotomies.

A recent review of the practicalities management of the acute abdomen has emphasized the importance of an all-day emergency theatre to deal with operations which otherwise would have had to be performed during the night but could in fact be safely deferred to the following morning.¹⁸⁶ This administrative aspect in the management of the acute abdomen is most important and will ensure a better level of overall surgical care and attention.

References

Breast cancer

1. Ranstam, J., Janzon, L. & Olsson, H. Rising incidence of breast cancer among young women in Sweden. *Br J Cancer* 1990, **61**: 120–122.
2. The WHO Collaborative Study of Neoplasia and Steroid Contraceptives. Breast cancer and combined oral contraceptives: results from a multinational study. *Br J Cancer* 1990, **61**: 110–119.
3. UK National Case-Control Study Group. Oral contraceptive use and breast cancer risk in young women: subgroup analyses. *Lancet* 1990, **335**: 1507–1509.
4. Chilvers, C.E.D. & Deacon, J.M. Oral contraceptives and breast cancer. *Br J Cancer* 1990, **61**: 1–4.
5. Williams, E.M.I., Jones, L., Vessey, M.P. *et al.* Short term increase in risk of breast cancer associated with full-term pregnancy. *Br Med J* 1990, **300**: 578–579.
6. International (Ludwig) Breast Cancer Study Group. Prognostic importance of occult axillary lymph node micrometastases from breast cancers. *Lancet* 1990, **335**: 1565–1568.
7. Chevallier, B., Morseri, V., Dauce, J.P. *et al.* A prognostic score in histological node negative breast cancer. *Br J Cancer* 1990, **61**: 436–440.
8. Locker, A.P., Horrocks, C., Gilmour, A.S. *et al.* Flow cytometric and histological analysis of ductal carcinoma *in situ* of the breast. *Br J Surg* 1990, **77**: 564–567.
9. O'Dwyer, P.J., Duffy, M.J., O'Sullivan, F. *et al.* CEA and CA 15–3 in primary and recurrent breast cancer. *World J Surg* 1990, **14**: 562–566.
10. Janicke, F., Schmitt, M., Hafer, R. *et al.* Urokinase-type Plasminogen activator (u-PA) antigen is a predictor of early relapse in breast cancer. *Fibrinolysis* 1990, **4**: 69–78.
11. Carty, N., Taylor, I., Roath, O.S. *et al.* Urinary tissue factor activity in malignancy. *Thromb Res* 1990, **57**: 473–478.
12. Van Limbergen, E., Van der Schueren, E., Van den Bogaert, W. *et al.* Local control of operable breast cancer after radiotherapy alone. *Eur J Cancer* 1990, **26**: 674–679.
13. Veronesi, U., Banfi, A., Salvadori, B. *et al.* Breast conservation is the treatment of choice in small breast cancer: long-term results of a randomized trial. *Eur J Cancer* 1990, **26**: 688–670.
14. Fisher, B., Redmond, C., Poissen, R. *et al.* Eight-year results of a randomised clinical trial comparing total mastectomy and lumpectomy with or without irradiation in the treatment of breast cancer. *N Engl J Med* 1990, **320**: 822–828.
15. Veronesi, U., Volterrani, F., Luini, A. *et al.* Quadrantectomy versus lumpectomy for small size breast cancer. *Eur J Cancer* 1990, **26**: 671–673.
16. Jacquemier, J., Kurtz, J.M., Amalric, R. *et al.* An assessment of extensive intraductal component as a risk factor for local recurrence after breast-conserving therapy. *Br J Cancer* 1990, **61**: 873–876.
17. Holland, R., Connolly, J.L., Gelman, R. *et al.* The presence of an extensive intraductal component following a limited excision correlates with prominent residual disease in the remainder of the breast. *J Clin Oncol* 1990, **8**: 113–118.
18. Matthei, W., Bourgeois, P., Delcorde, A. *et al.* Axillary dissection in breast cancer revisited. *Eur J Surg Oncol* 1989, **15**: 490–495.
19. Fentiman, I.S. & Mansel, R.E. The axilla: not a no-go zone. *Lancet* 1991, **337**: 221–223.
20. O'Dwyer, P.J. Axillary dissection in primary breast cancer. The benefits of node clearance warrant appraisal. *Br Med J* 1991, **302**: 360–361.
21. Du Toit, R.S., Locker, A.P. & Ellis, I.O. Evaluation of the prognostic value of triple node biopsy in early breast cancer. *Br J Surg* 1990, **77**: 163–167.
22. Andry, G., Suciu, S., Vico, P. *et al.* Locoregional recurrences after 649 modified radical mastectomies: incidence and significance. *Eur J Surg Oncol* 1989, **15**: 476–485.
23. Barr, L.C., Phillips, R.H., Brunt, A.M. *et al.* Salvage mastectomy after failed breast-conserving therapy for carcinoma of the breast. *Ann R Coll Surg Engl* 1991, **73**: 126–129.
24. Fisher, B., Redmond, C., Wickerman, D.L. *et al.* Systemic therapy in patients with node-negative breast cancer. A commentary based on two national surgical adjuvant breast and bowel project (NSABP) clinical trials. *Ann Intern Med* 1989, **111**: 703–712.
25. O'Reilly, S.M. & Richards, M.A. Node negative breast cancer. Adjuvant chemotherapy should probably be reserved for patients at high risk of relapse. *Br Med J* 1990, **300**: 346–348.
26. Mittra, I. Has adjuvant treatment of breast cancer had an unfair trial? *Br Med J* 1990, **301**: 1317–1319.
27. Harris, A.L., Cantwell, B.M.J. & Carmichael, J. Comparison of short-term and continuous chemotherapy (mitozantrone) for advanced breast cancer. *Lancet* 1990, **335**: 186–190.
28. Gazet, J.-C., Ford, H.T. & Coombes, R.C. Randomised trial of chemotherapy versus endocrine therapy in patients presenting with locally advanced breast cancer (a pilot study). *Br J Cancer* 1991, **63**: 279–282.
29. Roberts, M.M., Alexander, F.E., Anderson, T.J. *et al.* Edinburgh trial of screening for breast cancer: mortality at seven years. *Lancet* 1990, **335**: 241–246.
30. Hobbs, P., Kay, C., Friedman, E.H.I. *et al.* Response by women aged 65–79 to invitation for screening for breast cancer by mammography: a pilot study. *Br Med J* 1990, **301**: 1314–1316.
31. Ciatto, S., Grazzini, G., Iossa, A. *et al.* *In situ* ductal carcinoma of the breast – analysis of clinical presentation and outcome in 156 consecutive cases. *Eur J Surg Oncol* 1990, **16**: 220–224.
32. Holland, R., Hendriks, J.H.C.L., Verbeek, A.L.M. *et al.* Extent, distribution, and mammographic/histological correlations of breast ductal carcinoma *in situ*. *Lancet* 1990, **335**: 519–522.
33. Lucarotti, M.E., White, H., Deas, J. *et al.* Antiseptic toxicity to breast carcinoma in tissue culture: an adjuvant to conservation therapy? *Ann R Coll Surg Engl* 1990, **72**: 388–392.
34. Park, K.G.M., Chetty, U., Scott, W. *et al.* The activity of locally applied cytotoxics to breast cancer cells *in vitro*. *Ann R Coll Surg Engl* 1991, **73**: 96–99.

Upper gastrointestinal surgery

35. Maxton, D.G., Heald, J., Whorwell, J. *et al.* Controlled trial of Pyrogastron and cimetidine in the treatment of reflux oesophagitis. *Gut* 1990, **31**: 351–354.
36. Bardhan, K.D., Morris, P., Thompson, M. *et al.* Omeprazole in the treatment of erosive oesophagitis refractory to high dose cimetidine and ranitidine. *Gut* 1990, **31**: 745–749.
37. Dehn, T.C.B., Shepherd, H.A., Colin-Jones, D. *et al.* Double blind comparison of omeprazole (40 mg od) versus cimetidine (400 mg qd) in the treatment of symptomatic erosive reflux oesophagitis, assessed endoscopically, histologically and by 24 h pH monitoring. *Gut* 1990, **31**: 509–513.
38. Taylor, T.V. & Holt, S. Antireflux surgery. A time for reappraisal. *Br Med J* 1990, **300**: 1603–1604.
39. Timoney, A.G., Kelly, J.M. & Welfare, M.R. The Angelchik antireflux device: a 5-year experience. *Ann R Coll Surg Engl* 1990, **72**: 185–187.
40. Mughal, M.M., Bancewicz, J. & Marples, M. Oesophageal manometry and pH recording does not predict the bad results of Nissen fundoplication. *Br J Surg* 1990, **77**: 43–45.
41. Muller, J.M., Erasmi, H., Stelzner, M. *et al.* Surgical therapy of oesophageal carcinoma. *Br J Surg* 1990, **77**: 845–857.

42. Mannell, A. & Becker, P.J. Evaluation of the results of oesophagectomy for oesophageal cancer. *Br J Surg* 1991, **78**: 36–40.
 43. Earlam, R.J. & Johnson, L. 101 Oesophageal cancers: a surgeon uses radiotherapy. *Ann R Coll Surg Engl* 1990, **72**: 32–40.
 44. Khoury, G.A. Squamous cell carcinoma of the oesophagus: 10 years on. *Ann R Coll Surg Engl* 1991, **73**: 4–7.
 45. Earlam, R. An MRC prospective randomised trial of radiotherapy versus surgery for operable squamous cell carcinoma of the oesophagus. *Ann R Coll Surg Engl* 1991, **73**: 8–12.
 46. Oliver, S.E., Robertson, C.S., Logan, R.F.A. *et al.* What does radiotherapy add to survival over endoscopic intubation alone in inoperable squamous cell oesophageal cancer? *Gut* 1990, **31**: 750–752.
 47. Barr, H., Krasner, N., Raouf, A. *et al.* Prospective randomised trial of laser therapy only and laser therapy followed by endoscopic intubation for the palliation of malignant dysphagia. *Gut* 1990, **31**: 252–258.
 48. Rauws, E.A.J. & Tytgat, G.N.J. Cure of duodenal ulcer associated with eradication of *Helicobacter pylori*. *Lancet* 1990, **335**: 1233–1235.
 49. Cooperative Study Group. Double blind comparative study of omeprazole and ranitidine in patients with duodenal or gastric ulcer: a multicentre trial. *Gut* 1990, **31**: 653–656.
 50. Johnston, G.W., Spencer, E.F.A., Wilkinson, A.J. *et al.* Proximal gastric vagotomy: follow-up at 10–20 years. *Br J Surg* 1991, **78**: 20–23.
 51. Macintyre, I.M.C., Millar, A., Smith, A.N. *et al.* Highly selective vagotomy 5–15 years on. *Br J Surg* 1990, **77**: 65–69.
 52. Koruth, N.M., Dua, K.S., Brunt, P.W. *et al.* Comparison of highly selective vagotomy with truncal vagotomy and pyloroplasty: results at 8–15 years. *Br J Surg* 1990, **77**: 70–72.
 53. Taylor, T.V., Lythgoe, J.P., McFarland, J.B. *et al.* Anterior lesser curve seromyotomy and posterior truncal vagotomy versus truncal vagotomy and pyloroplasty in the treatment of chronic duodenal ulcer. *Br J Surg* 1990, **77**: 1007–1009.
 54. Holman, R.A.E., Davis, M., Gough, K.R. *et al.* Value of a centralised approach in the management of haematemesis and melaena: experience in a district general hospital. *Gut* 1990, **31**: 504–508.
 55. Sanderson, J.D., Taylor, R.F.H., Pugh, S. *et al.* Specialized gastrointestinal units for the management of upper gastrointestinal haemorrhage. *Postgrad Med J* 1990, **66**: 654–656.
 56. Wheatley, K.E., Snymann, J.H., Brearley, S. *et al.* Mortality in patients with bleeding peptic ulcer when those aged 60 or over are operated on early. *Br Med J* 1990, **301**: 272.
 57. Branicki, F.J., Coleman, S.Y., Fok, P.J. *et al.* Bleeding peptic ulcer: a prospective evaluation of risk factors for rebleeding and mortality. *World J Surg* 1990, **14**: 262–270.
 58. Teenan, R.P. & Murray, W.R. Late outcome of undersewing alone for gastric ulcer haemorrhage. *Br J Surg* 1990, **77**: 811–812.
 59. Hunt, P.S. & McIntyre, R.L.E. Choice of emergency operative procedure for bleeding duodenal ulcer. *Br J Surg* 1990, **77**: 1004–1006.
 60. Lin, H.J., Lee, F.Y., Kang, W.M. *et al.* Heat probe thermocoagulation and pure alcohol injection in massive peptic ulcer haemorrhage: a prospective, randomised controlled trial. *Gut* 1990, **31**: 753–757.
 61. de Dombal, F.T., Price, A.B., Thompson, H. *et al.* The British Society of Gastroenterology early gastric cancer/dysplasia survey: an interim report. *Gut* 1990, **31**: 115–120.
 62. Hallissey, M.T., Allum, W.H., Jewkes, A.J. *et al.* Early detection of gastric cancer. *Br Med J* 1990, **301**: 513–515.
 63. Bozzetti, F., Regalia, E., Bonfanti, G. *et al.* Early and late results of extended surgery for cancer of the stomach. *Br J Cancer* 1990, **77**: 53–56.
 64. Griffin, S.M., Chung, S.C.S., Woods, S.D.S. *et al.* Adenocarcinoma of the cardia: treatment by thoracoabdominal R3 radical gastrectomy. *Br J Surg* 1990, **77**: 937–939.
 65. Wilson, N.V., Geall, A., Kittermaster, R. *et al.* Thoracoabdominal total gastrectomy in the management of adenocarcinoma of the cardia. Is it worth it? *Ann R Coll Surg Engl* 1990, **72**: 329–334.
 66. Gill, P.G., Jamieson, G.G., Denham, J. *et al.* Treatment of adenocarcinoma of the cardia with synchronous chemotherapy and radiotherapy. *Br J Surg* 1990, **77**: 1020–1023.
 67. Buhl, K., Schlag, P. & Herfarth, C. Quality of life and functional results following different types of resection for gastric carcinoma. *Eur J Surg Oncol* 1990, **16**: 404–409.
 68. Cuschieri, A. Jejunal pouch reconstruction after total gastrectomy for cancer: experience in 29 patients. *Br J Surg* 1990, **77**: 421–424.
- Biliary disease**
69. Cheslyn-Curtis, S. & Russel, R.C.G. New trends in gallstone management. *Br J Surg* 1991, **78**: 143–149.
 70. Wastell, C. Laparoscopic cholecystectomy. *Br Med J* 1991, **302**: 303–304.
 71. Patterson-Brown, S., Garden, O.J. & Carter, D.C. Laparoscopic cholecystectomy. *Br J Surg* 1991, **78**: 131–132.
 72. Neugebauer, E., Troidl, H., Spangenberg, W. *et al.* Conventional versus laparoscopic cholecystectomy and the randomized controlled trial. *Br J Surg* 1991, **78**: 150–154.
 73. Nathanson, L.K., Shimi, S. & Cuschieri, A. Laparoscopic cholecystectomy: the Dundee technique. *Br J Surg* 1991, **78**: 155–159.
 74. Grace, P.A., Quereshi, A., Coleman, J. *et al.* Reduced postoperative hospitalization after laparoscopic cholecystectomy. *Br J Surg* 1991, **78**: 160–162.
 75. Darzi, A., El-Sayed, E., O'Morain, C. *et al.* Piezoelectric lithotripsy for gallstones: analysis of results in patients with extended selection. *Br J Surg* 1991, **78**: 163–166.
 76. Darzi, A., Reid, I., Kay, E. *et al.* Piezoelectric lithotripsy and soft tissue injury. Safety limits in the experimental and clinical setting. *Gut* 1990, **31**: 1110–1114.
 77. Schachler, R., Bird, N.C., Sauerbruch, T. *et al.* Extracorporeal shock wave lithotripsy of gallstones: an *in vitro* comparison between an electrohydraulic and a piezoceramic device. *Gut* 1991, **32**: 312–315.
 78. Chiverton, S.G., Inglis, J.A., Hudd, C. *et al.* Percutaneous cholecystolithotomy: the first 60 patients. *Br Med J* 1990, **300**: 1310–1312.
 79. Neoptolemos, J.P., Hall, C., O'Connor, H.J. *et al.* Methyl-tert-butyl-ether for treating bile duct stones: the British experience. *Br J Surg* 1990, **77**: 32–35.
 80. Motson, R.W. & Wetter, L.A. Operative choledochoscopy: common bile duct exploration is incomplete without it. *Br J Surg* 1990, **77**: 975–982.
 81. Pernthaler, H., Sandbichler, P., Schmid, Th. *et al.* Operative cholangiography in elective cholecystectomy. *Br J Surg* 1990, **77**: 399–400.
 82. Horton, R.C., Lauri, A. & Dooley, J.S. Endoscopic removal of common duct stones: current indications and controversies. *Postgrad Med J* 1991, **67**: 107–111.
 83. Worthley, C.S., Watts, J.M. & Toouli, J. Common duct exploration or endoscopic sphincterotomy for choledocholithiasis? *Aust N Z J Surg* 1989, **59**: 209–215.
- Pancreatic disease**
84. Clavien, P.-A., Burgan, S. & Moossa, A.R. Serum enzymes and other laboratory tests in acute pancreatitis. *Br J Surg* 1989, **76**: 1234–1243.
 85. Gudgeon, A.M., Heath, D.I., Hurley, P. *et al.* Trypsinogen activation peptides assay in the early prediction of severity of acute pancreatitis. *Lancet* 1990, **335**: 4–6.

86. Bassi, C., Vesentini, S., Nifosi, F. *et al.* Pancreatic abscess and other pus-harbouring collections related to pancreatitis: a review of 108 cases. *World J Surg* 1990, **14**: 505–512.
 87. Poston, G.J. & Williamson, R.C.N. Surgical management of acute pancreatitis. *Br J Surg* 1990, **77**: 5–12.
 88. D'Egidio, A. & Schein, M. Surgical strategies in the treatment of pancreatic necrosis and infection. *Br J Surg* 1991, **78**: 133–137.
 89. Villazon, A., Villazon, O., Terrazas, F. *et al.* Retroperitoneal drainage in the management of the septic phase of severe acute pancreatitis. *World J Surg* 1991, **15**: 103–108.
 90. Robin, A.P., Campbell, R., Palani, C.K. *et al.* Total parenteral nutrition during acute pancreatitis: clinical experience with 156 patients. *World J Surg* 1990, **14**: 572–579.
 91. Ihse, I., Borch, K. & Larsson, J. Chronic pancreatitis: results of operations for relief of pain. *World J Surg* 1990, **14**: 53–58.
 92. Imrie, C.W. Management of recurrent pain following previous surgery for chronic pancreatitis. *World J Surg* 1990, **14**: 88–93.
 93. Morel, P., Mathey, P., Corboud, H. *et al.* Pylorus preserving duodenopancreatectomy: long-term complications and comparison with the Whipple procedure. *World J Surg* 1990, **14**: 642–647.
 94. Hall, R.I., Rhodes, M., Isabel-Martinez, L. *et al.* Pancreatic exocrine function after a sutureless pancreatico-jejunostomy following pancreatico-duodenectomy. *Br J Surg* 1990, **77**: 83–85.
 95. Hatfield, A.R.W. Palliation of malignant obstructive jaundice – surgery or stent? *Gut* 1990, **31**: 1339–1340.
 96. Holbrook, A.G., Chester, J.F. & Britton, D.C. Surgical palliation for pancreatic cancer: will biliary bypass alone suffice? *J R Soc Med* 1990, **83**: 12–14.
 97. Zamboni, W.A., Fisher, K.S. & Ross, D.S. Surgical palliation for pancreatic carcinoma. *Postgrad Med J* 1991, **67**: 362–365.
 98. Bickerstaff, K.I., Berry, A.R., Chapman, R.W. *et al.* Endoscopic sphincterotomy for the palliation of ampullary carcinoma. *Br J Surg* 1990, **77**: 160–162.
 99. Kahn, P.J., Skornick, Y., Inbar, M. *et al.* Surgical palliation combined with synchronous therapy in pancreatic carcinoma. *Eur J Surg Oncol* 1990, **16**: 7–11.
 100. Bakkevoeld, K.E., Pettersen, A., Arnesjo, B. *et al.* Tamoxifen therapy in unresectable adenocarcinoma of the pancreas and the papilla of Vater. *Br J Surg* 1990, **77**: 725–730.
- Liver**
101. Karran, S. Progress in the assessment of liver blood flow in health and disease. *J R Coll Surg Edinb* 1990, **35**: 207–217.
 102. Hayes, P.C., Davis, J.M., Lewis, J.A. *et al.* Meta-analysis of value of propranolol in prevention of variceal haemorrhage. *Lancet* 1990, **336**: 153–156.
 103. Kahn, D., Terblanche, J., Kitano, S. *et al.* Injection sclerotherapy in adult patients with extrahepatic portal venous obstruction. *Br J Surg* 1990, **74**: 600–602.
 104. Mathur, S.K., Dalvi, A.N., Someswar, V. *et al.* Endoscopic radiological appraisal of gastric varices. *Br J Surg* 1990, **77**: 432–435.
 105. Greig, J.D., Garden, O.J., Anderson, J.R. *et al.* Management of gastric variceal haemorrhage. *Br J Surg* 1990, **77**: 297–299.
 106. Bird, G.L.A., O'Grady, J.G., Harvey, F.A.H. *et al.* Liver transplantation in patients with alcoholic cirrhosis: selection criteria and rates of survival and relapse. *Br Med J* 1990, **301**: 15–17.
 107. Neuberger, J.M., Gunson, B.K., Buckels, J.A.C. *et al.* Referral of patients with primary biliary cirrhosis for liver transplantation. *Gut* 1990, **31**: 1069–1072.
 108. Clarke, M.P., Kane, R.A., Steele, G Jr. *et al.* Prospective comparison of preoperative imaging and intraoperative ultrasonography in the detection of liver tumours. *Surgery* 1989, **106**: 849–855.
 109. Editorial. Lipiodol computed tomography for small hepatocellular carcinomas. *Lancet* 1991, **337**: 333–334.
 110. Belghiti, J. Resection of hepatocellular carcinoma complicating cirrhosis. *Br J Surg* 1991, **78**: 257–258.
 111. Ismail, T., Angrisani, L., Gunson, B.K. *et al.* Primary hepatic malignancy: the role of liver transplantation. *Br J Surg* 1990, **77**: 983–987.
 112. Ringe, B., Pichlmayr, R., Wittekind, C. *et al.* Surgical treatment of hepatocellular carcinoma: experience with liver resection and transplantation in 198 patients. *World J Surg* 1991, **15**: 270–285.
 113. Otto, G., Heuschen, U.A., Hofmann, W.J. *et al.* Primary hepatic malignancies: resection or liver transplantation? *Eur J Surg Oncol* 1990, **16**: 346–351.
 114. Al-Hadeedi, S., Choi, T.K. & Wong, J. Extended hepatectomy for hepatocellular carcinoma. *Br J Surg* 1990, **77**: 1247–1250.
 115. Paquet, K.-J., Koussouris, P., Mercado, M.A. *et al.* Limited hepatic resection for selected cirrhotic patients with hepatocellular or cholangiocellular carcinoma: a prospective study. *Br J Surg* 1991, **78**: 459–462.
 116. Hasegawa, H., Makuuchi, M., Yamazaki, S. *et al.* Central bisegmentectomy of the liver: experience in 16 patients. *World J Surg* 1989, **13**: 786–790.
 117. Dewar, G.A., Griffin, S.M., Ku, K.W. *et al.* Management of bleeding liver tumours in Hong Kong. *Br J Surg* 1991, **78**: 463–466.
 118. Karanjia, N.D., Rees, M., Schache, D. *et al.* Hepatic resection for colorectal secondaries. *Br J Surg* 1990, **77**: 27–29.
 119. Scheele, J., Stangl, R. & Altendorf-Hofmann, A. Hepatic metastases from colorectal carcinoma: impact of surgical resection on the natural history. *Br J Surg* 1990, **77**: 1241–1246.
 120. Hunt, T.M., Carty, N. & Johnson, C.D. Resection of liver metastases from a colorectal carcinoma does not benefit the patients. *Ann R Coll Surg Engl* 1990, **72**: 199–205.
 121. Schlag, P., Hohenberger, P. & Herfarth, Ch. Resection of liver metastases in colorectal cancer: competitive analysis of treatment results in synchronous versus metachronous metastases. *Eur J Surg Oncol* 1990, **16**: 360–365.
 122. Vogt, P., Raab, R., Ringe, B. *et al.* Resection of synchronous liver metastases from colorectal cancer. *World J Surg* 1991, **15**: 62–67.
 123. Hunt, T.M., Flowerdew, A.D.S., Birch, S.J. *et al.* Prospective randomized controlled trial of hepatic arterial embolization of infusion chemotherapy with 5-fluorouracil and degradable starch microspheres for colorectal liver metastases. *Br J Surg* 1990, **77**: 779–782.
 124. Goldberg, J.A., Kerr, D.J., Wilmott, N. *et al.* Regional chemotherapy for colorectal liver metastases: a phase II evaluation of targeted hepatic arterial 5-fluorouracil for colorectal liver metastases. *Br J Surg* 1990, **77**: 1238–1240.
 125. Andersson, R. & Bengmark, S. Conservative treatment of liver trauma. *World J Surg* 1990, **14**: 483–486.
 126. Hollands, M.J. & Little, J.M. The role of hepatic resection in the management of blunt liver trauma. *World J Surg* 1990, **14**: 478–482.
 127. Ciresi, K.F. & Lim, R.C. Jr. Hepatic vein and retrohepatic vena caval injury. *World J Surg* 1990, **14**: 472–477.
- Colorectal disease**
128. Rew, D.A., Wilson, G.D., Taylor, I. *et al.* Proliferation characteristics of human colorectal carcinomas measured *in vivo*. *Br J Surg* 1991, **78**: 60–66.
 129. Rowley, S., Newbold, K.M., Gearty, J. *et al.* Comparison of deoxyribonucleic acid ploidy and nuclear expressed p62 c-myc oncogene in the prognosis of colorectal cancer. *World J Surg* 1990, **14**: 545–551.
 130. Scott, N., Sagar, P., Stewart, J. *et al.* p53 in colorectal cancer: clinicopathological correlation and prognostic significance. *Br J Cancer* 1991, **63**: 317–319.

131. Maestro, R., Viel, A. & Boiocchi, M. Correlation between chromosome 5q deletions and different mechanisms of c-myc overexpression in human colorectal cancer. *Br J Cancer* 1991, **63**: 185–186.
132. Steele, R.J.C., Kelly, P., Ellul, B. *et al.* Epidermal growth factor receptor expression in colorectal cancer. *Br J Surg* 1990, **77**: 1352–1354.
133. Steele, R.J.C., Kelly, P., Ellul, B. *et al.* Immunohistochemical detection of epidermal growth factor receptors on human colonic carcinomas. *Br J Cancer* 1990, **61**: 325–326.
134. Sarode, R., Marwaha, N. & Gupta, N.M. Procoagulant activity of human tumours: existence of Xa and thrombin-like activities. *Br J Cancer* 1990, **61**: 29–31.
135. Gibson, P.R., van de Pol, E. & Doe, W.F. Cell associated urokinase activity and colonic epithelial cells in health and disease. *Gut* 1991, **32**: 191–195.
136. Carty, N., Taylor, I., Roach, O.S. *et al.* Urinary tissue factor activity in colorectal disease. *Br J Surg* 1990, **77**: 1091–1094.
137. Bingham, S.A. Diet and large bowel cancer. *J R Soc Med* 1990, **83**: 420–422.
138. Neoptolemos, J.P., Husband, D., Imray, C. *et al.* Arachidonic acid and docosahexaenoic acid are increased in human colorectal cancer. *Gut* 1991, **32**: 278–281.
139. Nicholson, M.L., Neoptolemos, J.P., Clayton, H.A. *et al.* Increased cell membrane arachidonic acid in experimental colorectal tumours. *Gut* 1991, **32**: 413–418.
140. Mullan, F.J., Wilson, H.K., Majury, C.W. *et al.* Bile acids and the increased risk of colorectal tumours after truncal vagotomy. *Br J Surg* 1990, **77**: 1085–1090.
141. Spiegelman, A.D., Owen, R.W., Hill, M.J. *et al.* Biliary bile acid profiles in familial adenomatous polyposis. *Br J Surg* 1991, **78**: 321–325.
142. Collins, J.E., Taylor, I. & Garrard, D.R. A study of desmosomes in colorectal cancer. *Br J Cancer* 1990, **62**: 796–805.
143. Kewenter, J., Engaras, B., Haglind, E. *et al.* Value of retesting subjects with a positive Hemoccult in screening for colorectal cancer. *Br J Surg* 1990, **77**: 1349–1351.
144. Thomas, W.M., Pye, G., Hardcastle, J.D. *et al.* Faecal occult blood screening for colorectal neoplasia: a randomized trial of three days or six days of tests. *Br J Surg* 1990, **77**: 277–279.
145. Stelling, H.P., Maimon, H.N., Smith, R.A. *et al.* A comparative study of fecal occult blood tests for early detection of gastrointestinal pathology. *Arch Intern Med* 1990, **150**: 1001–1005.
146. Fujita, M., Sugiyama, R., Kumanishi, Y. *et al.* Evaluation of effectiveness of mass screening for colorectal cancer. *World J Surg* 1990, **14**: 648–653.
147. Chamberlain, J. Is screening for colorectal cancer worthwhile? *Br J Cancer* 1990, **62**: 1–3.
148. Sampliner, R.E. Limitations of fecal occult blood testing. *Arch Intern Med* 1990, **150**: 945–946.
149. Juhl, G., Larson, G.M., Mullins, R. *et al.* Six-year results of annual colonoscopy after resection of colorectal cancer. *World J Surg* 1990, **14**: 255–261.
150. Barilari, P., Ramacciato, G., de Angelis, R. *et al.* The role of CEA, TPA and CA 19-9 in the early detection of recurrent colorectal cancer. *Int J Colorect Dis* 1989, **4**: 230–233.
151. Granowska, M., Mather, S.J., Britton, K.E. *et al.* ^{99m}Tc radioimmunoscinigraphy of colorectal cancer. *Br J Cancer* 1990, **62** (Suppl. X): 30–33.
152. Hertel, A., Baum, R.P., Lorenz, M. *et al.* Immunoscintigraphy using a technetium-99m labelled monoclonal anti-CEA antibody in the follow-up of colorectal cancer and other tumours producing CEA. *Br J Cancer* 1990, **62** (Suppl. X): 34–36.
153. Holting, Th., Schlag, P. & Georgi, P. Current status of immunoscintigraphy in colorectal cancer – results of 5 years' experiences. *Eur J Surg Oncol* 1990, **16**: 312–318.
154. Benyon, J., Mortensen, N.S. McC, Foy, D.M.A. *et al.* Preoperative assessment of mesorectal lymph node involvement in rectal cancer. *Br J Surg* 1989, **76**: 276–279.
155. Glaser, F., Schlag, P. & Herfarth, Ch. Endorectal ultrasonography for the assessment of invasion of rectal tumours and lymph node involvement. *Br J Surg* 1990, **77**: 883–887.
156. McArdle, C.S., Hole, D., Hansell, D. *et al.* Prospective study of colorectal cancer in the West of Scotland: 10-year follow-up. *Br J Surg* 1990, **77**: 280–282.
157. Karanjia, N.D., Schache, D.J., North, W.R.S. *et al.* 'Close shave' in anterior resection. *Br J Surg* 1990, **77**: 510–512.
158. Cawthorn, S.J., Parums, D.V., Gibbs, N.M. *et al.* Extent of mesorectal spread and involvement of lateral resection margin as prognostic factors after surgery for rectal cancer. *Lancet* 1990, **335**: 1055–1059.
159. Editorial. Breaching the mesorectum. *Lancet* 1990, **335**: 1067–1068.
160. Geoghegan, J.G. & Rosenberg, I.L. Experience with early anastomosis after the Hartmann procedure. *Ann R Coll Surg Engl* 1991, **73**: 80–82.
161. Novell, J.R. & Lewis, A.A.M. Peroperative observation of marginal artery bleeding: a predictor of anastomotic leakage. *Br J Surg* 1990, **77**: 137–138.
162. Berry, A.R., Souter, R.G., Campbell, W.B. *et al.* Endoscopic transanal resection of rectal tumours – a preliminary report of its use. *Br J Surg* 1990, **77**: 134–137.
163. Doudi, S., Wilson, N.M. & Heddle, R.M. Primary restorative colectomy in malignant left-sided large bowel obstruction. *Ann R Coll Surg* 1990, **72**: 393–395.
164. Stephenson, B.M., Shandall, A.A., Farouk, R. *et al.* Malignant left-sided large bowel obstruction managed by sub/total colectomy. *Br J Surg* 1990, **77**: 1098–1102.
165. Runkel, N.S., Schlag, P., Schwarz, V. *et al.* Outcome after emergency surgery for cancer of the large intestine. *Br J Surg* 1991, **78**: 183–188.
166. Loizou, L.A., Griff, D., Boulos, P.B. *et al.* Endoscopic Nd:YAG laser treatment of rectosigmoid cancer. *Gut* 1990, **31**: 812–816.
167. Von Dittfurth, B., Buhl, K. & Friedl, P. Palliative endoscopic therapy for rectal cancer with neodymium:YAG laser. *Eur J Surg Oncol* 1990, **16**: 376–379.
168. Barr, H., Krasner, N., Boulos, P.B. *et al.* Photodynamic therapy for colorectal cancer: a quantitative pilot study. *Br J Surg* 1990, **77**: 93–96.
169. Taylor, I. & Northover, J.M.A. Adjuvant therapy in colorectal cancer: need for a mega-trial. *Br J Surg* 1990, **77**: 841–842.
170. Laurie, J.A., Moertel, C.G., Fleming, T.R. *et al.* Surgical adjuvant therapy of large-bowel carcinoma: an evaluation of levamisole and the combination of levamisole and fluorouracil. *J Clin Oncol* 1989, **7**: 1447–1456.
171. Moertel, C.G., Fleming, T.R., MacDonald, J.S. *et al.* Levamisole and fluorouracil for adjuvant therapy of resected colon carcinoma. *N Engl J Med* 1990, **322**: 352–358.
172. Hafstrom, L., Domellof, L., Rudenstam, C.M. *et al.* Adjuvant chemotherapy with 5-fluorouracil, vincristine and CCNU for patients with Dukes' C colorectal cancer. *Br J Surg* 1990, **77**: 1345–1348.
173. Jarvinen, H.J. & Luukkonen, P. Comparison of restorative proctocolectomy with and without covering ileostomy in ulcerative colitis. *Br J Surg* 1991, **78**: 199–201.
174. Madden, M.V., Farthing, M.J.G. & Nicholls, R.J. Inflammation in ileal reservoirs: 'pouchitis'. *Gut* 1990, **31**: 247–249.
175. Levitt, M.D. & Lewis, A.A.M. Determinants of ileoanal pouch function. *Gut* 1991, **32**: 126–127.
176. Sugita, A., Sachar, D.B., Bodian, C. *et al.* Colorectal cancer in ulcerative colitis. Influence of anatomical extent and age at onset on colitis-cancer interval. *Gut* 1991, **32**: 167–169.
177. Lennard-Jones, J.E., Melville, D.M. & Morson, B.C. *et al.* Precancer and cancer in extensive ulcerative colitis: findings among 401 patients over 22 years. *Gut* 1990, **31**: 800–806.
178. Gyde, S. Screening for colorectal cancer in ulcerative colitis: dubious benefits and high costs. *Gut* 1990, **31**: 1089–1092.

179. Akbom, A., Helmich, C., Zack, M. *et al.* Increased risk of large bowel cancer in Crohn's disease with colonic involvement. *Lancet* 1990, **336**: 357–359.
180. Williams, J.G., Wong, W.D., Rothenberger, D.A. *et al.* Recurrence of Crohn's disease after resection. *Br J Surg* 1991, **78**: 10–19.
181. Weterman, I.T., Biemond, I. & Pena, A.S. Mortality and causes of death in Crohn's disease. Review of 50 years' experience in Leiden University Hospital. *Gut* 1990, **31**: 1387–1390.
182. Paterson-Brown, S. & Vipond, M.V. Modern aids to clinical decision-making in the acute abdomen. *Br J Surg* 1990, **77**: 13–18.
183. Ooms, H.W.A., Koumans, R.K.J., Ho Kang You, P.J. *et al.* Ultrasonography in the diagnosis of acute appendicitis. *Br J Surg* 1991, **78**: 315–318.
184. Vipond, M.N., Paterson-Brown, S., Tyrrell, M.R. *et al.* Evaluation of fine catheter aspiration cytology of the peritoneum as an adjunct to decision making in the acute abdomen. *Br J Surg* 1990, **77**: 86–87.
185. Raheja, S.K.J., McDonald, P.J. & Taylor, I. Non-specific abdominal pain – an expensive mystery. *J R Soc Med* 1990, **83**: 10–11.
186. Jones, P.F. Practicalities in the management of the acute abdomen. *Br J Surg* 1990, **77**: 365–367.

The acute abdomen