Elevated serum parathyroid hormone related protein and 1,25-dihydroxycholecalciferol in hypercalcaemia associated with adult T-cell leukaemia-lymphoma

S.R.D. Johnston and P.J. Hammond

Department of Medicine, Royal Postgraduate Medical School, Hammersmith Hospital, Du Cane Road, London W12 0HS, UK

Summary: Hypercalcaemia occurs in up to 80% of patients with adult T-cell leukaemia-lymphoma (ATLL) associated with human T-cell leukaemia virus-1 infection. Elevated serum levels of 1,25-dihydroxycholecalciferol, implicated in the pathogenesis of hypercalcaemia in lymphoma, and of parathyroid hormone-related protein (PTHrP), which is associated with hypercalcaemia of several solid malignancies, were demonstrated in a patient with ATLL hypercalcaemia. Treatment with bisphosphonates reduced the serum calcium but had no significant effect on the serum PTHrP levels. This case supports recent *in vitro* evidence for enhanced PTHrP expression in ATLL tumour cells and suggests that more than one tumour cell product may be involved in the pathogenesis of ATLL hypercalcaemia.

Introduction

Hypercalcaemia is a common finding in adult T-cell leukaemia lymphoma (ATLL).¹ A variety of tumour cell products have been implicated in the development of ATLL hypercalcaemia. Increased circulating concentrations of 1,25-dihydroxycholecalciferol (DHCC) have been associated with hypercalcaemia occurring in both Hodgkin's disease² and non-Hodgkin's lymphoma.³ Ectopic production of DHCC has been shown in human T-cell leukaemia virus-1 (HTLV-1) transformed lymphocytes⁴ suggesting that it also may become involved in the pathogenesis of hypercalcaemia in HTLV-1 associated ATLL.

The demonstration of parathyroid hormone-related protein (PTHrP) messenger RNA in various solid tumours complicated by hypercalcaemia implies that this peptide may play a major role in humoral hypercalcaemia of malignancy.⁵ Circulating serum PTHrP has been detected by radioimmunoassay in some patients with solid tumours associated with malignant hypercalcaemia including lung, breast and head and neck cancer.⁶ Evidence for the role of PTHrP in haematological malignancies is less clear cut. Although there has been increasing *in vitro* evidence for PTHrP expression in ATLL, there has

been little evidence for elevated serum levels in these patients. We report a case of hypercalcaemia in HTLV-1-associated ATLL in whom both the serum PTHrP and DHCC were found to be elevated and discuss the evidence implicating their role in the pathogenesis of ATLL hypercalcaemia.

Case report

A 52 year old woman from Guyana was admitted to hospital with bilateral facial nerve weakness, drowsiness and dehydration. Investigations revealed hypercalcaemia, with a serum calcium of 4.68 mmol/l and serum phosphate of 1.73 mmol/l. Further investigations showed a leucoerythroblastic blood film, with a peripheral lymphocyte count of 38,000/mm³. Serology for HTLV-1 was positive, and a bone marrow examination confirmed adult T-cell lymphoma with CD4+/CD8- T cells expressing the CD25 (TAC) antigen. In addition, marked osteoclast activity with bone resorption was seen.

The plasma urea and electrolytes were normal and the serum angiotensin converting enzyme was 32 IU/l (normal range < 52 IU/l). Serum parathyroid hormone (PTH) was undetectable. The 25-hydroxycholecalciferol level was suppressed at 5.40 ng/ml (normal range 7–50 ng/ml), with the 1,25-dihydroxycholecalciferol (DHCC) level elevated at 78 pg/ml (normal range 20–65 pg/ml). The serum PTHrP (1–86) measured by radioimmunoassay was significantly elevated at 0.9 pmol/l (nor-

Correspondence: S.R.D. Johnston, M.R.C.P., Department of Academic Biochemistry, Royal Marsden Hospital, Fulham Road, London SW3 6JJ, UK.

Accepted: 24 February 1992

mal range <0.23 pmol/l). After rehydration with normal saline disodium pamidronate (60 mg i.v.) was given. After 48 hours the calcium was lowered to 2.07 mmol/l with a serum phosphate of 0.41 mmol/l. Repeat measurement of the serum PTHrP showed persistent elevation at 0.51 pmol/l.

Although her general condition improved with treatment of the hypercalcaemia, she subsequently developed hyperinfection with *Strongyloides stercoralis*. She died 30 days after admission.

Discussion

A number of mediators of the humoral hypercalcaemia of malignancy (HMM) have been proposed including parathyroid hormone-related protein (PTHrP), various cytokines and 1,25-dihydroxycholecalciferol (DHCC).⁷ Cytokines appear to be more important in the development of local osteolytic hypercalcaemia resulting from metastatic disease. Cells of the monocytic-macrophage line are capable of converting 25-hydroxycholecalciferol to DHCC, and elevated levels of DHCC have been found in association with lymphomas,³ with synthesis of DHCC demonstrated in tumour cells.⁸ Since its isolation in 1987 it has become apparent that PTHrP is a common mediator of the HMM syndrome, with elevated serum levels found in 48–71% of solid tumours associated with HMM.⁶ In breast cancer, 92% of hypercalcaemic patients with bone metastases have elevated serum levels of PTHrP compared with 36% of normocalcaemic patients with bone metastases and 9% of normocalcaemic patients without bone metastases.⁹ Elevated serum levels of PTHrP are less common in HMM associated with haematological malignancies occurring in 30% of patients with myeloma⁶ and in four out of five patients with lymphoma in Kao's study.¹⁰

The incidence of hypercalcaemia in patients with HTLV-1-associated adult T-cell leukaemia-lymphoma is about 80% and, although the aetiology has not been fully elucidated, DHCC levels are often found to be suppressed.¹¹ There has been increasing

evidence implicating a role for PTHrP. Recently, immunocytochemical demonstration of PTHrP protein has been reported in lymphoblastic tissue from patients with ATLL.¹² Previous *in vitro* studies have demonstrated expression of PTHrP messenger RNA in both HTLV-1-infected T-cell lines,¹³ and in tumour cells from ATLL patients with hypercalcaemia.¹⁴ Enhanced expression of the PTHrP gene has been shown in HTLV-1-infected T-cells both in asymptomatic HTLV-1 carriers and in ATLL patients, but not in normal T-cells.¹⁵ Furthermore the HTLV-1 'tax' gene has been shown to transactivate the PTHrP gene promoter in co-transfection assays, and it is suggested that further activation of the gene during malignant transformation to ATLL may occur, thus accounting for the high incidence of hypercalcaemia in ATLL patients.

This is the first report of high circulating serum levels of PTHrP in a patient with HTLV-1-positive human adult T-cell leukaemia-lymphoma. Previous reports of elevated serum levels have been restricted to solid tumours associated with hypercalcaemia.⁶ The response to bisphosphonates reflected the probable role of PTHrP in causing the hypercalcaemia. It remains unclear whether other tumour products may also be involved in the development of hypercalcaemia in ATLL. The elevated 1,25-dihydroxycholecalciferol probably contributed to this patient's hypercalcaemia, indicating that the aetiology of HMM may be multifactorial. However, the *in vitro* evidence for PTHrP expression in these patients along with this report of elevated serum levels strongly implicate PTHrP as one mediator of the hypercalcaemia associated with ATLL.

Acknowledgements

We are grateful to Dr W.A. Ratcliffe, Department of Medicine, Queen Elizabeth Hospital, Birmingham, for performing the PTHrP assay, to Dr M. Hewison, Department of Medicine, Middlesex Hospital, London, for performing the 1,25-dihydroxycholecalciferol assay, and to Dr J. Cohen, Reader in Infectious Diseases, Hammsmith Hospital, who looked after the patient.

References

1. Catovsky, D., Greaves, M.F., Rose, M. *et al.* Adult T-cell leukaemia-lymphoma in blacks from the West Indies. *Lancet* 1982, **i**: 639–643.
2. Karmali, R., Barker, S., Hewison, M. *et al.* Intermittent hypercalcaemia and vitamin D sensitivity in Hodgkin's disease. *Postgrad Med J* 1990, **66**: 757–760.
3. Breslau, N.A., McGuire, J.L., Zerwekh, J.E. *et al.* Hypercalcaemia associated with increased serum calcitriol levels in three patients with lymphoma. *Ann Intern Med* 1984, **100**: 1–7.
4. Fetchick, D.A., Bertolini, D.R., Sarin, P.S. *et al.* Production of 1,25-dihydroxyvitamin D₃ by human T-cell lymphotropic virus-1 transformed lymphocytes. *J Clin Invest* 1986, **78**: 592–596.
5. Heath, D.A., Senior, P.V., Varley, J.M. & Beck, F. Parathyroid-hormone-related protein in tumours associated with hypercalcaemia. *Lancet* 1990, **335**: 66–69.
6. Budayer, A.A., Nissenson, R.A., Klein, R.F. *et al.* Increased serum levels of a parathyroid hormone like protein in malignancy associated hypercalcaemia. *Ann Intern Med* 1989, **111**: 807–812.
7. Gutierrez, G.E., Poser, J.W., Katz, M.S. *et al.* Mechanisms of hypercalcaemia of malignancy. *Bailliere's Clin Endocrinol Metab* 1990, **4**: 119–138.
8. Mudde, A.A., Van den Berg, H., Boshuis, P.G., Breedveld, F.C. *et al.* Ectopic production of 1,25-dihydroxyvitamin D by B-cell lymphoma as a cause of hypercalcaemia. *Cancer* 1987, **59**: 1543–1546.

9. Bundred, N.J., Ratcliffe, W.A., Walker, R.A. *et al.* Parathyroid hormone related protein and hypercalcaemia in breast cancer. *Br Med J* 1991, **303**: 1506–1509.
10. Kao, P.C., Klee, G.C., Taylor, R.L. & Heath, H. Parathyroid hormone-related peptide in plasma of patients with hypercalcaemia and malignant lesions. *Mayo Clin Proc* 1990, **65**: 1399–1407.
11. Dodd, R.C., Winkler, C.F., Williams, M.E. *et al.* Calitriol levels in hypercalcaemic patients with adult T-cell lymphoma. *Arch Intern Med* 1986, **146**: 1971–1972.
12. Moseley, J.M., Danks, J.A., Grill, V., Lister, T.A. & Horton, M.A. Immunocytochemical demonstration of PTHrP protein in neoplastic tissue of HTLV-1 positive human adult T cell leukaemia-lymphoma—implications for the mechanism of hypercalcaemia. *Br J Cancer* 1991, **64**: 745–748.
13. Motokura, T., Fukumoto, S., Takahashi, S. *et al.* Expression of parathyroid hormone-related protein in a human T-cell lymphotropic virus type-1 infected T-cell line. *Biochem Biophys Res Commun* 1988, **154**: 1182–1188.
14. Motokura, T., Fukumoto, S., Matsumoto, T. *et al.* Parathyroid hormone-related protein in human T-cell leukaemia lymphoma. *Ann Intern Med* 1989, **111**: 484–488.
15. Watanabe, T., Yamaguchi, K., Takatsuki, K. *et al.* Constitutive expression of parathyroid hormone-related protein in human T-cell leukaemia virus Type-1 (HTLV-1) carriers and adult T-cell leukaemia patients that can be transactivated by HTLV-1 tax gene. *J Exp Med* 1990, **172**: 759–765.