Diagnostic Images

The thrombolytic effect of heat

Joanne M. Currie and Douglas J.A. Adamson

Ward 47, Aberdeen Royal Infirmary, Foresterhill, Aberdeen, AB9 2ZB, UK

The patient

A woman of 76 who had a 13 year history of cold haemagglutinin disease presented on a cold February day with a sudden onset of breathlessness and pleuritic chest pain. A provisional diagnosis of pulmonary embolism was made, intravenous heparin was started, and her room was maintained at a temperature of 28°C. There were no obvious predisposing factors to pulmonary thromboembolism.

Investigations

Tc-99m perfusion lung scans were performed. The ventilation scans – not shown – were both normal.

Figure 1 On admission. Clockwise from top left are anterior, posterior, left posterior oblique and right posterior oblique views. There are multiple, unmatched perfusion deficits in both lungs, particularly the right. These appearances are consistent with multiple bilateral pulmonary emboli and total lung volume is reduced by 30%.

Correspondence: J.M. Currie, M.R.C.P.

Accepted: 29 June 1992

Figure 2 One week later. Clockwise from top left are anterior, posterior, left posterior oblique, and right posterior oblique views. This perfusion scan is normal and matched the ventilation scan (not shown).

Comment

Cold haemagglutinin disease usually occurs in the elderly and is caused by a monoclonal proliferation of B cells. The cold agglutinins (usually IgM kappa) are specific for red cell surface antigens and may cause spontaneous red cell clumping in the extremities, presenting as acrocyanosis in cold weather. Thrombosis resulting in gangrene occurs rarely.¹ Complement activation causes the haemolysis.

Resolution of a thrombus in one week is compatible with thrombolytic therapy but not with the simple anticoagulation our patient received (Dr F.W. Smith, personal communication). We believe that the pulmonary emboli were caused by red cell clumping related to the cold agglutinins. In such cases the effect of heat as a thrombolytic agent appears to be dramatic, returning the combined ventilation/perfusion scan to normal within a week.

Acknowledgements

We would like to thank Dr A.A. Dawson for permission to report this patient and Dr F.W. Smith for interpreting the isotope scans.

References

1. Cooper, R.A. & Bunn, H.F. Hemolytic anemias. In: Petersdorf, R.G., Adams, R.D., Braunwald, E., Isselbacher, K.J., Martin, J.B. & Wilson, J.D. (eds) *Principles of Internal Medicine*, 10th ed. McGraw-Hill Book Co., New York, 1983, pp. 1866–1867.