may relate both to duration of diabetes¹⁰ and to the severity of vitamin D deficiency and it has been suggested that vitamin D may play an important role in diabetes prevention in at-risk groups.⁵

Our observations suggest that vitamin D replacement may have a detrimental effect on glycaemic control in established diabetes. Although there was an increase in fasting insulin levels in two of the cases, as has been noted by others,^{3,5} our results suggest that the lack of perceived benefit on glycaemic control may be explained by an increase in insulin resistance. We are not aware of any previous similar reports but our findings rely on the FIRI index, which has not been fully validated against the

gold standard of measuring insulin resistance by the glucose clamp technique. Furthermore, it was not possible to exclude the influence of changes in lifestyle such as an alteration in diet or exercise level. Nevertheless, the Asian community is known to represent a group at particular risk of insulin resistance, which has recently been suggested to be an independent risk factor for ischaemic heart disease.¹² If our observations are reproducible, they question the benefit of vitamin D replacement in those with evidence of deficiency, especially in view of the associated effects on lipids. Vitamin D prophylaxis in the Asian community may circumvent such a problem.

- 1 Boucher BJ. Strategies for reduction in the prevalence of NIDDM; the case for a population-based approach to the development of policies to deal with environmental factors in its aetiology. *Diabetologia* 1995;38:1125-9.
- 2 Beaulieu C, Kestekian R, Havrankova J, Gascon-Barre M. Calcium is essential in normalizing intolerance to glucose that accompanies vitamin D depletion in vivo. *Diabetes* 1993;42:35-43.
- 3 Gedik O, Akalin S. Effects of vitamin D deficiency and repletion on insulin and glucagon secretion in man. *Diabetologia* 1986;29:142-5.
- 4 Kumar S, Davies M, Zakaria Y, *et al.* Improvement in glucose tolerance and beta-cell function in a patient with vitamin D deficiency during treatment with vitamin D. *Postgrad Med J* 1994;70:440-3.
- 5 Boucher BJ, Mannan N, Noonan K, Hales CN, Evans SJW. Glucose intolerance and impairment of insulin secretion in relation to vitamin D deficiency in East London Asians. *Diabetologia* 1995;38:1239-45.
- 6 Duncan MH, Singh BM, Wise PH, Carter G. A simple measure of insulin resistance (letter). *Lancet* 1995;436:120-1.
- 7 Finch PJ, Ang L, Eastwood JB, Maxwell JD. Clinical & histological spectrum of osteomalacia among Asians in S London. *Q J Med, New Ser* 1992;83:439-48.
- 8 Henderson JB, Dunnigan MG, McIntosh WB, Motaal AA, Hole D. Asian osteomalacia is determined by dietary factors when exposure to ultraviolet radiation is restricted: a risk factor model. *Q J Med* 1990;76:923-33.
- 9 Bansal S, Rizvi SNA, Rao MB, *et al.* Effect of hypocalcaemia on glucose tolerance, insulin release and free fatty acid levels in human subjects. *Postgrad Med* 1975;52:471-5.
- 10 Orwoll E, Riddle M, Prince M. Effects of vitamin D on insulin and glucagon secretion in non-insulin-dependent diabetes mellitus. *Am J Clin Nutr* 1994;59:1083-7.
- 11 Kocian J. Diabetic osteopathy: favourable effect of treatment with osteomalacia with vitamin D and calcium on high blood glucose levels. *Vnitř Lek* 1992;4:352-6.
- 12 Despres J-P, Lamarche B, Mauriege P, *et al.* Hyperinsulinemia as an independent risk factor for ischemic heart disease. *N Engl J Med* 1996;334:952-6.

Medical Anniversary

CARLO FORLANINI, 11 June 1847

Carlo Forlanini (1847-1918) was born in Milan, son of a doctor. He qualified as MD in Pavia (1870), worked at Ospedale Maggiore (1871-84) on skin and eye tuberculosis, and became Professor of Medicine in Turin in 1884. In 1889 he moved to Pavia where he became Professor of Medical Pathology (1889), Professor of Medicine, (1900) and Senator (1913). He died in Nervia in 1918. He is commemorated by the Carlo Forlanini Institute for Tuberculosis in Rome, which was founded in 1934. — D G James