syndrome in extensive haemangioma of the tongue and lip in a newborn infant.⁴

Thirdly, the substernal goitres reported in the literature have not been uniformly defined in relation to the proportion of the thyroid gland within the thorax. Therefore, it is rather difficult to compare the sizes and the results of reported series of substernal goitres. For the last decade, we and others⁵ have chosen to refer to any goitre in which more than 50% of its mass is inferior to the thoracic inlet as substernal.

LUC A MICHEL
JULIAN DONCKIER
Mont-Godinne University (UCL) Hospital,
Yvoir, B-5530, Belgium

Accepted 27 July 1998

- 1 Anders HJ. Compression syndromes caused by substernal goitres. *Postgrad Med J* 1998;74:327-9.
- 2 Michel LA, Bradpiece HA. Surgical management of substernal goitre. *Br J Surg* 1988;75:565-9.
- 3 Gadisseux P, Minette P, Trigaux JP, Michel L. Cerebrovascular circulation 'steal' syndrome secondary to a voluminous retrotracheal goiter. *Int Surg* 1986;71:107-9.
- 4 Ismagalov MF. The cerebrovascular circulation 'steal' syndrome in extensive hemangioma of the tongue and lip in a newborn infant. *Pediatrics* 1983;71:64-5.
- 5 Katic MR, Grillo HC, Wang CC. Substernal goitre. *Ann Thorac Surg* 1985;99:391-9.

Anaphylactoid reaction to hydroxycobalamin with tolerance of cyanocobalamin

Sir,

A patient with an anaphylactoid reaction to hydroxycobalamin but good tolerance of cyanocobalamin is described, which emphasizes the usefulness of challenge tests in cases of allergic or pseudoallergic reactions.

A 33-year-old woman with a history of Crohn's disease developed subacute combined degeneration of the spinal cord due to vitamin B12 deficiency. Replacement therapy with hydroxycobalamin was established at a dose of 10 mg intramuscularly every month with no problems for more than a year. Unexpectedly, 2 hours after a dose, the patient developed generalised urticaria and angioedema with involvement of the upper airway. Prick and intradermal tests performed with 5 mg/ml and 100 µg/ml of hydroxycobalamin, respectively, were negative. Under in-hospital observation the patient was given 2500 µg of hydroxycobalamin by the intramuscular route; 20 min later, she experienced pruritus on her palms, shortly followed by generalised urticaria, prominent lip and palpebral oedema, hoarseness and chest tightness. The patient was treated with epinephrine, methylprednisolone and chlorpheniramine with total recovery in 2 hours. A challenge test with benzyl alcohol, added as preservative, was carried out with no reaction. On the basis that the neurologic manifestations would progress without adequate replacement treatment, a desensitisation protocol was developed. Increasing doses of hydroxycobalamin, beginning with 0.05 µg, were administered every 15 min by the intramuscular route. Ten minutes after the injection of 125 µg of hydroxycobalamin, the same allergic reaction appeared. Premedication with antihistamines did not provide reliably effective protection from the hydroxycobalamin-induced reaction in the

patient. However, intramuscular challenge tests with cyanocobalamin up to 10 mg, performed on three different occasions, were followed by no reaction. At present, the patient receives 10 mg of cyanocobalamin monthly without problems.

Cobalamin is an organometallic vitamin which cannot be synthesized in the human body and must be supplied in the diet. The minimum daily requirement is about 2.5 µg. In patients with disease of the distal small intestine such as Crohn's disease, cobalamin deficiency may develop. In order to avoid clinical features of cobalamin deficiency, especially neurologic manifestations, replacement therapy is suggested. Because oral absorption is inadequate, replacement must be administered parenterally. The vitamin preparations which are used therapeutically are cyanocobalamin and hydroxycobalamin (both also called vitamin B12) given intramuscularly at monthly periods and maintained indefinitely. Allergic reactions to vitamin B12 are rare but can be observed even after several years of treatment.¹ James and Warin reported one patient with dyspnoea and urticaria in the course of a treatment with cyanocobalamin and hydroxycobalamin in which specific IgE could not be showed, suggesting an anaphylactoid reaction rather than a real allergic mechanism.² Recognising that a reaction is caused by direct histamine release may be important since treatment can generally be continued by lowering the dose of the drug. In the patient reported here, the immediate response obtained with low doses of hydroxycobalamin (125 µg) on rechallenge, the tolerance of previous doses of this drug (sensitisation period), together with the perfect tolerance of therapeutic doses of cyanocobalamin suggests an allergic mechanism even in the presence of negative skin tests. Even though the reaction developed only at or above a dose of 125 µg, it is difficult to explain this as an anaphylactoid mechanism, since the capacity of hydroxycobalamin and cyanocobalamin to induce direct release of histamine is quite similar. A reaction to an excipient rather than to the drug itself was ruled out because the only preservative in the formulation was benzyl alcohol (provided by the manufacturer) which was well tolerated by the patient on challenge. Up to now, positive skin tests with hydroxycobalamin have been described in only two patients.³⁻⁴ Accordingly, cyanocobalamin may be tried as an alternative in patients with a history of systemic reactions to hydroxycobalamin.

C VIDAL
Allergy Unit
A LORENZO
Division of Gastroenterology
Hospital de Conxo, Complejo Hospitalario
Universitario, Rúa Ramón Baltar sn, 15706
Santiago de Compostela, Spain

Accepted 27 July 1998

- 1 Ugwu CN, Gibbins FJ. Anaphylactic reaction to vitamin B12 appearing after several years of treatment. *Age Ageing* 1981;10:196-7.
- 2 James J, Warin RP. Sensitivity to cyanocobalamin and hydroxycobalamin. *BMJ* 1971;2:262.
- 3 De Blay F, Sager MF, Hirth C, et al. IgE-mediated reaction to hydroxycobalamin injection in patient with pernicious anaemia. *Lancet* 1992;339:1535-6.
- 4 Branco-Ferreira M, Clode MH, Pereira-Barbosa MA, Palma-Carlos AG. Anaphylactic reaction to hydroxycobalamin. *Allergy* 1997;52:118-9.

Salvage angioplasty following failed thrombolysis

Sir,

Drs Mahy and Jennings are correct to point out the dilemmas facing physicians responsible for the further management of patients with acute myocardial infarction and apparent failure to respond to thrombolytic therapy.¹ The lack of evidence supporting any particular management strategy is surprising given that up to 50% of patients fail to respond to thrombolytic therapy in the first few hours and that persistent ST segment elevation following acute myocardial infarction (AMI) is clearly associated with poor outcome. Purcell *et al.*² demonstrated a mortality of 18.2% in unselected patients with AMI and <50% resolution of ST segment elevation in the worst lead 60 minutes after the initiation of thrombolytic therapy. A substudy³ of the INJECT trial revealed a mortality of 17.5% in patients with ≤30% resolution of the summed ST segment elevation in leads reflecting the infarct zone. Even though it is frequently stated that such electrocardiographic (ECG) features are not 100% sensitive or specific for persistent arterial occlusion, the presence of such features must alert us to a patient who is at high risk of further adverse events. Salvage angioplasty has only been examined in one large prospective randomised study against conservative therapy.⁴ Despite a statistically significant reduction in the incidence of death or severe heart failure, this strategy has not been widely adopted nor examined further in the modern angioplasty era. This is surprising, given that this study probably underestimated the benefit of salvage angioplasty for a number of reasons. Firstly, high-risk patients, including those with a previous myocardial infarction who are perhaps more likely to benefit from attempts to open a second vessel, were excluded. Secondly, patients in this trial were taken on for salvage angioplasty relatively late after the onset of chest pain. Thirdly, intra-aortic balloon counterpulsation was rarely used, but is now known to reduce the risk of arterial occlusion following salvage angioplasty.⁵ Fourthly, the trial was performed without modern glycoprotein IIb/IIIa inhibitors, such as abciximab (Reopro®). These agents have been shown to be beneficial in high-risk angioplasty without increased risk of haemorrhage.⁶ Lastly, and most importantly, this trial was performed in the early 1990s before the modern coronary artery stent era. It is undoubtedly the case that the availability of coronary artery stents allows angioplasty in the context of AMI to be performed with greater success. We would go so far as to say that the results of the trials of immediate angioplasty following thrombolytic therapy, which universally demonstrated unfavourable outcomes with this strategy, have no relevance in the modern stent era. This is an area which commands further study. Our policy of performing salvage angioplasty in the context of <50% ST segment resolution in the worst lead 2 hours after the initiation of thrombolytic therapy has produced favourable results, especially if the patient presents promptly, receives thrombolysis promptly and the 2-hour ECG is scrupulously reviewed. Our experience is that this policy can reduce mortality from an expected 17-20% to 5%. Thus, patients with persistent ST elevation following thrombolytic therapy should be considered early for

coronary angiography with a view to salvage angioplasty. Some patients will have reperfused by the time of coronary angiography if the above ECG criteria are applied, but we believe that the potential benefits far outweigh the risks of an 'unnecessary' angiogram.

A G C SUTTON
M A DE BOLDER

Cardiothoracic Division, South Cleveland Hospital,
Marton Road, Middlesbrough TS4 3BW, UK

Accepted 27 July 1998

- 1 Mahy IR, Jennings JP. 'Rescue' after failed thrombolysis for acute myocardial infarction. *Postgrad Med J* 1998;74:355-7.
- 2 Purcell IF, Newall N, Farrer M. Change in ST segment elevation 60 minutes after thrombolytic initiation predicts clinical outcome as accurately as later changes. *Heart* 1997;78:465-71.
- 3 Schroder R, Wegscheider K, Schroder K, Dissman R, Meyer-Sabellek W. Extent of early ST segment elevation resolution: a strong predictor of outcomes in patients with acute myocardial infarction and a sensitive measure to compare thrombolytic regimens. A substudy of the International Joint Efficacy Comparison of Thrombolytics (INJECT) trial. *JACC* 1995;26:1657-64.
- 4 Ellis SG, Ribeiro-da Silva E, Heyndrickx G, *et al.* Randomised comparison of rescue angi-

- oplasty with conservative management of patients with early failure of thrombolysis for acute anterior myocardial infarction. *Circulation* 1994; 90:2280-4.
- 5 Ishihara M, Sato H, Tateishi H, *et al.* Intraaortic balloon pumping as adjunctive therapy to rescue coronary angioplasty after failed thrombolysis in anterior wall acute myocardial infarction. *Am J Cardiol* 1995;76:73-5.
- 6 The EPILOG Investigators. Platelet glycoprotein IIb/IIIa receptor blockade and low-dose heparin during percutaneous coronary revascularization. *N Engl J Med* 1997;336:1689-96.

Images in medicine

Foreign body aspiration

A 76-year-old woman with known squamous carcinoma of the post-cricoid region, previously treated with laryngopharyngectomy and radiotherapy, was referred to a tertiary hospital having aspirated her Blomsinger tracheo-oesophageal valve (size 10 mm, figure 1). Clinically she had a stridor, and was cyanosed, dyspnoeic and unconscious. Oxygen saturation was 85% on 60% inspired oxygen via a Venturi mask. Chest X-ray (figure 2) showed a massive left hydropneumothorax (pneumothorax was created by a previous aspiration) and the Blomsinger valve lodged at the right hilum. Fibre-optic bronchoscopy revealed the valve to be lodged in the right intermediate bronchus. When the valve was removed the patient rapidly improved, with an oxygen saturation of 98% on 40% inspired oxygen.

The anatomy of the bronchial tree is such that the right main bronchus is shorter and wider than the left, and it separates off from the vertical at about 25° compared to the left at 45°. This makes aspiration more common in the right lower and middle lobes.

Figure 1 Blomsinger tracheo-oesophageal valve

Figure 2 Chest X-ray

S P L MEGHJEE

Department of Anaesthesia, York District Hospital, York YO3 7HE, UK

EMMA THOMAS

Specialist Registrar in Oncology, Leeds General Infirmary, Leeds LS1 3EX, UK