Sudden unexpected death in epilepsy

S D Lhatoo, Y Langan, J W A S Sander

Summary

The majority of persons with epilepsy develop lasting remission from seizures, although mortality is significantly greater than that of the age-matched general population. Of the deaths that are thought to be directly related to seizures, sudden unexpected death in epilepsy is probably the commonest category; more so than status epilepticus or seizure-related accidents. Annual incidence rates vary from 1 in 200 patients with chronic epilepsy to about 1 in 1000 in more population-based studies. Young adults with severe, intractable epilepsy appear to be the most frequently affected group and may have even higher incidence rates. Other risk factors may also be important. An area of great research interest, several pathogenetic mechanisms have been postulated, centering mainly around cardiac rhythm and central hypoventilation. Given the frequent devastation caused by sudden unexpected death in epilepsy, the importance of seizure control is emphasised.

Keywords: epilepsy; sudden death

SUDEP: definition

A sudden, unexpected, non-accidental death in an individual with epilepsy, with or without evidence of a seizure having occurred (excluding status epilepticus), where autopsy does not reveal an anatomical or toxicological cause of death

Box 1

Epilepsy Research Group, University College London, Institute of Neurology, Queen Square, London WC1N 3BG, UK
S D Lhatoo
Y Langan
J W A S Sander

Accepted 7 June 1999

It has become increasingly clear that epilepsy can kill, not only through status epilepticus and accidents caused by seizures but through other mechanisms that cause sudden unexpected deaths. Sudden death is a real problem in the management of epilepsy and is now perhaps the commonest category of seizure-related death in patients with chronic epilepsy, often affecting young adults in this group.¹ Sudden unexpected death in epilepsy (SUDEP) is defined as a sudden, unexpected, non-accidental death in an individual with epilepsy with or without evidence of a seizure having occurred (excluding documented status epilepticus) and where autopsy does not reveal an anatomical or toxicological cause of death.² In 1910, Munson reported on mortality from the Craig Colony in New York which housed patients with intractable epilepsy and found that 99 out of 582 deaths were 'sudden'.³ Though some of these were seizure related accidental deaths, he concluded that most had occurred in the midst of a seizure, noting that "these deaths occur very rapidly at times, seizures not infrequently take place silently". Since then, there has been a somewhat reluctant acceptance of a small but significant risk of death directly attributable to seizures, and sudden death has appeared sporadically through the decades in several epilepsy mortality series as a category of death.⁴ With present day knowledge of an increased mortality in epilepsy,^{5–7} there has been renewed interest in the phenomenology of SUDEP and several studies have attempted to describe this enigma.

Epidemiological studies

It is clear from outcome studies of epilepsy that more than 70% of all patients with epilepsy enter lasting remission from the condition and up to 30% suffer chronic seizures that are difficult to control. Approximately 1 out of every 200 such patients with chronic epilepsy die suddenly and unexpectedly every year. But how common is this phenomenon in the community where the majority of patients do not have difficult, intractable epilepsy? The National GP Study of Epilepsy in the UK (NGPSE), which is an observational study of epilepsy in the community,⁸ has had only one confirmed SUDEP death in a prospective cohort of 564 patients with definite epilepsy followed up for approximately 8000 person years.⁹ The MRC Antiepileptic Drug Withdrawal Study had just two deaths attributable to SUDEP after 5000 person years of follow-up.¹⁰ These figures are reflective of the relative rarity of this phenomenon in patients who do not have chronic epilepsy, as in the NGPSE almost 70% of the cohort had achieved 5-year remission from seizures.¹¹ Similarly, the MRC study population was in remission and there is strong evidence that SUDEP is mainly a problem in the patient with intractable epilepsy.

Case ascertainment methods to determine the exact incidence of SUDEP have varied in different studies, but can be broadly categorised into studies that have used death certificates and coroner's registers,^{12–15} studies that have used treatment data, either medical,^{16–19} surgical,^{20–22} or palliative,²³ and those that have been based on data from epilepsy clinics or institutions.^{24–27} Coroner-based studies have shown incidences varying between 1:370 and 1:2100. Many of these studies, however, are flawed due to the fact that death certificates are notoriously unreliable; a large number do not accurately record the cause of death or the underlying conditions that could have contributed.²⁸ In addition, the prevalent epilepsy population is used as denominator – a presumed figure prone to variation and inaccuracy.²⁸ A recent case-control study found that annual incidence figures for SUDEP were 1.4 per 1000 patient years in both males and females.²⁹

SUDEP incidence rates of up to 1:200 per year have been reported by studies based on tertiary epilepsy clinic data,³⁰ and slightly higher figures have been reported from institutionalised patients who share features of seizure intractability with clinic subjects.^{24–26} These figures are probably close to the true incidence of SUDEP in patients with chronic epilepsy.

Table The incidence of SUDEP in different populations

Population studied	Annual incidence in population studied	Incidence in person years	Type of epilepsy	Comments
Epilepsy surgery patients ²⁰	1 in 150	—	Chronic	Severe epilepsy
Epilepsy clinic patients ³⁰	1 in 200	—	Chronic	Moderate to severe epilepsy
Patients from coroners' lists ¹³	1 in 1100	—	All	Mild, moderate and severe epilepsy
Patients from community based studies of epilepsy ⁹	—	1 in 8000	All	Mild, moderate and severe epilepsy
Patients from a case control study ²⁹	1.4 in 1000	—	All	Mild, moderate and severe epilepsy

Several studies have used anti-epileptic drug therapy lists for case ascertainment.^{17 16 19} More recent studies have been based on newer anti-epileptic drug registers such as those on lamotrigine,¹⁸ gabapentin³¹ and tiagabine.³² Some studies based on surgical data have shown particularly high incidences of sudden death (up to 1:50 per year), especially in post-surgical patients where the operative procedure has not been successful in containing seizures, reflective of a group with particularly bad epilepsy.²⁰ A more recent study of 791 patients who had received vagal nerve stimulation system implants (a seizure-reducing device that acts by stimulating the left vagal nerve), were followed up for 1335 person years and showed an incidence of definite or probable SUDEP of 4.5/1000 person years of follow-up.²³ The high mortality and incidence figures in most of these studies may reflect the severe epilepsy suffered by patients in these study populations and is likely to indicate the increasingly high risk of SUDEP in patients with chronic epilepsy.

Risk factors

SEIZURE STATUS

It is obvious from most studies that seizure control or the lack thereof, is vitally important in the mortality statistics of SUDEP and incidence figures are considerably higher in cohorts of patients with refractory and chronic seizures (hospital or clinic based) as compared to studies that are more community based. One surgical study found a particularly high incidence of SUDEP in patients unsuccessfully treated surgically for their epilepsy, with mortality approaching 1:50, in contrast to 1:150 when both pre- and post-surgical candidates were analysed together.²¹ In contrast, the NGPSE and MRC Drug Withdrawal studies, which both have large numbers of patients with mild epilepsy, have shown very low incidence figures for sudden death. It is important to note, however, that there have been reports of patients dying suddenly during their first-ever or second-ever seizure, though these are obviously uncommon occurrences. A recent case-control study showed that the relative risk of SUDEP increased with number of seizures per year (10.16; 95%CI=2.94–35.18), appearing substantially higher in patients with more than 50 seizures per year than in patients with two or less seizures per year.²⁹

AGE, SEX AND RACE

Patients with epilepsy within the 20 to 40 year age group appear to suffer higher mortality from this entity, and this has been a consistent finding in many studies.^{12 13 27} However, this does not mean that other age groups are immune and studies in children as well as older patients with epilepsy have shown significant mortality due to SUDEP. Apart from one study of children with epilepsy in a residential school, most studies have shown a male preponderance.^{13 25 27 30} One study found a higher incidence of sudden death in the African American population but this could reflect the higher incidence of epilepsy in this group.¹³

MEDICATION

Non-compliance with prescribed anti-epileptic medication has been implicated in the occurrence of sudden death and several patients have been found to have sub-therapeutic serum levels of anti-epileptic drugs.^{12 13 22 33} The significance of drug levels, however, is arguable as patients with well-controlled epilepsy can have low drug levels and vice versa. Conversely, there are reported instances of patients experiencing sudden arrhythmic deaths due possibly to the cardiac side-effects of carbamazepine,³⁴ though this is obviously not the case in the majority of SUDEP cases. The above-mentioned case-control study showed that patients on three anti-epileptic drugs were more likely to suffer SUDEP than patients on monotherapy, although it could be argued that patients with bad epilepsy are more likely to be prescribed polytherapy and the increased mortality reflects on refractory epilepsy rather than drug therapy.²⁹

Possible pathogenetic mechanisms of SUDEP

- cardiovascular
- central hypoventilation
- combined mechanisms

Box 2

NEUROLOGICAL STATUS

A higher incidence of SUDEP has been found in patients with neurological deficits and learning difficulties.²⁴ This may be attributed to the severity of epilepsy that is usually present in these patient groups and the higher all-cause mortality generally found in such patients.

OTHER

Alcohol abuse has also been implicated in some patients,¹³ as have psychotropic drugs.¹⁹ Recent head injuries have also been mentioned as a possible risk factor.

Circumstances of death

There is considerable evidence that the majority of sudden deaths in epilepsy are seizure related and circumstantial evidence of a generalised tonic-clonic seizure having occurred is often found in the form of superficial injuries, a bitten tongue or characteristic postures.^{13 30 35} This is strongly borne out by witnessed accounts of sudden death, one of which occurred during direct observation at a telemetry unit.³⁶ However, less than half of sudden deaths in epilepsy are witnessed and it is yet unclear whether all such deaths are due to seizures.

AUTOPSY FINDINGS

Autopsy findings, by definition, should not indicate an anatomical or toxicological cause of death, though this does not necessarily imply that no abnormalities are found on post-mortem examination. Pulmonary oedema is often found, although this is usually of insufficient severity to be an attributable cause of death.^{12 13 37 38} Similarly, increased liver, cardiac and brain weights are also found. Some studies have found brain lesions on autopsy^{12 13 38} that could have caused seizures in those cases. A recent study noted the presence of interstitial cardiac fibrosis in patients who had epilepsy-related sudden deaths.³⁹

Pathogenetic mechanisms

Seizure activity remains central in the various hypotheses that have been put forward to explain the pathogenetic mechanisms that underlie sudden death. These studies can be broadly categorised into those that have looked at cardiovascular causes and those that have emphasised the role of central hypoventilation during seizures.

Various rhythm abnormalities have been implicated in SUDEP, but although autonomic disturbances are common enough observations in the fitting, hospitalised, patient, there is no convincing evidence that malignant tachyarrhythmias cause significant mortality in epilepsy. There are case reports, however, of significant bradyarrhythmias and sinus arrest occurring in patients with temporal lobe epilepsy, suggesting that these could be a cause of death.⁴⁰⁻⁴² Significant ST segment changes⁴³ and QTc interval prolongations have been observed with epileptiform electroencephalogram (EEG) discharges in patients with chronic epilepsy, although the investigators also found that this prolongation remained within normal limits in almost all cases.⁴⁴ Intrinsic cardiac disease has been implicated and in one recent study a significant number of patients were found to have interstitial cardiac fibrosis, though whether this is directly related to a mechanism of death is debatable.³⁹ The influence of cardio-active anti-epileptic drugs too, remains speculative.⁴⁵

In an interesting study where seizures were chemically induced in sheep, central hypoventilation was found to be a mechanism of death.⁴⁶ Endogenous opioids that may be released during seizures have been implicated in the causation of this central hypoventilation and could well account for the suppression of respiratory drive.⁴⁷ In another study on anaesthetised and ventilated animals, elevated left atrial and pulmonary vascular pressures were found – a possible mechanism for the pulmonary oedema found in patients who have died from SUDEP, though why this should happen in the first place is unclear.⁴⁸

In another study, transient bradyarrhythmias were found to occur in association with apnoea and or a change in respiratory pattern in patients experiencing seizures, suggesting that both cardiovascular as well as central mechanisms are important and that cardiorespiratory brainstem reflexes may play a role in SUDEP.⁴⁹ In summary, however, although there are several postulates, the true mechanism of sudden death remains unclear.

Conclusion

Seizure control is likely to be an important issue in SUDEP and evidence from many studies strongly suggests an important role played by uncontrolled

Epilepsy Bereaved

PO Box 112, Wantage, OX2 8XT, UK
Tel 01235 772852/772850

Epilepsy Bereaved is a registered charity that provides valuable support and information to friends and relatives of patients with epilepsy who have suffered sudden unexpected death. Information packs on epilepsy are available from the National Society for Epilepsy helpline at 01494 601400

Box 3

Learning points

- 30% of all patients with epilepsy develop chronic seizures
- approximately 1 out of every 200 patients with chronic, severe epilepsy suffer SUDEP every year
- central hypoventilation and cardiac arrhythmias are possible mechanisms
- control of epilepsy is likely to be an important preventive measure
- all chronic epilepsy patients should be referred to specialist epilepsy units

Box 4

epilepsy. There is no doubt that the traditional emphases on drug compliance, minimisation of potential seizure precipitants, and referral of appropriate patients to epilepsy specialist units are as important as ever.⁵⁰ Although the incidence of SUDEP in community-based populations may be low, its occurrence is often devastating to relatives, carers and physicians. In patients with chronic epilepsy, where the incidence is considerably higher, the role of adequate supervision is equally important, especially amongst children and the institutionalised. As newer anti-epileptics appear on the market and the provision of both curative and palliative surgery for intractable epilepsy becomes more established, the incidence of this often tragic complication will hopefully lessen. The role of further studies examining the risk factors involved in SUDEP in this cannot be under emphasised and detailed studies such as a case-control study presently under way at this centre, should be encouraged.

- 1 Nashef L, Sander JWAS. Sudden unexpected deaths in epilepsy - where are we now? *Seizure* 1996;5:235-8.
- 2 Nashef L. Sudden unexpected death in epilepsy: Terminology and definitions. *Epilepsia* 1997;38: S6-S8.
- 3 Munson JF. Death in epilepsy. *Med Rec* 1910;77.
- 4 Rodin EA. *The prognosis of patients with epilepsy*. Springfield: Charles Thomas, 1968;
- 5 Cockerell OC, Johnson AL, Sander JW, Hart YM, Goodridge DM, Shorvon SD. Mortality from epilepsy: results from a prospective population-based study [see comments]. *Lancet* 1994;344:918-21.
- 6 Hauser WA, Annegers JF, Elveback LR. Mortality in patients with epilepsy. *Epilepsia* 1980; 21:399-412.
- 7 O'Donoghue MF, Sander JW. A historical perspective on the mortality associated with chronic epilepsy. *Acta Neurol Scand* 1997;96: 138-41.
- 8 Sander JW, Hart YM, Johnson AL, Shorvon SD. National General Practice Study of Epilepsy: newly diagnosed epileptic seizures in a general population. *Lancet* 1990;336:1267-71.
- 9 Lhatoo SD, Langan Y, MacDonald BK, Zeidan S, Sander JWAS. Sudden unexpected death: a rare event in a large community based prospective cohort with newly diagnosed epilepsy and high remission rates. *J Neurol Neurosurg Psychiatry* 1999;66:692.
- 10 Medical Research Council Antiepileptic Drug Withdrawal Study Group. Randomised study of antiepileptic drug withdrawal in patients in remission. *Lancet* 1991;337:1175-80.
- 11 Cockerell OC, Johnson AL, Sander JW, Shorvon SD. Prognosis of epilepsy: a review and further analysis of the first nine years of the British National General Practice Study of Epilepsy, a prospective population-based study. *Epilepsia* 1997;38:31-46.
- 12 Leestma JE, Kalelkar MB, Teas SS. Sudden unexpected death associated with seizures: analysis of 66 cases. *Epilepsia* 1984;25:84-8.
- 13 Leestma JE, Walczak T, Hughes JR, Kalelkar MB, Teas SS. A prospective study on sudden unexpected death in epilepsy. *Ann Neurol* 1989; 26:195-203.
- 14 Terrence CF, Wisotzkey HM, Perper JA. Unexpected, unexplained death in epileptic patients. *Neurology* 1975;25:594-8.
- 15 Langan Y, Nolan N, Hutchinson M. The incidence of sudden unexpected death in epilepsy in South Dublin and Wicklow. *Seizure* 1998;7:355-8.
- 16 Derby LE, Tennis PS, Jick H. Sudden unexplained death among subjects with refractory epilepsy. *Epilepsia* 1996;37:931-5.
- 17 Jick SS, Cole TB, Mesher MD, Tennis PS, Jick H. Sudden unexplained death in young persons with primary epilepsy. *Pharmacoepidemiol Drug Safety* 1992;1:59-64.
- 18 Leestma JE, Annegers JF, Brodie MJ, et al. Sudden unexplained death in epilepsy: observations from a large clinical development program. *Epilepsia* 1997;38:47-55.
- 19 Tennis PS, Cole TB, Annegers JF, Leestma JE, McNutt M, Rajput A. Cohort study of incidence of sudden unexplained death in persons with seizure disorder treated with antiepileptic drugs in Saskatchewan, Canada. *Epilepsia* 1995; 36:29-36.
- 20 Dasheiff RM. Sudden unexpected death in epilepsy: a series from an epilepsy surgery program and speculation on the relationship to sudden cardiac death. *J Clin Neurophysiol* 1991; 8:216-22.
- 21 Sperling MR, O'Connor MJ, Saykin AJ, Plummer C. Temporal lobectomy for refractory epilepsy. *JAMA* 1996;276:470-5.
- 22 Vickery BG. Mortality in a consecutive cohort of 248 adolescents who underwent diagnostic evaluation for epilepsy surgery. *Epilepsia* 1997; 38:S67-S69.
- 23 Annegers JF, Coan SP, Hauser WA, Leestma JE, Duffel W, Tarver B. Epilepsy, vagal nerve stimulation by the NCP system, mortality, and sudden, unexpected, unexplained death. *Epilepsia* 1998;39:206-12.
- 24 Klenerman P, Sander JW, Shorvon SD. Mortality in patients with epilepsy: a study of patients in long term residential care. *J Neurol Neurosurg Psychiatry* 1993;56:149-52.
- 25 Lip GY, Brodie MJ. Sudden death in epilepsy: an avoidable outcome? *J R Soc Med* 1992;85: 609-11.
- 26 Nashef L, Fish DR, Garner S, Sander JW, Shorvon SD. Sudden death in epilepsy: a study of incidence in a young cohort with epilepsy and learning difficulty. *Epilepsia* 1995;36:1187-94.
- 27 Timmings PL. Sudden unexpected death in epilepsy: a local audit. *Seizure* 1993;2:287-90.
- 28 Zielinski JJ. Epilepsy and mortality rate and causes of death. *Epilepsia* 1974;15:191-201.
- 29 Nilsson L, Farahmand BY, Persson PG, Tomson T. Risk factors for sudden unexpected death in epilepsy: a case control study. *Lancet* 1999; 353:888-93.
- 30 Nashef L, Sander JWAS, Fish DR, Shorvon SD. Incidence of sudden unexpected death in an adult outpatient cohort with epilepsy at a tertiary referral centre. *J Neurol Neurosurg Psychiatry* 1995;58:462-4.
- 31 Anon. Neurontin. In: *Physician's Desk Reference*. Montvale NJ: Medical Economics, 1996.
- 32 Leppick IE. Tiagabine: the safety landscape. *Epilepsia* 1995;36(suppl 6):S10-S13.
- 33 George GR, Davis GG. Comparison of anti-epileptic drug levels in different cases of sudden death. *J Forensic Sci* 1998;43:598-603.
- 34 Boesen F, Andersen EB, Kehn E, et al. Cardiac conduction disturbances during carbamazepine therapy. *Acta Neurol Scand* 1983;68:49-52.
- 35 Nashef L, Garner S, Sander JW, Fish DR, Shorvon SD. Circumstances of death in sudden death in epilepsy: interviews of bereaved relatives. *J Neurol Neurosurg Psychiatry* 1998;64: 349-52.
- 36 Bird JM, Dembny KAT, Sandeman D, Butler S. Sudden unexplained death in epilepsy. *Epilepsia* 1997;38:S52-S56.
- 37 Earnest MP, Thomas GE, Eden RA, Hossack KF. The sudden unexplained death syndrome in epilepsy: demographic, clinical, and postmortem features. *Epilepsia* 1992;33:310-16.
- 38 Hirsch CS, Martin DL. Unexpected death in young epileptics. *Neurology* 1971;21:682-90.
- 39 Natelson BH, Suarez RV, Terrence CF, Turizo R. Patients with epilepsy who die suddenly have cardiac disease. *Arch Neurol* 1998;55:857-60.
- 40 Dasheiff RM, Dickinson LJ. Sudden unexpected death of epileptic patient due to cardiac arrhythmia after seizure. *Arch Neurol* 1986;43: 194-6.
- 41 Fincham RW, Shivapour ET, Leis AA, Martins JB. Ictal bradycardia with syncope: a case report. *Neurology* 1992;42:2222-3.
- 42 Howell SJ, Blumhardt LD. Cardiac asystole associated with epileptic seizures: a case report with simultaneous EEG and ECG. *J Neurol Neurosurg Psychiatry* 1989;52:795-8.
- 43 Tiagarani S, Rasmussen V, Dam M, Pedersen S, Hogenhaven H. ECG changes in epilepsy patients. *Acta Neurol Scand* 1997;96:72-5.
- 44 Tavernor SJ, Brown SW, Tavernor RME, Gifford C. Electrocardiograph QT lengthening associated with epileptiform EEG discharges - a role in sudden unexplained death in epilepsy? *Seizure* 1996;5:79-83.
- 45 Tomson T, Kennebeck G. Arrhythmia, heart rate variability and antiepileptic drugs. *Epilepsia* 1997;38:S48-S51.
- 46 Johnston SC, Horn JK, Valente J, Simon RP. The role of hypoventilation in a sheep model of epileptic sudden death. *Ann Neurol* 1995;37: 531-7.
- 47 Ramabadran K, Bansinath M. Endogenous opioid peptides and epilepsy. *J Clin Pharm Ther* 1990;28:47-62.
- 48 Johnston SC, Darragh TM, Simon RP. Postictal pulmonary edema requires pulmonary vascular pressure increases. *Epilepsia* 1996;37:428-32.
- 49 Nashef L, Walker F, Allen P, Sander JW, Shorvon SD, Fish DR. Apnoea and bradycardia during epileptic seizures: relation to sudden death in epilepsy. *J Neurol Neurosurg Psychiatry* 1996;60:297-300.
- 50 Langan Y, Nashef L. Sudden unexpected death. *Neurol Rev Int* 1999; in press.