CASE REPORTS

Recurrent acute heart failure caused by sliding hiatus hernia

C-W Siu, M-H Jim, H-H Ho, F Chu, H-W Chan, C-P Lau, H-F Tse

Postgrad Med J 2005;81:268–269. doi: 10.1136/pgmj.2004.023416

The case is reported of a 75 year old woman who presented with recurrent nocturnal episodes of acute pulmonary oedema. The cause was uncertain as she had normal cardiothoracic ratio on chest radiography and normal left ventricular systolic and diastolic function by transthoracic echocardiogram. Another transthoracic echocardiogram was repeated when she was recumbent for an hour and had a full stomach. It showed a striking finding of severe left atrial compression by an external structure. Computed tomography of the thorax showed an intrathoracic mass behind the left atrium causing external compression of the left atrium suggestive of a sliding hiatus hernia. Cardiac catheterisation confirmed the diagnosis by showing a pronounced rise of pulmonary capillary wedge pressure in the recumbent position compared with the sitting up position.

A 75 year old woman presented with recurrent episodes of shortness of breath and chest pain in the previous three months requiring multiple admissions. The diagnosis of acute pulmonary oedema was made but no cause could be found on previous admissions. Her cardiothoracic ratio was normal on chest radiography, her left ventricular function, both systolic and diastolic, were normal by transthoracic echocardiogram. Her symptoms occurred typically at bedtime, especially after a heavy dinner, and were associated with orthopnea, paroxysmal nocturnal dyspnea, and ankle oedema. Physical examination showed regular pulses with a normal blood pressure finding of 124/61 mm Hg. The jugular venous pressure was raised, the heard sounds were normal, and no murmur could be heard. There was bilateral ankle oedema as well as basal crackles heard over both lungs. An electrocardiogram showed normal sinus rhythm without any ischaemic or hypertensive changes. Careful examination of the chest radiograph showed congested lung field with mild bilateral pleural effusion compatible with acute pulmonary oedema. There was also a round shadow behind the heart with an air-fluid level within it. Blood tests including complete blood counts, renal and liver function test, and creatinine kinase activity were within normal limits. Transthoracic echocardiography was repeated when the patient was in the supine position for an hour and had a full stomach. It showed normal left ventricular function but the left atrium was severely compressed by an extrinsic structure confirmed by multiple views (fig 1). Spiral computed tomography of the thorax showed a large hiatus hernia with intrathoracic extension. The hernia was located behind the left atrium causing anterior shift of the heart (fig 2). Subsequently coronary angiography showed normal coronary anatomy. Right heart catheterisation showed that baseline right atrial pressure and pulmonary capillary wedge pressure during prolonged supine positioning were 8 mm Hg and 18 mm Hg respectively. However, after sitting upright for 30 minutes, the right atrial pressure and pulmonary capillary wedge pressure decreased to 5 mm Hg and 6 mm Hg respectively,

confirming the diagnosis of significant left atrial compression by the sliding hiatus hernia. She was successfully treated with conservative measures including frequent small meals, avoidance of a late dinner, and sleeping in slanting position using several pillows. She had no further recurrence of acute pulmonary oedema in the subsequent 12 months.

DISCUSSION

Hiatus hernia is a common condition and its incidence increases with age.¹ It does not produce symptoms itself in most patients, but may contribute to the pathogenesis of reflux oesophagitis. Infrequently, sliding hiatus hernia may become incarcerated and strangulated, which may subsequently lead to acute chest pain, dysphagia, and a mediastinal mass.² Furthermore, cardiac compression with haemodynamic collapse has been reported in patients with complicated or large hiatus

Figure 1 Echocardiogram in apical four chamber view showing extrinsic compression of the posterior wall (arrows) of the left atrium by a large mass. LV, left ventricle; LA, left atrium.

Figure 2 Computed tomogram of the thorax showing a large, mixed type hiatus hernia compressing the left atrium from posterior aspect (arrow). LV, left ventricle; LA, left atrium; HH, hiatus hernia; Ao, descending aorta.

hernia.³⁻⁴ To our knowledge, this is the first reported case of recurrent acute heart failure caused by sliding hiatus hernia. As reported previously,⁵⁻⁶ hiatus hernia may mimic a left atrial mass on transthoracic echocardiography, and is usually shown by spiral computed tomography as shown in this case. However, the clinical significance of these findings remains unclear. In this case, we performed detail cardiac haemodynamic measurements during supine and upright posture, and clearly showed the direct compressive effect of the hiatus hernia on the left atrium. This resulted in an increase in pulmonary capillary wedge pressure and subsequently contributed to the development of acute pulmonary oedema in this patient. This case shows that hiatus hernia is a potentially reversible cause of recurrent acute heart failure; accurate diagnosis and successful treatment of hiatus hernia can prevent further recurrence of acute heart failure.

Authors' affiliations

C-W Siu, M-H Jim, H-H Ho, F Chu, H-W Chan, C-P Lau, H-F Tse, Department of Medicine, Queen Mary Hospital, The University of Hong Kong, Hong Kong

Correspondence to: Dr H-F Tse, Cardiology Division, Department of Medicine, The University of Hong Kong, Queen Mary Hospital, Hong Kong, China; hftse@hkucc.hku.hk

Submitted 23 April 2004

Accepted 23 July 2004

REFERENCES

- 1 Goyal RK. Diseases of the esophagus. In: *Harrison's principles of internal medicine*. Maidenhead: McGraw-Hill, 1994:1355-63.
- 2 Maziak DE, Todd TR, Pearson FG. Massive hiatus hernia: evaluation and surgical management. *J Thorac Cardiovasc Surg* 1998;**115**: 53-60.
- 3 Ito H, Kitami M, Ohgi S, et al. Large hiatus hernia compressing the heart and impairing the respiratory function. *J Cardiol* 2003;**41**:29-34.
- 4 Kalra PR, Frymann R, Allen DR. Strangulated gastric volvulus: an unusual cause of cardiac compression resulting in electromechanical dissociation. *Heart* 2000;**83**:550.
- 5 Yang SS, Wagener P, Dennis C. Hiatal hernia masquerading as left atrial mass. *Circulation* 1996;**93**:836.
- 6 Airolidi L, Rizzotti M, Sarasso G, et al. False left atrial mass caused by hiatal hernia: a clinical case and review of the literature. *Cardiologia* 1998;**43**:635-8.

Acute liver failure: a message found under the skin

M Meier, A Woywodt, M M Hoeper, A Schneider, M P Manns, C P Strassburg

Postgrad Med J 2005;**81**:269-270. doi: 10.1136/pgmj.2004.023382

Acute liver failure is a rare syndrome with rapid progression and high mortality. It is characterised by the onset of coma and coagulopathy usually within six weeks but can occur up to six months after the onset of illness. Viral hepatitis, idiosyncratic drug induced liver injury, and acetaminophen ingestion are common causes. This report describes the case of a 35 year old man who presented with acute liver failure shortly after binge drinking. Repeated history taking disclosed a gluteal disulfiram implant that the patient had received to treat his alcohol dependence. The patient recovered with maximum supportive care after surgical removal but without liver transplantation. This case illustrates that only meticulous history taking will disclose the sometimes bewildering causes of acute liver failure.

Acute liver failure is characterised by liver cell dysfunction leading to coagulopathy and hepatic encephalopathy, mainly attributable to viral, acetaminophen, or drug induced liver injury. Fulminant hepatitis is a rare but potentially fatal adverse reaction that may occur after the use of disulfiram, a drug used to treat alcoholism. We report a case of a 35 year old man who experienced acute liver failure associated with a gluteal disulfiram implant and alcohol misuse.

CASE REPORT

A 35 year old man first presented to a primary hospital in April 2003 with fatigue, vomiting, and vague abdominal complaints. His medical history included ongoing alcohol

misuse despite various attempts of treatment. An alcohol binge had occurred three days before admission. On examination by the admitting physicians, he was jaundiced and drowsy. Initial laboratory studies showed increased aspartate aminotransferase (24012 U/l), total bilirubin (150 µmol/l), and blood alcohol (7.7 mmol/l). Transfer to our medical intensive care unit was arranged with a tentative diagnosis of alcohol induced liver failure.

On admission, the patient appeared acutely ill with pronounced jaundice, hepatic foetor, and hepatomegaly. Auscultation and percussion of heart and lungs were normal and the patient had no clinical signs of liver cirrhosis or portal hypertension. A 2 cm scar in his left lateral gluteal region was noted. Laboratory studies in our hospital on admission confirmed a massive increase in aspartate aminotransferase (60 620 U/l), alanine aminotransferase (16726 U/l), lactate dehydrogenase (38180 U/l), glutamate dehydrogenase (12211 U/l) total bilirubin (179 µmol/l), and lactate (5.2 mmol/l). Severe coagulopathy with thrombocytopenia was present (INR 8.29; factor V 12%; 16 000/µl platelets), which precluded liver biopsy. Abdominal ultrasound showed hepatic oedema and excluded cirrhosis. The portal vein, hepatic artery, and hepatic veins were all patent. In view of progressive encephalopathy the patient was sedated and intubated. Cerebral oedema and haemorrhage were excluded by cranial computed tomography. Fluid refractory hypotension ensued, vasopressor support was begun, and anuric renal failure prompted continuous veno-venous haemodialysis. Fresh frozen plasma, platelets, packed red cells, factor XIII, and fibrinogen were given. Further laboratory tests excluded common causes of acute liver failure like viral hepatitis A-C, Wilson's, and liver autoimmune diseases as

Figure 1 Disulfiram implant excised from the left gluteal region.

judged by the absence of autoantibodies (ANA, SMA, LKM, SLA). A comprehensive drug screen was negative. High urgency orthotopic cadaveric liver transplantation was considered but declined on the basis of ongoing alcohol misuse in accordance with policies of German organ transplant legislation and Eurotransplant.

The medical history was scrutinised again to shed light on the aetiology of the liver failure. The patient was not receiving any medication and denied recreational or occasional exposure to drugs or toxic substances. It transpired that our patient had received a subcutaneous implantation of the oral drug disulfiram (Espéral) in his left buttock in Poland three months previously to “get rid of the drinking”. The implant was then immediately excised (fig 1). After surgical removal and under further supportive treatment the patient made an uneventful recovery after six days in the intensive care unit during which hepatic synthesis and detoxification normalised. He was then discharged to the referring hospital without neurological sequelae.

DISCUSSION

The main differential diagnosis in a 35 year old patient with acute liver failure would include alcohol induced liver disease, acetaminophen intoxication, viral hepatitis (predominantly HBV) as well as drug reactions and other rare diagnoses such as autoimmune hepatitis, Wilson’s disease, and Budd-Chiari syndrome.¹

Disulfiram has been in use for adjunctive treatment of severe alcoholism since 1948.² A thiuram derivative, it inhibits the second step of ethanol metabolism by inhibition of acetaldehyde dehydrogenase.³ This leads to immediate accumulation of acetic aldehyde and results in nausea, flushing, and vertigo. By virtue of this action it exerts a penalising effect on alcohol consumption.^{2,3} However, disulfiram has been widely abandoned because of its unfavourable safety profile. Inadvertent ingestion of alcohol may cause severe acetic aldehyde reaction requiring medical assistance.⁴ Fulminant hepatitis after the use of disulfiram usually occurs within the first two months after disulfiram treatment, with symptoms suggestive of acute hepatitis including fatigue, malaise, anorexia, nausea, vomiting, abdominal pain, jaundice, fever, rash, and pruritus.⁵ The pathophysiology, however, has not been elucidated.⁵ Both accumulation of toxic metabolites such as carbon disulfide, an end product of the disulfiram metabolism, and immunological mechanisms have been suggested.^{5,6} Forns *et al* concluded that disulfiram hepatotoxicity is mainly produced by the accumulation of toxic metabolites,⁶ whereas many case reports are consistent

Learning points

- Acute liver failure is a rare syndrome with rapid progression and high mortality.
- Acetaminophen ingestions, viral hepatitis and idiosyncratic drug toxicity are common causes.
- Idiosyncratic drug induced hepatitis attributable to disulfiram is a rare but well described cause.
- History taking in liver failure should include occupational exposure to toxins, alternative therapies, and herb ingestion.
- Several sets of criteria have been proposed to identify patients who will only survive with liver transplantation.

with a hypersensitivity reaction and include clinical findings such as eosinophilic infiltrates, arthralgia, fever, rash, and pruritus.⁵ Depot preparations of disulfiram have been described in the literature albeit without proper evaluation of their benefit-hazard ratio.⁷ Notably, concomitant alcohol misuse opens the possibility of aggravated reactions to drugs.⁵ Based on the literature, we believe that an idiosyncratic adverse drug reaction of disulfiram is the most probable pathophysiological mechanism, which is compatible with the course of the disease.

CONCLUSION

Our patient experienced liver failure associated with a gluteal disulfiram implant and alcohol misuse. This case illustrates that acute liver failure can have a bewildering aetiology while concomitant alcohol misuse opens the possibility of aggravated reactions to drugs such as disulfiram induced toxic hepatitis. Maximum supportive care was started only after the implant had been discovered and appreciated as a potentially reversible cause of hepatotoxicity.

Authors’ affiliations

M Meier, A Woywodt, Division of Nephrology, Hanover Medical School, Germany

A Schneider, M P Manns, C P Strassburg, Division of Gastroenterology, Hepatology and Endocrinology, Hanover Medical School

M M Hoepfer, Division of Respiratory Medicine, Hanover Medical School

Correspondence to: Dr M P Manns, Department of Gastroenterology, Hepatology and Endocrinology, Hanover Medical School, Carl-Neuberg-Strasse 1, D-30625 Hanover, Germany; mpmanns@aol.com

Submitted 23 April 2004

Accepted 30 August 2004

REFERENCES

- 1 Ostapowicz G, Lee WM. Acute hepatic failure: a Western perspective. *J Gastroenterol Hepatol* 2000;**15**:480–8.
- 2 Chick J, Gough K, Falkowski W, *et al*. Disulfiram treatment of alcoholism. *Br J Psychiatry* 1992;**161**:84.
- 3 Petersen EN. The pharmacology and toxicity of disulfiram and its metabolites. *Acta Psychiatr Scand* 1992;**369**:7–13.
- 4 Miller NS, Gold MS. *Alcohol in drugs of abuse: a comprehensive series*. New York: Plenum, 1991.
- 5 Rabkin JM, Corless CL, Orloff SL, *et al*. Liver transplantation for disulfiram-induced hepatic failure. *Am J Gastroenterol* 1998;**93**:830–1.
- 6 Forns X, Caballeria J, Bruguera M, *et al*. Disulfiram-induced hepatitis. Report of four cases and review of the literature. *J Hepatol* 1994;**21**:853–7.
- 7 Johnsen J, Morland J. Depot preparations of disulfiram: experimental and clinical results. *Acta Psychiatr Scand Suppl* 1992;**369**:27–30.