subacute endocarditis, or any badly damaged heart, also those cases in which the heart cannot be controlled with digitalis, and finally those occasionally seen in which anginal attacks have ceased with the onset of fibrillation.

CASES IN WHICH THE ADMINISTRATION OF QUINIDINE MAY BE CONSIDERED.

(a) If there is no evidence of structural damage (cardiogram). (b) Cases of recent onset, but these may occasionally clear up themselves. (c) Cases without serious valvular defects. (d) In Graves's disease if muscle is fair and an operation is in view.

If quinidine is used let the patient get fairly free of his digitalis, at the same time keeping the heart controlled; commence with 5 gr. and increase daily. First day, 5 gr.; second day, 10 gr.; third day, 15 gr.; fourth day, 20 gr.; fifth day, 25 gr.; sixth day, 30 gr.; seventh day, 40 gr., divided into three doses.

If during this period rhythm has reverted, keep same dose for two days and reduce rapidly to 5 gr. daily. If at the end of the period it has not changed stop the quinidine; it is toxic to the heart, and if the change does not occur soon it is doubtful if it is wise to force the drug. In any case it is wiser when giving quinidine to have the patient's condition controlled occasionally by means of cardiograms.

RADIOGRAMS AS AN AID TO DIAGNOSIS IN LUNG DISEASES.*

BY

A. GOODMAN LEVY, M.D. LOND.,

PHYSICIAN TO THE CITY OF LONDON HOSPITAL FOR DISEASES OF THE CHEST.

THE remarks I have to make are derived solely from the impressions which I have received in the course of routine work as physician to this hospital.

X RAY EVIDENCE.

Let us first consider what is the relative value of the X ray evidence in lung disease.

An old teacher of mine, Dr. Frederick Roberts, used to sum up examination of the chest as follows: (1) Inspection, (2) palpation, (3) percussion, (4) auscultation. To this summary of physical signs I feel we should now add (5) visualisation or X ray examination. It need not always be employed, but it should be always available. Please note carefully I am not encouraging you to resort to X rays as a short cut to diagnosis. The X ray picture is a physical sign as much as the recognised physical signs, and it should rank with

them, and be interpreted in conjunction with them. If reliance is placed on the interpretation of the X ray picture alone you are liable to serious mistakes.

Nevertheless, it must be admitted that sometimes radiograms afford information that all the other signs and symptoms taken together do not give. For instance, a new growth may be found in its early stage, although totally unsuspected. The congenital absence of the diaphragm on the left side was one of the surprises I have encountered, and a long list of instances might be given. For this reason the routine X ray examination of every chest seems to me very desirable.

Sometimes the X ray causes the physician to modify or change his diagnosis; generally it is useful in cases of doubtful diagnosis, but in the great majority of instances it simply serves as a check on diagnosis. Let me particularly warn you against the detailed and positive reports which are submitted by some X ray specialists. The frequent allusion in those reports to hilar phthisis has to be received with much scepticism—it is, in fact, better disregarded *in toto*—it is entirely unjustifiable to condemn a patient to a diagnosis of phthisis on the evidence of such reports. So also such refined points as the activity or non-activity of tubercular foci, I can assure you, are not worthy of serious attention. Regard the broad indications of your X ray not as a method of complete diagnosis, but as physical signs, and you will be in the right path.

TUBERCULOUS DISEASE OF THE LUNGS.

It is my experience that X rays usually demonstrate an extent of disease which is in advance of the clinical signs, but in cases with very definite physical signs the X ray may be in accordance with the signs as regards extent. It is especially in so-called early cases with few signs and little constitutional disturbance that we may see more than we expected, and I can show you plates of such cases in which the disease is seen to be actually very extensive, but much of what the radiogram shows is old mischief, the actual activity being localised to a small area. My present opinion is that it will probably be found that any tubercular disease of the lung sufficiently developed to give rise to symptoms and signs can throw a definite X ray shadow.

Does the converse hold good? I can say no more than this: When the diagnosis from physical signs is doubtful and the X ray throws no definite shadow, this is strong evidence in support of a negative diagnosis. I have not so far come across a case which could be definitely diagnosed as tubercular and which showed no X ray evidence, but it is likely that early miliary tubercle will not be revealed by the X rays.

Pleural Exudates.—The presence of fluid in considerable quantities is generally readily diagnosable, but it is quite easy to be in error, and to show plates which demonstrate errors both

* Abstract of a Post-Graduate Lecture delivered at the City of London Hospital for Diseases of the Chest on June 23rd, 1926.

in the physician's opinion and the radiologist's report. One of these plates shows fluid as was diagnosed, but it also indicates the presence of a new growth which was not diagnosed. Whenever pleural effusion exists, an opportunity should be taken of examining the lungs by X rays, either when the fluid has been absorbed or soon after aspiration. We may then look for signs of tubercle or of new growth.

[A series of plates were given which showed a new growth demonstrated in this way.]

Bronchiectasis.—This condition is generally easily recognisable in X ray photographs, but occasionally the details are obscured and of doubtful interpretation. In such cases recourse should be made to the injection of lipiodol into the trachea; this sinks into and delineates the dilated bronchioles in a very striking fashion. This substance likewise often renders valuable assistance in defining the course and origin of sinuses in the chest wall.

TECHNIQUE IN RADIOGRAPHY.

I think I have said enough to demonstrate the exceedingly valuable service X ray examination of the lungs affords. The use made of this method of diagnosis in this hospital is very considerable; in fact, on an average, some 50 to 60 per cent. of the in-patients are X rayed at some time during their residence. The proportion would be higher with increased facilities, and we have under consideration a system whereby every case shall be X rayed on admission. I consider this routine examination very desirable, simply as an aid to clinical diagnosis.

To make this routine examination effective, however, a high standard of technique must be maintained. At present many X ray pictures are good, but too many are bad. The number of requests for repeats have been too large. These should be, I am certain, unnecessary, and I consider it should be as possible to produce uniformly good X ray photographs as it is to produce technically perfect photographic studio portraits—it is largely a matter of standardising all procedures. It is good technique which we physicians look for before anything else. There is a natural disposition to make our own interpretations, and an expert interpretation from a bad plate does not satisfy us.

There is another point of technique which, in my opinion, requires urgent attention—that is, the accurate standardisation of position. The point of view should always be the same, or only purposely varied—we then know what to look for. This is especially important when investigating the chest, for it is frequently desirable to note alterations in the size or position of various structures. Now, unless the point of view is exactly the same in each case, judgment becomes impossible. The incidence of divergent rays upon objects in different positions and planes produces a distortion in relative size and position so that it is impossible to compare photographs in which the position of the source

of the rays has varied even to a small degree. These remarks apply with special force to serial photographs.

Further, the source of the rays should be as far removed from the object as possible—the further away the more the rays approach to parallelism and the exaggeration of the shadows of the nearer objects is reduced; a more true picture is thus obtained.

Technical points such as these are of great importance in our special work in this hospital. They seem to be at present insufficiently appreciated, but their careful consideration, which I hope we shall achieve in time, will double the value of X ray work to us physicians.

"PROSTATIC" NEURASTHENIA.

BY

J. JOHNSTON ABRAHAM, C.B.E., D.S.O., M.D.,
F.R.C.S. ENG.,

SURGEON, LONDON LOCK HOSPITALS; SENIOR SURGEON,
KENSINGTON GENERAL HOSPITAL.

If there is one sort of patient more than another who is pre-eminently a source of infinite weariness to everyone with whom he comes in contact, it is probably that unfortunate individual known as the "prostatic" neurasthenic.

He is a constant worry to himself, to his family doctor, to the series of consultants—physicians, neurologists, urologists, electro-therapeutists, and probably, in a last despairing fling, to the psychoanalysts—to whom he has been sent in the hope that some one or other of them may light on a cure. Nobody loves him. Indeed, the only person who really appears to hail his advent with delight is the quack with his pills for "lost vitality," to whom he turns eventually in despair when orthodox medicine has failed him. A short account of how he may be treated successfully may therefore be of interest.

His history is generally as follows: At some unfortunate time he acquires an attack of urethritis, sometimes gonococcal, more often due to secondary organisms such as staphylococci or coliform bacilli. For this attack he is treated more or less adequately; he never develops any definite complications; and in the efflux of time his urine clears.

But in the interval he has suffered a mental trauma. He is naturally what is called a nervous man. He is generally very intelligent. He takes a keen interest in his symptoms. He reads up the subject, examines his urine for flakes, looks for signs of recurrence, and worries continually.

Occasional attacks of phosphaturia and stinging at the penile tip after micturition occur, and keep up his fear. He develops neuralgia along the cord