THE POST-GRADUATE MEDICAL JOURNAL

VOL. I.

MAY, 1926.

No. 8.

DYSPEPSIA.*

BY

E. P. POULTON, M.D. OXF., F.R.C.P. LOND.,
PHYSICIAN TO GUY'S HOSPITAL.

DYSPEPSIA may be defined as a disorder of the functions of the upper alimentary tract. These functions are concerned with the preparation of the swallowed food and its transmission into the intestines for absorption. Disorders give rise to a variety of symptoms—e.g., pain, discomfort, nausea, &c.—and since the disorders and the symptoms are always associated, the term dyspepsia may be used to denote the symptoms, and in this sense dyspepsia is a symptom-complex. In the old meaning of the term, dyspepsia was used to denote those conditions from which organic diseases could be excluded, but nowadays it is used in a wider sense. The conditions previously called by this name are probably best called “primary dyspepsia.” Meunier quotes with approval the following definition: “A dyspeptic is one who suffers from pain in the stomach and complains of it”; but this definition is too restricted. The term “indigestion” is used in a still wider sense than dyspepsia, since it also includes disturbances in the absorption of food.

SYMPTOMS.

The symptoms of dyspepsia may be briefly enumerated *Pain* or *Discomfort* which may be experienced in the chest (heartburn), in the epigastrium, in the neighbourhood of the umbilicus, and in the region of the right and left costal margin. It is either made worse or improved by taking meals. The pain is of two kinds: (1) a pain felt deeply in the organ that produces it, and associated with tenderness on deep pressure; and (2) a referred pain in the skin due to a reflex from the organ through the spinal cord, at this level spreading out over the corresponding sensory area. It is often noticed as a soreness of the skin when it is touched or rubbed. Dr. Ryle has suggested that the presence of referred pain always means an organic lesion. This is not necessarily the case, since in experiments by Dr. W. W. Payne and myself it was noticed some hours after producing pain in the œsophagus experimentally. *Nausea*, felt chiefly at the back of the throat, a sense of *sinking* felt in the stomach and at other times a sense of *fullness* often accompanied by a desire to bring up wind, or *flatulence*. *Regur-*

gitations into the mouth, which may be acid or neutral, and the final result of reflex disturbances—viz., *vomiting*, which may terminate the attack. Certain other reflex manifestations may also be mentioned—i.e., loss of appetite, congestion of the face, palpitation, which includes tachycardia and extrasystoles; then there are drowsiness, malaise, and tiredness, hiccough and headache, while prolonged indigestion may have a depressing effect on the mental state of the patient, like any other painful disease.

The method that Dr. Payne and I adopted in order to investigate the cause of the symptoms of dyspepsia was to tie a rubber finger-stall tightly over the end of a catheter, inside which a small piece of metal tube had been inserted. The finger-stall contained a lead weight and was connected by rubber tubing to a syringe and manometer. The whole apparatus was filled with water, air being completely excluded, and the amount of water in the bag could be adjusted by means of the syringe. The bag was passed down the œsophagus into the stomach, through the pylorus into the duodenum, and it was also passed through a gastro-jejunostomy opening of one patient into the jejunum. Its position could be accurately determined by means of X rays and by means of a spirit-level, the position of the bag with respect to the manometer could be determined, and the height of the column of water in the manometer above this level gave the pressure exerted by the viscus on the bag. In patients suffering from “heartburn” it was observed that in some cases there was an increase of the mean pressure in the œsophagus with numerous contractions, while when pain was absent the pressure was only slightly above the atmosphere, although the effect of swallowing was to increase the pressure to as great an extent as was noticed when pain was present. In another case the œsophagus was shown to be dilated, while there was apparently a spasm in two places, one high up behind the top of the sternum and the other just above the cardia. The mean pressure was less than that of the atmosphere. We were, in fact, examining the intrathoracic part of the œsophagus. Every time the patient swallowed, the peristaltic movements passed downwards and were associated with pain. These peristaltic movements could also be observed in the normal œsophagus, particularly at the beginning of an examination, and in the case of my colleague a momentary pain was also observed. In a patient with gastro-jejunostomy pains were noticed in the stomach and in the jejunum, and here again an increase of pain was found to coincide with a period of increase of the mean pressure of the organ associated with large

* A Lecture delivered before the Fellowship of Medicine on March 4th, 1926.

movements of contraction and relaxation. In a patient with a "hypertonic stomach" a remarkable correspondence between the sensations and the movements of the stomach and duodenum were observed. When the stomach was empty his sensations formed a definite cycle. First there was no pain, then slight pain, then more pain, and an eructation often ended the cycle and the pain vanished. The eructation followed immediately after a very great increase of pressure in the organ and the pressure fell immediately afterwards. During the pain the mean pressure in the viscus rose. The movements and also the sensations were completely abolished by food. We also carried out observations by means of a kymograph to determine the exact timing of the pain. These observations showed that when the pain was prolonged over a certain period there was at the same time, as already described, an increase of the mean pressure with large contractions and relaxations; but, when the pain was momentary, there was only one contraction, and we found that pain was always signalled when the contraction was over—i.e., during the succeeding relaxation. Another interesting point came out of our observations, since we succeeded in getting this patient to swallow two small bags, one of which was passed

We passed bags containing air into the œsophagus and connected them with an apparatus for filling them with a measured quantity of air, and also a recording mechanism so that we obtained graphic records on a kymograph. The presence of a foreign body in the œsophagus undoubtedly gives rise to a number of descending peristaltic movements. When only a little air is put into the bag, a momentary sensation of nausea is experienced—a sensation felt in the throat—and here again the sensation was recorded after the contraction had passed the bag—i.e., during the succeeding relaxation. If more air were passed into the bag the sensation became painful, although the element of nausea could still be appreciated; but, finally, the sensation of pain alone remained. In all these cases it was after the contraction that the sensation was observed. When the bag was considerably distended with air the sensation of pain was experienced during a period of some seconds, and here again our records show that the painful sensation was felt during the time that the œsophagus was relaxed, and disappeared as the contraction passed downwards and compressed the bag. We also obtained X ray pictures, using a bag, composed of two rubber balloons, one inside the other, with a thin layer of bismuth suspension between the balloons, air being forced

into the inner balloon, so that a picture of the bag's opaque wall was obtained. When the air-bag was fully distended with 30 or 40 c.cm., the pain was continuous and no contraction succeeded in getting past the bag at all. When the bag contained less air peristaltic contractions succeeded in passing down the œsophagus, narrowing and elongating the bag as they went. The pain was found to vanish at the moment that the bag was compressed and elongated by the peristaltic wave. The sensation of sinking in the stomach was also associated with a momentary lowering of pressure, and in this case the muscle-fibres are probably relaxed; but there was no rise in the mean pressure.

We can now discuss a theory as to the genesis of

visceral sensation, and I am going, as far as possible, to omit all reference to the word "tone," which has been used in different senses by different people. The stomach, the bladder, and to a less extent the œsophagus are capable of having a small or a large capacity depending normally on the amount of their contents. Sherrington talks about the posture of the organ being small or large ("postural tone").

FIG. 1.

Diagram illustrating variations of pressure in the œsophagus (A) when the patient complained of pain; (B) when pain was absent. S, represents the movement following a normal swallow. As high a pressure was recorded as was obtained when the patient had pain. (Figs. 1 and 2 from *Quarterly Journal of Medicine*, W. W. Payne and E. P. Poulton, 1923, xvii., 53.)

into the stomach and the other one into the duodenum. We found that exactly the same movements were present in the duodenum as in the stomach; but were felt just one second later. This seemed to show that the pylorus was widely patent at the time.

Dr. Payne and I carried out further observations, the results of which have not yet been published.

A big posture is obtained by lengthening or rearrangement of the muscle-fibres of the viscus. Such a lengthening or rearrangement normally occurs gradually, as the contents of the organ gradually increases. If the lengthening occurs suddenly, we get a sensation. This at first can be appreciated by the individual as a sensation of nausea if it occurs in the oesophagus, or a sensation

body—the finger, ulcer, or whatever it may be—and I would ask you to test our theory the next time you do a rectal examination. I have always found that when the anal sphincter contracts firmly on the examining finger, the patient's sense of pain or discomfort disappears for the moment.

Judging by what I have said, the size of the organ is not of so much importance in causing sensations

FIG. 2.

Tracing of the movements in the duodenum (upper curve) and stomach (lower curve). The arrows mark a painful sensation and corresponding points on the tracing. There is no significance in the different sizes of the two tracings because two different types of recording apparatus were used.

of emptiness and sinking if it occurs in the stomach. If the sudden lengthening is more pronounced this sensation alters to that of pain. The latter tends to disappear after a time, when the muscle-fibres take up their normal lengthened posture without stretching, or it disappears intermittently every time a peristaltic contraction shortens the fibres concerned. If the fibres are greatly stretched, as by the presence of an unyielding foreign body in the oesophagus, no contraction can get by at all and the pain is continuous. After a time adjustment may occur and the fibres become capable of contracting. The pain then becomes intermittent or griping, and finally may disappear altogether, even though the foreign body still remains in position. This theory takes no account of spasm causing pain, and hence it is opposed to the ordinary views prevalent at the present time. In our opinion, the ordinary spasm, such as is produced in the anus, is nothing but a series of peristaltic contractions produced by the presence of a foreign

as the fact that its muscle-fibres are being suddenly extended. In the case of the stomach, there are the "hypertonic"—high placed and transverse, stomachs, and the so-called "hypotonic" stomachs—elongated and dropped; and both these types are compatible with perfect health and with complete absence of dyspeptic symptoms. As a matter of fact, the position of the stomach depends upon the build of the individual. The stout man, with broad chest, horizontally placed ribs, has a highly placed stomach; while the long-chested man, with more vertical ribs, wide intercostal spaces, and narrow costal angle, has what is called a dropped stomach. The position of the stomach is not in itself of importance, but the liability of its muscle-fibres to undergo sudden extension. Symptoms are experienced when a stomach of, say, normal position, becomes suddenly dropped. It is the change which is important. I should like to mention Dr. Barclay's observations on the effect of smelling asafetida, which caused a sensation of

nausea and sinking, and at the same time the stomach dropped while the patient was being observed under the X ray screen.

Flatulence is usually regarded as due to swallowed air. Fermentation in the stomach is almost certainly only a very rare cause, but another factor which has not been considered in modern times is the evolution of carbon dioxide which occurs when the alkaline contents of the duodenum and the acid contents of the stomach are mixed together. An evolution of gas under these circumstances was found to occur by Beaumont over 90 years ago.

Vomiting that so often terminates the sensations of a dyspeptic attack is due to reflex contractions of the abdominal wall. Barclay found that if the lesser curvature of the stomach was stimulated by electricity the abdominal muscles contracted, while the stomach remained stationary. Vomiting is not necessarily a painful process, as I can testify myself, when a bag was pushed out of the œsophagus by an abdominal contraction without any sensation at all. However, vomiting as it ordinarily occurs in dyspepsia is extremely painful, but this is probably because the contents are forced out by the stomach through a closed cardia.

Since the symptoms of dyspepsia are due to associated motor defects in various organs, their study often cannot determine what is the organic lesion responsible for these disturbances, since an organic lesion may cause at one time reflex disturbances in the œsophagus, another time in the stomach, another time in the duodenum, and so forth. It is true that there may be a tendency for certain lesions, such as a duodenal ulcer, to produce a definite clinical picture, but it is nothing more than a tendency, and diagnosis based on clinical observation alone would be nearly as often wrong as right. An instance will make this clear. It is commonly held that duodenal ulcer is associated with a hypertonic stomach, hunger pains, and hyperchlorhydria, but Conran found that X ray appearances suggestive of duodenal ulcer occurred in 23 per cent. of patients in whom the stomach was dropped, and a dropped stomach in 49 per cent. of cases was associated with gastric hyperacidity, while hypo-acidity was only present in 21 per cent. For diagnosis we must rely on other means. In lesions of the stomach and duodenum, X rays will help to diagnose the presence of a chronic peptic ulcer and carcinoma. A test-meal will indicate the presence of carcinoma, while occult blood will be present in the stools in both these types of lesions and also commonly in gastritis, according to my experience. Of the lesions outside the stomach that cause dyspepsia the most important are the gall-bladder—i.e., cholecystitis or gall-stones—appendicitis, pancreatitis, and tabes dorsalis. Occult blood will not be present in the stools with these lesions. X rays will help in the diagnosis of gall-bladder disease and appendicitis, and with regard to the former we have the introduction of the new dyes “tetra-bromphenol phthalein” and the iodine compound. Pancreatitis and tabes must be

diagnosed by their special tests. In considering these lesions removed from the stomach, but causing dyspepsia, a distinction must be made between “referred pain” and “reflex pain.” Appendicitis may cause referred pain in the area of skin which is innervated by the same segment of the spinal cord, but reflex pain, in contrast to the former, means that the afferent impulses pass up and down the spinal cord, causing disturbances in organs at a distance with corresponding symptoms. Thus, the presence of heartburn—an œsophageal disturbance—has been noted after pressing on an inflamed appendix. Gall-stones with jaundice produce painful gastric sensations. Stimulation of the duodenum electrically at operation has been seen to cause contraction of the greater curvature of the stomach. There are organs in other systems of the body that may cause reflex dyspeptic disturbances—the heart, or vessels in angina pectoris, renal calculus, or movable kidney—and more general agencies, such as pregnancy, sensitiveness to foreign protein, hypoglycæmia caused by insulin, wasting disease, anxiety, and lead poisoning.

With regard to gastritis as a cause of dyspepsia, I cannot resist quoting to you observations by Beaumont carried out on his patient, Alexis St. Martin, between 90 and 100 years ago, because his observations are of fundamental importance, and they are not known as widely as they should be. You will remember that St. Martin's stomach was lacerated as the result of a gunshot wound, and a permanent fistula resulted. Beaumont's observations were carried out over a series of years.

“The inner coat of the stomach in its natural and healthy state is of a light or pale pink colour, varying in its hues according to its full or empty state. It is of a soft or velvet-like appearance, and is constantly covered with a very thin, transparent, viscid mucus, lining the whole interior of the organ. Immediately beneath the mucous coat, and apparently incorporated with the villous membrane, appear small, spheroidal, or oval-shaped glandular bodies from which the mucous fluid appears to be secreted.

“By applying aliment or other irritants to the internal coat of the stomach, and observing the effect through a magnifying glass, innumerable minute lucid points and very fine nervous or vascular papillæ can be seen arising from the villous membrane and protruding through the mucous coat, from which distils a pure, limpid, colourless, slightly viscid fluid. This fluid, thus excited, is invariably distinctly acid. The mucus of the stomach is less fluid, more viscid or albuminous, semi-opaque, sometimes a little saltish, and does not possess the slightest character of acidity. On the application of aliment the action of the vessels is increased, the colour brightened, and the vermicular motions excited.

“In febrile diathesis, or predisposition, from whatever cause—obstructed perspiration, undue excitement by stimulating liquors, overloading the stomach with food, fear, anger or whatever depresses or disturbs the nervous system—the villous coat becomes sometimes red and dry, at other times pale and moist, and loses its smooth and healthy appearance; the secretions become vitiated, greatly diminished, or entirely suppressed; the mucous coat scarcely perceptible; the follicles flat and flaccid, with secretions insufficient to protect the vascular and nervous papillæ from irritation.

“There are sometimes found, on the internal coat of the stomach, eruptions or deep red pimples, not numerous, but

distributed here and there upon the villous membrane, rising above the surface of the mucous coat. These are at first sharp-pointed and red, but frequently become filled with white purulent matter. At other times irregular, circumscribed red patches, varying in size or extent from half an inch to an inch and a half in circumference, are found on the internal coat. These appear to be the effect of congestion in the minute blood-vessels of the stomach. There are also seen at times small aphthous crusts in connexion with these red patches. Abrasion of the lining membrane, like the rolling up of the mucous coat into small shreds or strings, leaving the papillæ bare for an indefinite space, is not an uncommon appearance.

"These diseased appearances, when very slight, do not always affect essentially the gastric apparatus. When considerable, and particularly when there are corresponding symptoms of disease, as dryness of the mouth, thirst, accelerated pulse, &c., no gastric juice can be extracted, not even on the application of alimentary stimulus. Drinks received are immediately absorbed or otherwise disposed of, none remaining in the stomach ten minutes after being swallowed. Food taken in this condition of the stomach remains undigested for 24 or 48 hours or more, increasing the derangement of the whole alimentary canal and aggravating the general symptoms of disease.

"Whenever this morbid condition of the stomach occurs, with the usual accompanying symptoms of disease, there is generally a corresponding appearance of the tongue. When a healthy state of the stomach is restored, the tongue invariably becomes clean.

"IX. July 28th.—9 o'clock A.M.; weather clear: wind N.W., brisk; thermometer 66°; stomach empty; not healthy; some erythema, and aphthous patches on the mucous surface. St. Martin has been drinking ardent spirits pretty freely for eight or ten days past; complains of no pain, nor shows symptoms of any general indisposition; says he feels well, and has a good appetite.

"XII. August 3rd.—7 o'clock A.M. Inner membrane of stomach unusually morbid; the erythematous appearance more extensive, and spots more livid than usual; from the surface of some of which exuded small drops of grumous blood, the aphthous patches larger and more numerous, the mucous covering thicker than common, and the gastric secretions much more vitiated. The gastric fluids extracted this morning were mixed with a large proportion of thick ropy mucus, and considerable mucopurulent matter, slightly tinged with blood, resembling the discharge from the bowels in some cases of chronic dysentery. Notwithstanding this diseased appearance of the stomach, no very essential aberration of its functions was manifested. St. Martin complains of no symptoms indicating any general derangement of the system, except an uneasy sensation, and a tenderness at the pit of the stomach, and some vertigo, with dimness and yellowness of vision on stooping down and rising again; has a thin yellowish-brown coat on his tongue, and his countenance is rather sallow; pulse uniform and regular; appetite good; rests quietly, and sleeps as well as usual."

Finally, I quote the following experiment, which shows that Beaumont had really elicited the origin of hunger pains.

"Experiment 65.—To ascertain whether the sense of hunger would be allayed without the food being passed through the œsophagus, he fasted from breakfast time till 4 o'clock P.M., and became quite hungry. I then put in at the aperture three and a half drachms of lean boiled beef. The sense of hunger immediately subsided and stopped the borborygmus or croaking noise caused by the motion of air in the stomach and intestines, peculiar to him since the wound, and almost always observed when the stomach is empty."

I will now shortly discuss the principles of treatment. It is a familiar circumstance with dyspeptic patients that, though they can tolerate one kind of food, another, which one would suppose to be very

similar in texture, is not at all well tolerated. This condition has been called "selective dyspepsia," and there has been a tendency to attribute it more to mental disturbances than to any real alteration in the response of the stomach to types of food. Beaumont's observations, however, suggest the fact that the stomach really has a very different power of dealing with what are apparently very similar substances. The time that the stomach takes to get rid of various types of food are tabulated in the book, and the results, I must admit, are surprising. Thus, an uncooked egg, whether whipped or not, leaves the stomach in one and a half hours; a soft-boiled egg in three hours; and a hard-boiled egg and fried egg in three and a half hours; baked or roast potatoes leave the stomach in one and three-quarters hours, boiled potatoes in three and a half hours; pigs' feet, soured and boiled, leave the stomach in one hour; roast pork, consisting of both fat and lean, leaves the stomach in five and a quarter hours; fresh salmon trout, boiled, leaves the stomach in one and a half hours; salted salmon, boiled, in four hours; cereals, such as rice, sago, tapioca, barley, vary between one and two hours. These observations, to my mind, open a possible field of investigation of the greatest importance in the treatment of dyspepsia. A sound rule in treatment is based on the fact that protein foods, such as meat, call forth and require a highly acid juice for their digestion, while fat which, when mixed with other foods, forms a pellicle on the surface of the gastric contents, causes a diminution in the secretion of hydrochloric acid and a regurgitation of bile and remains a long time in the stomach. Consequently, fat and protein food should, as far as possible, be taken at separate meals. Small quantities of food taken at frequent intervals is another sound principle. The intake of fluids must be regarded from two points of view. There is what Meunier calls the "water of hydration," which is all the fluid which is required to replace that lost in the urine and sweat, and there is the fluid, much smaller in quantity, which should be taken with a dry meal in order to give it the porridgy consistence which makes it most suitable for the gastric juice to act on. The "water of hydration" should be taken half an hour before meals so that it leaves the stomach quickly and alone. Beverages should be taken warm; they may include water, weak tea, cocoa, milk, and a beverage containing diastase, which may be prepared by putting a tablespoonful of malted barley in a teacupful of water and standing this in a saucepan of boiling water for ten minutes. No alcohol should be allowed. The most valuable drugs are the carbonates—the so-called alkalies. These act by liberating carbon dioxide in the stomach, which stimulates contraction, and so prevents stretching of the muscle-fibres. The carbonates may be given in the form of sodium bicarbonate gr. xxx., or prepared chalk gr. xxx., or magnesium carbonate gr. xxx., or the two latter may be mixed. It is unnecessary to use the

expensive bismuth carbonate. The liberation of CO_2 may be facilitated in certain cases by immediately following the dose of carbonate with citric acid gr. xxx. dissolved in 1 oz. of chloroform water. In cases where the hydrochloric acid in the gastric juice is diminished, hydrochloric acid may be given by mouth with advantage, especially in rosacea. Powdered kaolin has been used with the idea of covering over abrasions of the mucous membrane. I am sceptical of its value, and I am also sceptical of the value of the various bitters very commonly prescribed.

The application of local heat to the epigastrium is a treatment of dyspepsia hallowed by long use. Observations by Dr. Payne and myself have given an indication as to how such counter-irritants are effective. Rubbing the skin over the sternum, with or without a counter-irritant, caused a profound effect on the movements of the œsophagus. In one case a sensation of "tingling," experienced some time after the rubbing had taken place, was associated with very considerable peristaltic activity of the œsophagus.

Local heat may be applied most simply in the form of a flannel bandage or a rubber hot water

bottle. Thermogene wool, antiphlogistine, and poultices may be used.

When pain is severe and the patient unable to vomit, complete relief may be obtained by the removal of the contents of the stomach by means of a tube. In cases of gastroptosis or, as has been already described, in cases where the stomach is apt to undergo a sudden drop with corresponding sensations, exercise such as walking on the hands and feet strengthens the abdominal muscles. Massage of the abdominal muscles is of value, but one of the most effective of all treatments is by means of a belt. But it is important that considerable pressure should be exercised by the belt on the lower part of the abdomen just above the symphysis pubis. It may be advisable to have a conical pad made and fastened inside one of the more usual forms of belt, so that this pressure may be exercised. It probably acts by preventing the sudden dropping of the stomach, and its value can be assessed by an observation of Leven's that by pressing the fist into the epigastrium of these patients pain is produced, which may be completely alleviated by pressing in with the other fist just above the symphysis.

REMARKS ON MANIPULATIVE SURGERY OR BONESETTING.*

BY

FRANK ROMER, M.R.C.S., L.R.C.P. LOND.

FROM time to time much criticism has been published by the lay papers on the after-treatment of bone and joint injuries by our profession, whilst the treatment by bonesetting or manipulative surgery—as it is now termed—by unqualified practitioners, has been proportionately extolled. The question occasionally arises as to the probable origin of this humble branch of surgery. Dr. Wharton Hood, to whom the profession was originally indebted for the elucidation of the bonesetter's secrets, held the opinion that "the first bonesetter was the servant or unqualified assistant of a surgeon who had known exactly what could be done by sudden movements and how the movements should be executed." This theory is, of course, possible. Waterton, the naturalist, who many years ago wrote much on the subject, says that "this art is practised not only in England but throughout Europe as well, and that in Spain he bears the significant name of *Algebrista*." As a matter of fact he does not; the name is *cirujano algebrista*, in France he is termed *un rebouteur*, but Waterton is correct in saying this mode of treatment exists throughout Europe. Personally

I believe that some such rough-and-ready method of treatment has always existed, but that its utility got overlooked during the time surgery was developing into a definite science. As our professional ancestors progressed in their knowledge of pathology, such a condition as tubercular disease began to be more generally understood, and the importance of quietude in this and other acutely inflamed joints was soon realised. In consequence of the general appreciation of the value of absolute physiological rest in these cases, it was not long before it was considered essential for the treatment of all painful and swollen joints, no matter from what cause. The cult of complete rest became to be deemed so necessary that it was never considered advisable to use a joint as long as the least pain was experienced on attempted movement, thereby reminding one of the boy who was forbidden to enter the water till he had learned to swim. The universal adoption of these principles by the profession caused anything in the way of more vigorous treatment of injuries to remain in the hands of the less educated classes who, "rushing in where angels fear to tread," frequently obtained brilliant results from their unorthodox methods. Thus it happened that many vendors of herbs and simples resident in country places came to specialise in this business, which, unlike the majority of folklore remedies, contained, in the words of *The Lancet*, "some long-forgotten truth."

HISTORY OF BONESETTING.

The only book on this subject for many years was by Dr. Wharton Hood, who, in 1871, from personal observation of a celebrated bonesetter called Hutton, was able to disclose to us the methods

* A Lecture delivered before the South-West London Post-Graduate Association on March 17th, 1926.