PLATE III.

Mr. C. Price Thomas

Lung Abscess

FIG. 1.

FIG. 2.

FIG. 3.

These three films taken at intervals of 4 weeks between first and second and 2 weeks between second and third show the progress of a case of spontaneous cure. It is in the absence of such evidence that surgical treatment is undertaken. Fig. 1 also shows the fluid level so characteristic when seen.

PLATE IV.

Mr. C. Price Thomas - - - - - Lung Abscess

FIG. 5.

FIG. 4.

The above A P and Lateral films of a case already draining into the bronchial tree. Note the absence of fluid level, owing to granulations preventing entry of air to cavity. Their value in localisation is also well shewn, the abscess being posterior and near the mid line; ribs 7 and 8 would need resection from transverse processes forward.

LUNG ABSCESS.

By C. PRICE THOMAS, F.R.C.S.

(Assistant-Surgeon to Brompton Hospital for Consumption and Diseases of the Chest;
Assistant-Surgeon to Westminster Hospital.)

Definition.

The term, *Pulmonary Abscess* is reserved for an abscess in the parenchyma of a lung in which the bronchial tree is normal as opposed to the abscess which is associated with a pre-existing bronchiectasis and which is termed a bronchiectatic abscess.

Ætiology.

Pulmonary abscess in about 60 per cent. of cases follows operations on the upper respiratory tract; Whittemore, in 100 cases, gives the causes as follows:—

66 cases after operations on the upper air passages under general anæsthesia, 22 following some type of pneumonia; septic infarction in 3 cases, tracheo-oesophageal fistula complicating carcinoma of oesophagus in 1 case; in 8 cases the cause could not be determined.

The commonly accepted modes of infection are: (1) by the blood stream; (2) by the lymphatics; (3) *via* the bronchi.

It has been very difficult to produce pulmonary abscess experimentally, either by the introduction of infected material into the bronchi or into the blood stream. Hunter, however, by introducing glass into the bronchi of rabbits to produce bronchial irritation and then injecting streptococci into the blood stream, succeeded in producing an acute abscess similar to that met with in human pathology, and more recently Löffler induced pulmonary infarction by injecting sterile paraffin into the blood stream and then occluding the bronchus of the corresponding area with a sterile foreign body. These combined procedures resulted in a pulmonary abscess of the chronic putrid type. It is interesting to note that neither pulmonary infarction nor bronchial obstruction, separately, had the desired effect and also that although sterile agents were used, a putrid abscess resulted, the organisms in this case probably finding entry through the bronchus.

It is probable that both vascular and bronchial occlusion play a part in the production of lung abscess in the bulk of cases, although in those cases of septic embolism from a distance such as a sinus thrombosis or appendicitis, it seems probable that vascular embolism is solely responsible.

In the cases following tonsillectomy or other operations on the upper air passages, the mechanism for the production of bronchial occlusion or vascular embolism is present and it seems probable from experimental evidence that if sufficient care be taken to prevent aspiration of blood or fragments of tissue into the lung and so avoid bronchial occlusion (one of the favourable factors) that this type of case already dignified by the name of post tonsillectomy abscess may be to a great extent eliminated.

The cases of abscess following pneumonia occur most probably as a direct result of the inflammatory process as in abscess in other parts of the body, although it is possible that there may be an extension by lymphatic spread from the bronchial tree, the alveoli of which are filled with secretion, into the lung parenchyma.

In those cases following abdominal operations which result primarily from vascular embolism, it is interesting to recall that lobar and lobular atelectasis is a

not uncommon feature after laparotomy. It may be that bronchial occlusion may be a feature in the production of this group as well. Pulmonary abscess and subphrenic abscess are not uncommonly associated without direct communication and these are possibly produced by lymphatic spread.

The organisms infecting a pulmonary abscess are very varied, streptococci, anaerobes and spirochaetes and the fusiform bacilli are the commonest, although the staphylococcus, pneumococcus and others have been described. The foul smell which is associated with the large percentage of abscesses is due to anaerobes and the spirochaetes, and the black colour of the discharges met with in some cases to the *B. melanogenicum*.

Symptomatology.

The history is very varied. The onset may be dramatic or insidious, there may or may not be a history of a preceding operation, either a few days or weeks before, or of pneumonia.

The outstanding feature in the early stages is *pain* which may be fairly well localized or diffuse on the affected side and may vary from a fairly sharp pain to a dull ache; in some cases pain is entirely absent. The amount of pain apparently depends on the proximity of the abscess to the pleura.

Cough is a feature of the disease, but in its early stages it is practically absent and it is when the abscess is near a bronchus, producing bronchial irritation, that the cough commences. At first, the cough is non-productive and later, there is some frothy sputum which is non-offensive, but the sputum only becomes foul and purulent after the abscess has ruptured into a bronchus (the latter event occurs from 2 to 6 weeks after the onset of the disease). There is, however, foetor in the breath some few days before the abscess ruptures. There may be haemoptysis, rarely fatal, some time before the abscess has ruptured into a bronchus, but this occurs as a rule after this has happened. There is usually a pyrexia up to 101-102° although it may be higher, and the disease sometimes starts with a rigor; the pulse rate is raised in the early stages up to about 100 and in uncomplicated cases it rarely exceeds 120; the respiration rate in an uncomplicated case is not markedly raised and, except in very large abscesses, or in very frail people with a low vital capacity, there is rarely dyspnoea despite the fact that there may be slightly cyanotic facies. Clubbing of the fingers is common and may occur very rapidly, well marked clubbing developing in 4-6 weeks.

In early cases, there may be nothing but a raised temperature and a slight increase in pulse rate with malaise and nothing else that may direct attention to the lungs, and it is only the antecedent history and the having in mind the possibility of lung abscess which will lead to the investigations which will provide the diagnosis.

Physical signs.—Physical signs may be very scanty; in the early stages if the abscess is peripheral, that is close to the chest-wall, there may only be a little dullness with perhaps deficient air-entry and some fine râles in the surrounding lung. If the abscess is deeply seated in the lung with normal lung overlying it, physical signs will be completely absent. When the abscess is discharging through a bronchus, the physical signs will vary according to the position of the opening in relationship to the accumulated discharges. If the opening is above the level of the fluid content, there may be cavernous or amphoric breathing and whispered pectoriloquy and if the opening is below the level of the fluid, coarse bubbling râles. At any time during the course of the disease, the physical signs

may be those of consolidation, localized to the area of the abscess; to sum up, the physical signs may be completely absent or any of those met with in chest disease.

Blood examination will shew a polymorphonuclear leucocytosis.

X-rays are undoubtedly the *most valuable aid to diagnosis* and are often the only means by which it can be made. The picture varies according to the stage of the disease at which it is taken, that is whether before or after the abscess has ruptured into a bronchus.

All cases should have antero-posterior and lateral radiograms taken in the erect posture and in those at the apex it is valuable also to take oblique ones as well to obviate the dense shadow of the pectoral girdle.

In those cases where the abscess has ruptured into a bronchus and thus permitted the entry of air into the cavity, the abscess shews as a roughly circular clear area with a surrounding zone of consolidation and a fluid level; it is often of value to take a picture with the patient lying on the affected side as this shews an alteration in the fluid level. (Fig. 1, Plate III).

In cases before rupture into a bronchus, the shadow is roughly circular but is uniformly opaque owing to the absence of air in the cavity; in some cases even after rupture into a bronchus there is no evidence of a fluid level, this being due to the valve-like action of granulations allowing only of escape of pus and preventing air getting in.

If the abscess is complicated by an empyema, the diagnosis may be obscured until the pus has been removed either by aspiration or operation, as the whole side of the chest may merely show a more or less uniform opacity.

The injection of lipiodol into the bronchial tree serves to differentiate pulmonary abscess from a bronchiectatic abscess by outlining the normal bronchi. The lipiodol never enters an acute abscess owing to the granulations at the drainage opening, but when the abscess has become chronic and granulations are not present, it is quite possible to fill the cavity.

Needling the chest as a diagnostic procedure should never be undertaken when there is the slightest suspicion that the condition is a pulmonary abscess. It is a sound principle in thoracic surgery never to needle a chest until the chest has been radiographed and a diagnosis has been made. *Diagnostic needling should be confirmatory and not a preliminary step in making a diagnosis.* The reason for condemning exploratory puncture is that in any given case the chances are that the lung may not be adherent to the chestwall and a needle introduced into a pulmonary abscess across a free pleura most probably, though not inevitably, will lead to infection of the pleural sac with dire results.

The diagnosis of pulmonary abscess rests mainly on the X-ray picture and on the character of the sputum after the abscess is draining into a bronchus; it is, however, important to remember that pulmonary neoplasms may get infected and give rise to X-ray appearances, physical signs and sputum identical with an abscess and in patients over 45, in whom there is no definite history, it is well to keep the possibility of malignant disease in mind. The treatment probably will be more or less the same but the prognosis is of course completely altered.

Complications.

The complications of Pulmonary Abscess with or without operation are the following.

Cerebral Abscess is the most dread of them all, as in the well established case it is fatal in the large percentage of instances. The bulk of surgeons veto operation even if the abscess is localizable, as the operation is inevitably fatal and a certain small percentage of cases which have all the symptoms of a cerebral abscess will, if left alone, settle down and survive. In some cases, however, there results a recrudescence of symptoms even years after with fatal consequences. Brain abscesses are the result of embolism from the thrombosed vessels in the abscess wall, the clot of course being septic. They are often multiple, attacking both sides of the brain; it is not uncommon to find postmortem two abscesses, one large and a small one on the opposite side, the small one only having given rise to symptoms during life. It is interesting to note also that sometimes the brain abscess is chronic with thick walls in association with a more recent acute encephalitis, an obvious flare of an old focus.

Empyema. This may occur spontaneously or as the result of treatment. When it occurs spontaneously, it is generally of gradual onset and consequently of limited extent, it being rare to meet with a complete pyothorax in this connection. The occurrence of a spontaneous empyema does definitely increase the mortality but not anything like to the extent that the empyema which occurs as the result of injudicious treatment does. The reason of the special seriousness of the latter type is probably its rapid onset; the sudden flooding of a normal pleural cavity with the pus from a ruptured abscess is identically the same as the rupture of an appendix abscess into a free abdomen. In each case the subserous lymphatics are in the optimum condition for the absorption of toxic products and before the patient, who is already suffering from toxæmia, can put up any barrier of resistance, the balance is turned against him. In those cases of gradual onset, on the other hand, the pleural and sub-pleural tissues gradually formed a barrier of granulation tissue and hence absorption is decreased and the organism as a whole is capable of dealing with the new focus. It is this lack of appreciation between the modes of onset of empyema complicating abscess which leads to conflicting statements as to the prognostic significance of this complication.

Pulmonary Gangrene. This hardly is a complication of lung abscess, but rather a more severe degree of the original disease. It is said by some to be due to infection by the streptococcus putridus; it is of course fatal when extensive.

The opposite lung. During any part of the disease, there may be an aspiration to the opposite side resulting in broncho-pneumonia which may lead to a fatal termination.

Suppurative Pericarditis may terminate the case, usually in association with empyema.

Hæmorrhage and lardaceous disease may complicate as in suppuration elsewhere.

Treatment.

Treatment resolves itself under two headings, Medical and Surgical.

There is no shadow of doubt that certain pulmonary abscesses are cured spontaneously; the abscess ruptures into a bronchus and is coughed up with an apparent clinical and radiological restoration to normal. (Figs. 1, 2 and 3, Plate III.)

General. In the early stages of the disease before the abscess has ruptured, rest in bed with the usual medical treatment of a pyrexia is indicated. If the cough is troublesome and non-productive, a sedative such as linctus heroin should be given so as to give the patient rest. When, however, rupture into a bronchus has occurred, all efforts should be directed to aid evacuation of the abscess. This may be achieved by a stimulating expectorant, e.g., creosote by mouth, on a mask, or in a creosote chamber. In view of the presence of spirochaetes, salvarsan substitutes, intravenously or intramuscularly, may be used, although their value is very problematical and any beneficial effect they may have is probably due to the effect of arsenic as a tonic and not as a spirochæticide.

Postural drainage is of definite value and should be practised as frequently as the patient's condition will allow; the optimum position for emptying the cavity is usually found by the patient himself, although study of the radiograms will sometimes indicate a position not already tried.

Bronchoscopy. The chief factors preventing free bronchial drainage are the small size of the opening, and the presence of exuberant granulations at its mouth, and to obviate these bronchoscopic aspiration may be instituted.

Bronchoscopy should be done under local anaesthesia, with preliminary omnopon and scopolamine, the soft palate and posterior pharyngeal wall painted with 15 per cent. cocaine on swab applicators and the superior laryngeal nerves anaesthetised by placing a swab with the same solution in each vallecula and held there. If the patient is instructed to breathe quickly through the mouth, retching is reduced to a minimum. 10 mins. of 20 per cent. cocaine is now injected into the trachea and by coughing the solution is distributed and the larynx and whole bronchial tree anaesthetised. The patient should be instructed to spit after each application in order to minimise the risk of cocaineism.

When the bronchoscope is introduced, pus will be seen in the bronchi of the affected side and this is aspirated. A powerful suction pump is necessary, the flexible end of the aspirating tube being introduced, if possible, into the cavity. Often, however, granulations obstruct the bronchus or the opening is too small, when the granulations should be painted with 15-20 per cent. silver nitrate or the opening dilated with dilators. Some writers advocate lavage of the cavity with a variety of antiseptics, although this does not seem to enhance the value of the procedure. Bronchoscopic aspiration is of value in those cases which can be cured by bronchial drainage although their number is small, but in the large majority it seldom leads to cure. The procedure, however, does definitely decrease the toxæmia and increases the patient's sensation of well-being, the sputum also tends to lose a great deal of its offensiveness and its quantity may diminish.

Artificial Pneumothorax. In view of the value of *artificial pneumothorax* in the closure of cavities in tuberculous disease, it was only natural that it should be tried in pulmonary abscess. Some writers still advocate its use in this connection and many successful cases are recorded; it is necessary, however, to sound a note of warning as pneumothorax can be followed by very grave consequences. Needless to say, pneumothorax is never instituted unless there is free drainage into a bronchus and it is fairly universally accepted that peripheral abscesses should not be subjected to a pneumothorax. The most dangerous complication of this treatment is rupture into the pleural cavity and when it is

considered that the disease is an acute inflammatory one, it is not surprising that the majority are to a greater or lesser degree adherent to the chest wall. When one also remembers the oedematous and friable condition of the surrounding lung, it will be seen that the risk of rupture is by no means small and practical experience confirms this, at least in dealing with the peripheral type of case. The result of flooding a relatively normal pleura with the highly toxic contents of an abscess has already been noted and this risk is the chief reason for the unpopularity of this method of treatment.

It may be concluded that central or hilar abscess would be ideal for pneumothorax, but even in these circumstances rupture is liable to occur. Added to this risk is also the fact that even though rupture does not occur, the effect of the pneumothorax may not be as happy as was anticipated, firstly, because drainage may not be as free after treatment as it was before, and, secondly, although drainage may be free, the cavity may not be collapsed. It is true that Forlanini succeeded in closing a cavity after two years of refills, but at the present time the risks of this method would hardly be justifiable.

Surgical Drainage. The bulk of pulmonary abscesses in the acute stage call for surgical drainage. Every case should have a preliminary period of medical treatment with postural drainage, and, if necessary, bronchoscopic aspiration in the hope that recovery will result, but these methods should not be persisted in for more than 6 weeks after bronchial drainage has been established. Persistence beyond this point lays the patient open to chronicity of the abscess and secondary bronchiectasis, both of which preclude of successful result by external drainage. During the course of medical treatment, note should not merely be kept of the clinical condition and quantity and character of the sputum, but serial radiograms should be made and any indication that the disease is stationary or progressive is a sign that active measures should be taken. In the very acute cases, where toxæmia is so grave as to imperil life, surgical intervention will be necessary even before the abscess has ruptured into the bronchial tree.

Before undertaking surgical treatment it is essential that good lateral as well as antero-posterior views be taken a day or two immediately prior to the operation so that the present state of affairs can be estimated and the abscess accurately localised. It is only by means of the A.P. and Lateral views that the latter can be done. (Figs. 4 and 5, Plate IV.) Some people also have stereoscopic films taken, but except in apical lesions, where they may be of value, they cannot supplant the A.P. and Lateral views. In apical lesions, owing to the opacity of the pectoral girdle, it is practically impossible to get a satisfactory lateral view, and oblique and stereoscopic films will be of value.

Anaesthesia. Before discussing the question of anaesthesia, it is well to review one or two of the problems involved. The patients as a rule are suffering from a fair degree of toxæmia and any general anaesthetic used must be of the least toxic order, e.g., nitrous oxide. The lung underlying is acutely inflamed with the vessels in the cavity wall filled with septic emboli and this infected tissue at some stage in the operation has to be traversed by a needle which puts the infected area into communication with the airway system of the lung either through puncture of sound lung or through the cavity itself; in other words, a track is formed which communicates the site of the thrombosed vessels with the intrapulmonary air pressure. Nitrous oxide is used as a general anaesthetic usually, as this does not increase the already existing toxæmia and

also allows of anaesthesia without abolishing the cough reflex, and so decreases the risk of aspiration. But it presents certain difficulties. Even when there is no positive pressure, there is always a certain amount of congestion and its value, because the cough reflex is not abolished, is a disadvantage in that the cough is also not controllable and the tendency between the two factors is to increase the intrabronchial pressure so that during the needling a fragment of infected clot may be forced into the general circulation and a cerebral abscess result; the following illustrative case demonstrates the possibility.

G. K., aged 57 years, suffering from an abscess at right apex, following S.M.R. Operation, two stages, both under gas and oxygen anaesthesia. First operation, resection segments of 1st, 2nd and 3rd ribs, posteriorly, post-operative period uneventful. Second operation, needling and opening of abscess which was packed. The same day and the following, patient was unconscious, flaccid paralysis left arm and leg and face. Next day, conscious but completely mentally deranged. For 5-6 days following, mental condition clearer, periods of complete lucidity, alternating with periods of derangement. At the end of the 4th day, movement in arm and leg returned but remained weak, in 10 days, apparently complete recovery. Patient, however, still complained 5 months later, that on turning sharply, he comes down heavily on left heel.

The question arises in the above case whether or not the embolism was a gas embolism or not. It is more than probable, however, that some infected material was carried along and thus the condition must be considered as a chronic cerebral abscess.

Local anaesthesia is the anaesthetic of choice, as it fulfils the conditions nearest the ideal. The operation can be performed quite painlessly, the patient's cough reflex is retained and the patient can give warning that he wishes to cough so the risk of needling during an increase in the intrapulmonary pressure can be to a great extent eliminated. Local anaesthesia with novocaine is also relatively non-toxic.

The operation is most often performed in two stages. The first consists in resecting ribs and introducing an extrapleural pack to induce adhesions and the second carried out, 7 to 10 days later, is the finding and opening of the abscess.

The abscess is localised as accurately as possible by the means at one's command, both clinical and radiological, the latter being most valuable. The patient is placed on the table in a position which is most comfortable for him and which at the same time gives good access for the surgeon.

The incision is made vertically to allow of resection of lengths of 2 to 3 ribs and is placed over the site of the abscess. The skin is infiltrated along the line of the incision with $\frac{1}{2}$ per cent. or 1 per cent. novocaine and then the muscular layers down to the level of the ribs; anaesthesia will be sufficient to carry on with the operation as soon as the infiltration has been done. The coverings of the chest-wall are divided down to the ribs; it is advisable at this point to inject the intercostals of the ribs to be resected, injecting 2 to 3 ccs. of novocaine under the subcostal groove and at the upper border of each rib to be removed, and by the time the bleeding points have been tied off, anaesthesia will be complete; it is well to warn the patient before stripping the periosteum that he will hear a scraping noise and also a crack when the rib is cut; the upper border of the rib should of course be stripped from behind forwards and the lower border from before backwards because of the direction of the external intercostals. About 2 to 3 inches of each rib is removed. After the ribs have been resected, the intercostal bundles including the muscle and vessels are tied at the level of the cut ends of the

ribs both in front and behind, and the intervening portion of the bundle is removed; this leaves the pleura with the thin strips of periosteum fully exposed.

It is now time to assess whether the two layers of pleura are adherent or not, and it is not out of place to stress here that if there is the slightest doubt about it, it is absolutely essential to treat the case as though they were not. The extra week's wait will rarely affect the course of the disease and the slightest risk of needling or opening the abscess across a free pleura and producing an empyema should be avoided at all costs. To reverse an old surgical aphorism, "when in doubt, do *not* drain" sums up the position. When the pleura is undoubtedly adherent, it will be found to be fibrous and hard, nearly stony hard, and this is the only condition which permits proceeding; many cases will, of course, be adherent, but experience has shown that even when one is certain that the lung is adherent, the area of adhesion is irregular and a free pleura is opened in part of the incision when it is least expected.

The great majority of cases need a two-stage operation, and after the intercostal bundles have been resected, a swab soaked in iodine is laid over the outer surface of the pleura; it is an advantage to incorporate a metal ring in the swab as this will shew clearly in the subsequent radiograms whether localisation has been accurate or not. The chest-wall is sewn up in layers over the swab, preferably with a small tube for drainage. There is always a large effusion around the swab in these cases, and it has been occasionally found that the effusion has been so large as to change the relationship of the abscess to an otherwise accurately localised rib section. Not all the effusion is drained off by the tube and a certain amount of compression inevitably occurs, occasionally, with very satisfactory results in small abscesses which have thereby been emptied and cured by extra pleural packing only.

The extra pleural pack is left for 7 to 10 days and by the end of this time, adhesions are sufficiently firm to carry on with the second stage. In about 1 per cent. or 2 per cent. of cases, however, the pleura may not be firmly adherent. In one case, for example, after 12 days between the stages and with what appeared to be a firmly adherent pleura, nearly half the area was found to be non-adherent on incision; air-tight suture of pleura to lung was immediately performed but the patient developed an empyema with gravely toxic symptoms, although, fortunately, ultimate recovery resulted. Occasionally, in basal cases, instead of a plastic adhesion of the layers, an effusion occurs generally of limited extent. This demands repeated aspiration through an openly packed wound until the lung becomes adherent.

The second stage involves needling of the lung and drainage. The patient is given omnopon and scopolamine and is placed on the table in the same position as for the previous stage; if care is taken the wound can be re-opened without any anaesthetic, but, if necessary, infiltration of the muscular layers at a little distance from the wound will produce perfect anaesthesia; the pleura is of course anaesthetic, as the nerves were cut at the previous operation.

When the wound has been reopened, and swabbed clean, the lung is needled; this should be the first time that a needle has been introduced into the chest. If and when pus is aspirated, the needle is left *in situ* as a guide into the cavity; sometimes merely gas is aspirated as the cavity may contain only a small quantity of pus at the time of operation, the foul odour of the gas and perhaps a few flakes of pus will be sufficient indication that the cavity has been entered. On the day of operation it is advisable to discourage the patient's cough and of course avoid

postural drainage so that the cavity may be as full as possible. Before needling the lung, it is well to warn the patient not to cough if he can possibly help it, and that if he wishes to, he must give the operator due warning so that the needle may be withdrawn before he coughs.

Diathermy, where possible, should be used instead of a knife as it tends to seal the vessels beyond the area of thrombosis and so minimises the risk of embolism. Also hæmorrhage is much more easily controlled and it is also possible by means of a loop end to remove portions of lung tissue with ease and little hæmorrhage.

The cavity is opened into along the needle, having previously coagulated the surface with a button diathermy; when the cavity is open, portions of lung substance and pleura are removed with the loop. In the peripheral abscess, it is sometimes possible to remove the outer wall of the abscess and so saucerise the cavity; with those more deeply placed it is wiser to be content with enlarging the hole of entry so as to establish efficient drainage rather than attempt saucerisation which would necessitate removal of a great deal of normal lung.

The contents of the abscess vary enormously from foul thin purulent fluid to thick, solid, inspissated material, also generally very foul; in those cases which have not already ruptured into a bronchus, it is not uncommon to find the contents odourless, these cases having been apparently embolic in origin. The contents of the abscess are gently removed and the cavity explored with the finger. Great care must be exercised in doing this so as to avoid any risk of detaching the adherent lung from the pleura and also to avoid disseminating emboli from the abscess wall. The cavity is lightly but uniformly packed with roll gauze impregnated with a mixture of vaseline and paraffin wax, taking care to remove the excess of paraffin from the gauze before packing as the paraffin will be squeezed out into the bronchi draining the cavity. The packing is carried on until the whole wound is packed, the superficial part of the wound is more firmly packed to keep it open, superficial dressings are now applied and the patient returned to bed.

In cases which are done in one stage, apart from the risk of opening the free pleura, there is only one practical consideration and that is the liability to get a spreading cellulitis of the chest-wall, consequently the superficial part of the wound should also be lightly packed to allow of adequate drainage. The danger of cellulitis in the two-stage operation obviously is not so great as the tissues around the wound have been closed off by the inflammation induced by the presence of the swab; this inflammation, by the way, is of course sterile.

The wound is not touched except for removal of the outer dressings for 4 to 5 days and during this time the temperature remains raised and the patient continues to bring up foul sputum. This should cause no concern. After 4 to 5 days, the gauze plugging is removed. An anaesthetic, gas and oxygen, will be necessary for very nervous patients, and the wound redressed in a manner similar to formerly; the plugging need only be changed every other day until all the sloughs have separated, and then a daily inspection is carried out. A thoracoscope is very valuable for examining deeply situated abscesses. After a varying period of time, plugging becomes impracticable and a tube is inserted instead, the tube being shortened and made smaller according to the exigencies of the case; in all these cases, there are one or more bronchial fistulæ into the cavity and the tube should not be discarded until these are closed. The discharge is at first very offensive and at this time there seems to be some grounds for the administration of arsenic intravenously as it certainly seems to cut the period of offensiveness

short. However, in any case with adequate drainage the foul smell diminishes and disappears and the final discharge which is already sweet becomes mucopurulent and finally mucoid; on no account must irrigation of the cavity be undertaken in view of the bronchial fistulæ.

Healing takes place as in abscesses elsewhere, by fibrous tissue and it is the pull of this tissue added to the cough and infection of the bronchi which is responsible for the production of secondary bronchiectasis. It will be appreciated that the ideal time for healing is when the walls of the abscess are soft and come together easily. It is in the cases that are allowed to become chronic that secondary bronchiectasis is most liable to occur, as in these there is considerable fibrosis in the wall of the abscess before open drainage is established, and of course the element of infection and cough has a longer time to have its effect. In cases that are in what might be called the subacute stage, it is well to allow of relaxation of the lung either by a generous rib resection or by a supplementary diaphragmatic paralysis.

The average time taken for healing in these cases is from 2 to 4 months, occasionally some take less and others considerably more, in those that take longer there is usually a fair degree of chronicity associated with some degree of secondary bronchiectasis. The complications after operation are the same as those already discussed and the added one of bronchial fistula, this complication in the absence of an associated bronchiectasis, if the cavity has been obliterated, is amenable to treatment and closure can be achieved by one of the many methods devised.

A certain number of cases are either not recognised or are not presented for treatment until the abscess has been present for 9 months to a year and by this time the condition is a thoroughly chronic one with thick walls and in some cases with the cavity more or less completely epithelialised. It is hopeless to expect that these cases will be cured by open drainage. Even when epithelialisation has not commenced, the ultimate result will be non-closure of the cavity and a secondary bronchiectasis and where epithelialisation has started, closure obviously can never be achieved by simple drainage. The only treatment which will cope with this type of case is lobectomy.

In the new growth of the lung which has become infected, the diagnosis, although it may be suspected, is only made on exploration. These cases are usually not amenable to radical treatment but drainage does get rid of their toxæmia and makes them comfortable. The importance in the recognition of these cases is the prognostic significance.

Conclusions.

The important points to be remembered in connection with this disease are:—

- (1) That the onset is often insidious and physical signs may be scanty or absent.
- (2) The great importance of radiology in the diagnosis and management of the case.
- (3) That needling should never be undertaken until the diagnosis has been established and only when it has been established without doubt that the two layers of pleura are adherent and even then only as an immediate preliminary to open drainage.
- (4) The importance of establishing open drainage as soon as the case under conservative treatment has ceased to improve.