immunity with which the Medical School of St. Mary's Hospital has been for so long identified, and that future work may lead to a convergence of the "cellular" and "humoral" aspects of immunological research of the activities of the streptococci.

SUMMARY.

The examination of agglutination, toxin and antitoxin relationships of the various "pathogenic" streptococci suggests that there is a very close relationship between all the members of the group. The further analysis of this close relationship is the problem before us.

RHEUMATISM.*

BY

CHARLES SUNDELL, M.D. LOND.,

PHYSICIAN IN CHARGE OF THE CHILDREN'S DEPARTMENT, PRINCE OF WALES'S GENERAL HOSPITAL; SENIOR PHYSICIAN TO THE SEAMEN'S HOSPITAL, GREENWICH.

FOR the purpose of this lecture it is proposed to confine the conception of rheumatism within definite limits excluding both osteo-arthritis with its atrophic and hypertrophic bone changes and rheumatoid arthritis with its rarefaction of bone, muscular atrophy, and its undoubted relation to absorption from septic foci obvious or occult. When these diseases are excluded there remains a large group of rheumatic conditions with which alone it is proposed to deal to-day.

Rheumatism is met with in two forms, acute and chronic. There is evidence that these are variations of one underlying state differing only in the mode and intensity of the response of the tissues to the irritation of a single morbid cause.

At first sight it is, perhaps, surprising that the dramatic signs and symptoms of rheumatic fever in a child should be regarded as identical in origin with the lumbago of his grandparent. But experience has much support to lead to this view. Careful inquiry into past history will again and again reveal the fact that an adult sufferer from fibrositis or "neuritis" has been marked as the prey of rheumatism during the years of his early childhood by growing pains, tonsillitis, or acute fibrile arthritis. It is not uncommon for the progress of a definite acute rheumatic fever into a slow crippling chronic rheumatic process to occur, or for alternations of acute joint affections and attacks of chronic fibrositic and muscular pains to be seen in the same individual. Such observations emphasise the latency of the rheumatic state, the chronicity of the disease, and its similarity in this particular to gout.

In connexion with the latency of this disease mention may be made of its frequent appearance

in an occult form, sapping energy and destroying the sense of well-being. Children are particularly prone to this manifestation; they may suffer for weeks or months from fatigue before definite limb pains or joint swelling make their appearance. Delayed or restless sleep at night may be followed by intense sleepiness in the morning and lack of energy during the day. Such symptoms of ill-health in a child should always arouse the suspicion of rheumatism and the call for extra rest be generously answered. In the adult a very similar state of affairs is often seen. For weeks or months before the appearance of undoubted rheumatic signs there may exist a lowering of mental and physical vitality that makes work and play burdensome and clouds all joy in life. Under appropriate antirheumatic treatment these unpleasant symptoms are among the earliest to disappear.

Rheumatism may be described as a chronic state of the body characterised by certain chemical and physiological stigmata, essentially chronic in its course, liable to periods of great activity, marked by long periods of latency and by slowly progressing phases of slight activity, influencing particularly fibrous and synovial and muscle tissues, and producing a definite lowering of vitality both physical and nervous.

CAUSES OF RHEUMATISM.

Infection with germs from mucous surfaces is for many a satisfactory explanation of the origin of this disease. The supporters of this view must, however, explain why: (1) case-to-case infection is unknown; (2) why members of certain families are peculiarly prone to the disease in successive generations; (3) why the great antirheumatic specific, salicylate, so powerful in rheumatism is so powerless in other infections with the same or closely allied organisms.

Careful considerations of problems such as these force one to the conclusion that the upholders of the infection theory see only part of the truth. If germs play a part in rheumatism, as they undoubtedly do in rheumatoid arthritis, may it not be that their rôle is a subsidiary one and that the real fundamental cause of the disease is to be sought in some aberration of body chemistry and physiology that renders the tissues vulnerable to germ attack?

Three features of the rheumatic state are of great interest and importance: (1) During periods of latency, quiescence, or merely fibrositic pain the skin is abnormally dry. (2) During such periods the temperature is always abnormally low. (3) During spontaneous active rheumatism and under the influence of artificially induced fever sweating is profuse and strongly acid in reaction. These points merit a little elaboration. The dry skin of the rheumatic is sometimes obvious; sometimes questioning is necessary to elicit the fact that sweating is always very slight or that

* A Lecture delivered at the North-East London Post-Graduate College.

formal habits of normal sweating have been lost. In the case of children habitual dryness of skin may be interrupted by occasional night-sweats when they are asleep in a warm bed. In the absence of other cause for such night-sweating rheumatism should always be suspected.

The low temperature of the rheumatic is a constant feature in the adult, rarely does it stand above 97.6°F . In children whose thermal regulation is notoriously unstable brief irregularities may occur from slight causes, but apart from these exceptions the rule of the low temperature holds good in children as in adults.

Acid sweat is, perhaps, the most constant feature of all. It is found in all conditions that are rheumatic in origin. In acute rheumatic fever it is spontaneous, in chronic conditions it has to be induced by artificial measures. In the normal individual the reaction of sweat from clean skin is slightly acid, that from the leg being slightly more acid than that from the arm or trunk; most acid of all is the sweat from the palms and soles. In rheumatism the same rules hold good, but if any local manifestation of the disease is present as, e.g., an inflamed knee or stiff neck, the sweat from the skin in such an area is found to be more acid than in any other part of the body.

Chemical tests show that this abnormal acidity of the sweat is due to the presence of lactic acid. This is a product of muscle metabolism that is increased under conditions of venous congestion, impaired nutrition, or weakened innervation of the muscle concerned. It is normally broken down by processes of oxidation to CO_2 and H_2O , but traces are to be found in the urine. Under circumstances of severe muscular work the content of lactic acid in the blood, lymph, and urine may rise temporarily to ten times its normal.

We have found so far that the chronic rheumatic state is associated with a dry skin and low temperature and that the acute rheumatic state is associated with free sweating and fever. By physical measures it is possible to make the chronic rheumatic state behave in regard to temperature and sweating exactly like that of the acute rheumatic.

The question arises: What happens to acute rheumatic fever if it is left alone? Before the discovery of the antipyretic and analgesic effects of the salicylate group of drugs rheumatic fever was treated by rest, warmth, local anodynes, alkalies, and opium. Judging from the writings of physicians of those days recoveries were as frequent and relapses no more common than they are to-day. It is well recognised that salicylates are powerless to control the heart lesion of rheumatism.

With some misgivings I began some years ago to treat rheumatic fever without drugs, relying entirely on the eliminative power of the skin to deal with the condition. The results have been most encouraging. Temperature comes down gradually, pain is of short duration, and joint swellings subside quickly. Sweating continues to be profuse and strongly acid for one to three weeks

after pain and swellings have disappeared. There appears to be no greater incidence of heart complications than under orthodox methods, transient apical murmurs during the febrile period disappearing completely as convalescence is established. Acute rheumatism appears, therefore, to be a self-curative disease. Encouraged by such results, one may endeavour to reproduce them in chronic rheumatic diseases by stimulating fever and free sweating—i.e., by conversion of a chronic state into an acute self-curative process.

Such a procedure is the aim of the "pyretic treatment" of rheumatism initiated by Percy Wilde, of Bath. Its employment in cases of chronic rheumatism has given me completely satisfactory results. The stimulation of fever and sweating is achieved by means of hot packs once or twice a day, or the use of a special couch in which the patient lies in an atmosphere of moist warmth at $100\text{--}105^{\circ}\text{F}$. A rise of body temperature of $3^{\circ}\text{--}4^{\circ}$ is quickly obtained and profuse sweating occurs.

Each treatment lasts 20 minutes to half an hour; if he is otherwise capable the patient can continue his ordinary mode of life while undergoing the course. Until a balance is obtained between the degree of febrile response and the amount of sweating there may be a temporary increase in the severity of the symptoms or the appearance of rheumatic pains and stiffness in joints hitherto free from discomfort. This stage never lasts longer than three or four baths and is succeeded by rapid amelioration of all symptoms; the mental depression and lassitude referred to above are usually the first symptoms to disappear. As the course of packs or baths is proceeded with it is found that the acidity of the sweat becomes less and less; finally after 12–30 baths it becomes normal and cure can be pronounced; long before the result has been attained the patient feels completely cured, but unless he remains under treatment until his sweat is normal in reaction he is liable to suffer sooner or later from a recurrence of his symptoms. These packs and baths have an educative action upon the skin, and it is usual to find that the habit of adequate sweating induced by them persists after the baths themselves have been discontinued.

It has been my aim to lay before you suggestions as to the nature of rheumatism, the essential identity of the acute and chronic rheumatic condition, the self-curability of acute rheumatism, and the curability of chronic rheumatism by simple physical measures through the conversion of a chronic into an acute process.

THE new pathological department and research institute of the Victoria Park Hospital for Diseases of the Chest, the opening of which on July 19th by the Minister of Health is referred to at p. 174 of our present issue, was presented to the hospital by the Prudential Assurance Company, and its equipment was provided for by an anonymous donor of £500, and by the Ladies' Association of the hospital, which raised £400.