INTRODUCTORY

By MAURICE DAVIDSON, M.A., M.D., F.R.C.P.

(Physician to the Brompton Hospital for Diseases of the Chest, Consulting Physician to the London County Council)

The increased employment of radiography in the diagnosis of chest disease during the last quarter of a century is one of the many notable advances in the science and art of medicine. So great has been the improvement in radiographic technique and in the efficiency of radiological interpretation that X-ray diagnosis has now reached a very high standard of accuracy. Like other scientific advances, radiology has in the course of its development met with no little scepticism and even with active opposition from clinicians. This recalcitrant attitude, however, is becoming more and more a matter of past history. The time has long since gone by when X-ray examination could be regarded as a mere ancillary of clinical medicine. In the diagnosis of pulmonary disease most of the medical profession have at last come to regard it as a *sine qua non*: the successful education of the clinician in this respect is greatly to be welcomed, and is a matter for congratulation to the radiologists, who have had to bear the brunt of much trivial and ill-informed criticism. Now the fruit of their labours is ripe, and radiology has won its merited place in the combination of various methods of examination essential to modern team-work.

One of the most outstanding features of X-ray work to-day is the employment of mass radiography, in which increasing interest is being shown by all sections of our profession. The introduction of this method of examination on a really large scale is certainly an innovation; its universal adoption and its application to vast numbers of the civil population are far-reaching measures which raise many important questions and difficulties entirely outside the range of the purely technical aspect of the subject. In the following articles the reader will find a digest of the matter presented by various writers, each dealing with the question from a different point of view. As in other subjects, it is impossible to avoid a certain amount of controversy, especially where new and comparatively untried methods are concerned; in this symposium the pros and cons are set forth. It is part of the Editorial policy to encourage the free and impartial consideration of all matters that concern the health of the nation and the welfare of the profession and its clientèle. In the following introductory pages, which I have been asked to write in place of an Editorial, I shall endeavour fairly to consider the difficulties which universal introduction of mass radiography is likely to entail, with due regard to some of the criticisms that have been advanced, while at the same time insisting upon those features which have led me to regard it as a valuable contribution to preventive medicine and as a method which must, sooner or later, become an integral part of our medical service.

The introduction of any great scheme of investigation involving mass examination of large numbers of persons in a community will inevitably present difficulties, especially in the early stages of its adoption; this, as I have already pointed out, is particularly true of mass radiography applied to vast numbers of the industrial population. It is, however, unfair to adduce this as an argument against the method. Difficulties are made to be overcome, and criticisms to be of any real value must be constructive. Dr. Brailsford's controversial article is useful in so far as it is provocative and of potential service as a stimulus to those in search of guidance as to how best to deal with the increase of pulmonary tuberculosis already in evidence since the outbreak of this war, and likely to appear in even more serious degree within the next few years. His observations on the Team and its work and on the reading of the films are, perhaps, useful in calling attention to some of the administrative troubles which may possibly have been underestimated by those who are pressing for the universal adoption of the scheme: he does moreover rightly refer to what is probably the most important problem, viz., the fact that at present we lack sufficient skilled personnel, equipment, and accommodation to treat the cases found in the present system, let alone the extra numbers, hitherto unsuspected, which will be discovered by the new method. These, however, are not insuperable obstacles, and there are points in some of the other articles in this series which suggest constructive efforts to meet and to overcome them. Many of his objections ignore the whole weight of experience of expert team-workers who have devoted some years to investigation of the clinico-radiological questions involved; not a few of them consist in a contradiction of propositions which no expert phthisiologist would venture to assert: e.g. "*No wise physician would be prepared to accept a chest as normal*

merely because the radiograph showed no abnormality" (my italics). No wise physician of the present day has ever done so! Other trenchant criticisms of Dr. Brailsford's unhappy aversion to mass radiography will be found in Dr. Brian Thompson's well-written commentary with its fine sense of proportion and appreciation of relative values. This article, expressed as it is in clear and restrained language, provides a good judicial summary of the arguments. Dr. Thompson is of opinion that the possibilities offered by the present E.M.S. hospital network for post-war expansion are likely to offset the risk of our being suddenly overwhelmed by large numbers of fresh tuberculous cases discovered by mass radiography. Whether this hope will be fulfilled in practice remains to be seen, but it is at least a reasonable answer to some of the objections that have been raised.

The admirable review by Toussaint and Pritchard of the modern pathological conception of young-adult phthisis and its relation to early diagnosis is, perhaps, the most important contribution to this general survey of the question of mass radiography. The following passage may be quoted as providing information which is a vital preliminary to any clear understanding of the plea for extension of the present limited scope of radiographic examination: "Recent surveys have shown that some 40 per cent, more or less, of adolescents may be going out into the world of office, factory, and workshop without any previous encounter with tuberculous infection. . . . *It is almost certainly in this group that there arise many of the acute lesions, fatal within a few months to a year of onset, in the absence of diagnosis and treatment, and preceded even shortly before the onset of the disease by radiograms showing lungs quite clear of abnormal shadows*" (my italics). Realising the practical difficulties of a universal scheme applied to a whole industrial population, these authors have given an interesting account of an experiment in selective mass radiography which they have carried out in Bermondsey and Southwark based on the symptomatology of tuberculosis. The encouragement to the general practitioner to co-operate with the specialist, which is one of the outstanding merits of their scheme, receives valuable testimony in the article by Dr. Scott, who has collaborated with Toussaint and Pritchard from the time of its inception. The results are of considerable interest and importance, and although the scheme from the point of view of those who advocate universal mass radiography has certain obvious lacunae, as a workable solution of the general practitioner's difficulties and as an example in co-operative effort it has much to commend it.

Space does not permit me to enter in any detail into some of the more remote problems that will inevitably be raised by the introduction of mass radiography on a national scale, e.g. the financial involvements, the various delicate questions of an ethical or political character which may arise, and which will need the most careful and tactful handling if the scheme is not to be wrecked, the difficult question of the appropriate periods for re-examination of individuals, and the possible danger of creating phthisiophobia and general hypochondria. It is sufficient here to note that these points are realised by most of those who advocate mass radiography, and that they have been at least considered in some of the discussions that have already taken place, in the Royal Society of Medicine and elsewhere. In commending this number of our Journal to its readers, I would remind them in conclusion of the value of the long-term policy and of the necessity for a better appreciation of the vast amount of expense and suffering entailed by our present inadequate and wasteful method of dealing with pulmonary tuberculosis. This malady provides one of the best examples of the need for the medical profession to put its house into better order. We have at the present time a great opportunity for reaping the reward of all the valuable knowledge that has emerged in the last few years from the co-operative researches of chest physicians, radiologists, and pathologists in the natural history of this disease.