2. *First-Aid Burn Cream* (No. 9).

Cetyl Trimethyl Ammonium Bromide (Cetavlon)	1 per cent
Sulphanilamide	3 "
Castor Oil	25 "
Beeswax	1.8 "
Wool Fat	1.8 "
Cetyl Alcohol	5.0 "
Glycerine	10 "
Water to produce	100 "

THE CARE OF THE BURNED PATIENT

By R. P. OSBORNE, F.R.C.S.

(Plastic Surgery Unit, Stoke Mandeville Hospital, Aylesbury, Bucks.)

History

Tannic acid, for long the most popular method of treating burns, was introduced by Davidson (1925), and good support was given in England by Wilson (1929) and Mitchener (1938). Because tannic acid lacks any bactericidal action, Aldrich (1933) favoured gentian violet, and later the triple dye solution (1937).*

Since the War began in 1939, there have been many articles written (Mowlem 1941, Wells 1942, Erb et al 1943) relating to liver necrosis following the use of tannic acid. Present writings appear to favour—saline treatment (McIndoe 1940), plaster of Paris (Barnes and Trueta 1941; Fleet and Ackman 1944), the Koch method of pressure dressings (Allen and Koch 1942; Silver and Reid 1942), or the envelope method (Bunyan 1941, Douglas 1944).

Objects

Whichever method is chosen, the objects should be the same:—

- (1) To obtain a skin covering as soon as possible.
- (2) To maintain full function.●
- (3) To prevent deformity.
- (4) To prevent infection and cross infection.
- (5) To eliminate pain.
- (6) To reduce to a minimum the length of time between receipt of the injury, and return to normal duty.

* From the Plastic Surgery Unit, Lancashire County Hospital, Whiston.

Planning

There should be a definite plan of campaign in every hospital, large and small, understood and adopted by all who may be called upon to take part in the treatment of the case. By this means better results will be obtained and the value and comparative value of any method better assessed.

Ackmann (1944) and his colleagues have drawn up a plan which is a model of perfection. It is sad to note that a complacent attitude still exists in some hospitals: burns must not be regarded as minor problems and their treatment left in the hands of the enthusiastic but inexperienced junior residents.

First Aid

This will depend upon the materials to hand at the site of the accident, and will vary from applying the cleanest material available, to a sulphonamide preparation (Colebrook 1944) spread as a cream on gauze, covering this with wool and a firmly applied crêpe bandage. The general measures consist of warmth, warm sweetened drinks such as cocoa or tea, calm reassurance: the use of narcotics, depending upon the presence of a doctor and the absence of any other undiagnosed injury. Several cases of partial thickness burns in Service personnel have been treated since "D" Day, and the results of applying 5 per cent sulphanilamide cream on gauze, followed by bandaging, without any cleansing of the burned area, were very gratifying. Particularly so if the dressings had been renewed at least every 48 hours for, naturally, the cotton bandages used tended to become loose and the wound exposed, as movements of the individual, and passage of time occurred. None of the areas were free from bacterial contamination, all were clinically clean and very rapidly healed with 100 per cent function after the application of Stannard envelopes. It is agreed that they would have healed just as speedily by any of the other methods, but the advantage gained by using the envelopes was the immediate freedom of movement.

Definitive Therapy

(a) *Local—limbs*

Since reporting on the use of envelopes in the *British Journal of Surgery* (1943), the author has continued to use these whenever possible. Douglas (1944) has since issued a detailed account of this method as used in the U.S.A., and to him should go the credit for first introducing the procedure (1936 and 1939). Nothing further need be added to the detailed account already given except to

emphasize the necessity to elevate the limbs in the early stages, or whenever there is oedema if the case is seen late, and to apply pressure by means of wool and crêpe bandages outside the envelope between the irrigations. During the irrigations active movements are encouraged. The other instruction requiring emphasis is the use of splints outside the envelope, e.g. for deep burns of the hand—made of plaster with the hand, fingers and wrist in the position of function.

(b) Local—face

Envelopes cannot be used for burns of the face, and here the suggested procedure is to cleanse the affected area by very gentle use of gauze soaked in electrolytic sodium hypochlorite solution 1 in 100 at 100° F., or 1% C.T.A.B. (If the scalp is involved, it is advisable to shave the area and a wide margin around—otherwise infection is apt to follow, especially in women.) The area is dried by means of filtered air through a hair dryer, covered with a single layer of tulle gras; over this, gauze four layers thick soaked in the electrolytic sodium hypochlorite 1 in 100; a layer of wool and a firm crêpe bandage completes the dressing. In widespread burns only the nostrils and vermilion of the lips are left uncovered.

The after-treatment is to remove the bandage and wool twice a day and resoak the gauze with the same solution of electrolytic sodium hypochlorite 1 in 100 at 100° F. by means of an undine. The whole dressing is taken down and renewed every 48 hours. Healing occurs quicker than on other parts of the body in partial-thickness burns, and correspondingly, the areas are fit for grafting at an earlier date in cases of full-thickness loss.

(c) Local—Other Areas

Envelope therapy can be used for burns involving the thorax and upper abdomen or back, using a covering designed in the form of a smock; for burns involving wide areas a nightdress of the envelope material is available which is used in combination with a specially designed "irrigation bed" (Goldberg, 1944).

(d) General Measures

These begin the moment the patient is admitted to Hospital, treating the shock by means of warmth (not exceeding 70° F.), warm, sweetened drinks by mouth when possible, and giving a suitable drug to relieve the pain. Temperature and respiration are recorded four-hourly, and the pulse rate hourly. A fluid intake and output record is commenced. Plasma given intravenously is best started at once and continued during the process of cleansing and application of local covering—a

process which is not begun until the patient has made a good recovery from the shock.

Other general measures continue throughout the whole treatment of the patient and will be described under the appropriate headings.

The Circulatory System

Blood pressure readings are insufficient to assess the condition of a burned patient (Olson 1943). More reliance can be placed upon the haemoglobin, haematocrit, and plasma protein levels, together with the blood count. The more extensive the burn, the greater the need for repeated readings, for without them the general treatment becomes a haphazard procedure.

Plasma is lost from the capillaries and therefore plasma must be replaced—either empirically with the first bottle run in quickly, and a subsequent maintenance rate of 80–100 drops per minute, or based on calculations made from a formula (Harkins 1940, Black 1940, or Elkington 1940).

In major burns blood transfusions will be required after the first few days; in the severe cases at weekly intervals—particularly if there is infection. A difficulty then arises following upon the loss of suitable veins due to thrombosis: when all available have been used the transfusion can be given into bone marrow. A state of anaemia will delay the successful treatment of the infection and render the "take" of grafts less certain.

Infection

Although the area burnt is rendered sterile at the time of injury, it is only a matter of hours before bacterial contamination occurs. An opportunity of gauging the effect of sulphonilamide cream as a first-aid dressing in civilian life, and admitted to hospital within a few hours, has not yet occurred. Several cases were admitted from Normandy where this measure had been used, but all showed contamination.

When the patient is admitted to hospital one of the aims is to prevent further contamination, and reduce the possibility of cross-infection (Clowes 1943), two requirements which are well met using the envelope therapy (admittedly, with much less certainty when the irrigation bed is used). No matter what local application is used, whenever the area is exposed, all taking part, and the onlookers, should be gowned and masked—not forgetting a mask for the patient.

The organisms commonly found are *Staph. albus*, *Staph. aureus*, diptheroid bacilli, haemolytic and non-haemolytic streptococci, *B. Subtilis*, and unhappily *B. coli*, *B. proteus*, and *B. pyocyaneus*. Generally the contamination is mixed.

The methods of combating the bacteria and their effects are the sulphonamides, locally or by mouth, penicillin, local or systematic, propamidine, euflavine preparations, and zinc peroxide. When sulphonamides are used, the concentration level of the drug in the blood must be watched: high levels are to be expected if it is used locally over large areas. Altemeier (1944), Meleney (1944), and Whipple (1943) have recently suggested that sulphonamides are more effective by mouth than used locally. *B. coli*, *B. proteus*, or *B. pyocyaneus* are the most difficult to eliminate and neither sulphonamides nor penicillin are of value. Many local applications have been suggested, e.g. 10 % mercurochrome, household vinegar, 2 % acetic acid, 5 % gentian violet, potassium tellurite, proflavine with sulphathiazole, neutral proflavine sulphate, etc.

It is well-nigh impossible to keep a full thickness loss area free from bacteria, but all efforts to control the infection should be made, and this includes attempts to free the area of any dead tissue. Any continued infection leads to anaemia, exhaustion, loss of weight, fall of plasma proteins, loss of appetite, and all the usual signs of fever. At the same time there will be a delay in the appearance of granulations sufficiently healthy to permit skin grafting, and early skin grafting is the very factor one aims at in this depth of burn.

Hypo-proteinaemia

Loss of protein occurs early due to the exudation of plasma from the surface of the burn, and into the subjacent tissues, and to some extent this is replaced by the intravenous use of plasma. How much of the plasma given is retained is difficult to say, but to give saline would merely serve to increase the total loss. It is often stated that the first infusion of plasma should be given rapidly, and thereafter a steady drip, but the value of this is doubted by Lee and Wolff (1942).

Later the plasma proteins may be further depleted because of infection, loss of nitrogen in the urine, and the demands for tissue repair, and since an adequate protein reserve is necessary for wound-healing (Ravdin 1940), it is essential that efforts be made to replace the lost proteins.

Wolff and Lee (1942) have published a chart whereby, if the plasma protein level, haematocrit value, and body weight are known, the plasma protein deficit can be read off. The treatment of such a deficit is in the form of blood transfusions, concentrated serum infusion, high protein diet (e.g. eggs, milk, meat and brewers yeast). In cases where the surface area burnt is large, it may not be possible to replace the proteins in

the amount necessary by mouth; in this connection much work is being done on the intravenous use of amino acids, and food by means of a stomach tube (Taylor 1943). It is interesting to observe that workers in America suggest from their experiments that hypo-proteinaemia has some bearing in connection with the formation of bed-sores (Mulholland 1943).

Skin Grafting

Save in small areas in unimportant sites, full-thickness loss burns will require to be grafted. Ideally, these areas would be excised and grafted at the time of the initial cleansing if this was possible, for considerable time now wasted in waiting for the removal of dead tissue would be saved. The reason it is not carried out then, is the absence of certainty as to the exact extent of the full-thickness loss. Dingwall (1943) suggests giving 10 cc. of 20 % sodium fluorescein intravenously, and then examining the burned area in ultra violet light screened with a Woods filter. Under this light, areas of full-thickness loss are seen as sharply demarcated blue-black patches, in contrast to the surrounding yellow-green colour of normal fluorescein containing skin. As it is, the aims are to get the surface ready for grafting by removal of all dead tissue, reducing bacterial contamination as much as possible—especially the haemolytic streptococcus, and *B. pyocyaneus*, correcting any anaemia, and so producing flat, firm, bright red granulations (Barrett Brown 1943, and Padgett 1943). This preparation takes three to five weeks, and when these aims have been achieved, thin Thiersch grafts may be applied. However, when large areas are involved, and bacterial contamination is heavy with much discharge, time can be saved by using thin Thiersch grafts applied in the form of postage stamps. This type of graft is prepared as described by Gabarro (1943), but the author (1944) applies a Stannard envelope over the area grafted whenever possible, a method suggested by Professor T. P. Kilner. A pressure dressing of wet gauze, wool, and crêpe bandage is applied outside the envelope. On the third day this pressure dressing is removed, the envelope gently lifted off (it is *not* removed), the tulle gras covering the grafts and granulations, and an irrigation with electrolytic sodium hypochlorite (1 in 20) carried out, taking care not to direct the stream directly on to the grafts. Thereafter an irrigation is performed once daily until the seventh day, when the pressure dressings are discarded. Irrigations are then continued until all the spreading grafts coalesce, and the patient is encouraged to carry out a full range of movements during these irrigations. By this means the chance

of the grafts failing to take because of infection is vastly reduced, ample drainage being available. The combination of good drainage and moist dressing is emphasised by Padgett (1943) as a necessity when grafting granulated areas. This postage stamp method is of additional value when, because of the extent of the burn, the amount of donor area is limited.

A great advance in the treatment of full thickness burns will be made when the time between the injury and the application of a graft is reduced to a matter of days rather than weeks. Ackman and his colleagues (1944), using an occlusive pressure dressing over 5 % sulphathiazole emulsion can skin graft at the time of the first dressing on the 12th to 14th day (with this dressing streptococcus pyogenes *B. pyocyaneus*, and *B. proteus* have been absent). This is a highly creditable saving of time.

Early separation of the dead tissue in full-thickness burns is the most important local change; the earlier it can be achieved and grafting proceed, the less the drain on the patient's resistance. It is the presence over a long period of a large infected wound which is responsible for the deterioration of the general condition, and this is in turn responsible for the delay in improvement of the local condition. Therefore, one looks forward with considerable interest to the results of the experiments of Connor and Harvey (1944), who are using acids so prepared as to lower the pH of the surface of the wound. So far, their results suggest that a slough can be removed in 72 hours without damage to the underlying tissue, so that grafting can proceed at once. Should their further experiments confirm the present findings, a tremendous advance will have been made since, in addition to eliminating the damaging general effects of the open wound; the local dysfunction due to the formation and contracture of scar tissue will be reduced to a minimum.

After Treatment of a Grafted Area

Since thin grafts are applied in order to reduce the chance of "failure to take," the newly covered area is at first delicate. Whilst active movements are encouraged, friction against hard articles must be avoided in the early days: blister formation has been seen following the use of a billiard cue, table tennis bat, and garden spade, in enthusiastic Service patients.

Daily wax baths followed by massage with lanoline, and a further application of lanoline when retiring into bed for the night, are the best methods of improving the grafted area—keeping it smooth and supple, and, at the same time, it is getting thicker. The patient or parents, can

learn to apply the lanoline, which should be continued morning and evening for three months after discharge from hospital.

Rehabilitation exercises should begin as soon as possible when the patient is in bed and continued when he gets up, thus restoring his general fitness.

Homografts of Skin

Homografts of skin may be indicated when the burns are extensive and the patient's general condition poor, or when the amount of donor area is small as in children. The use of the grafts will serve to improve the patient's condition by limiting the exudation from the burnt surfaces. They will not survive for more than a few weeks, and by this time it is to be hoped that general measures have so improved the patient, that his own donor areas can be used. Fortunately, the need for homografts is not great, otherwise difficulties in obtaining donors might at some date be solved by the creation of "skin banks," in addition to the "blood banks" which are so invaluable.

Penicillin

When sulphanilamides either by mouth or used locally, fail to eliminate the streptococcus from a burn, the use of penicillin is of very great value. The elimination of this organism serves to reduce the effects of infection, and to render the take of grafts more certain. If the *Staph. aureus* is producing large quantities of pus which would tend to "lift off" grafts, again treatment with penicillin prior to grafting is to be advocated.

B. pyocyaneus, *B. proteus*, and *B. coli* are not eliminated by penicillin and, unfortunately, they can be responsible for the production of a great deal of pus.

In mixed infections penicillin will eliminate the streptococci and staphylococci, but the effects of *B. pyocyaneus*, *B. proteus* and *B. coli* may become more marked.

If penicillin is to be used it can be applied either locally as a powder or cream, or systemically. To use it as a fluid either for irrigation or in compresses would be sheer waste.

Bodenham (1943) has reported on the use of penicillin powder and cream in 75 cases of burns and wounds; and in the prophylactic use of the powder under grafts on non-infected surfaces in 15 cases. He recommends the use of a cream (100 Oxford units per gramme) for the treatment of burns.

Battle (1944) working in charge of a Maxillo-Facial Unit with the C.M.F., reports the use of the calcium salt in sulphathiazole or sulphanilamide powder (2,500 Oxford units per gramme) on 48

granulating areas due to burns in 33 cases. This powder in all cases was applied to the granulating area under the grafts at the time of grafting. Their preliminary preparation varied, however, and on this variation Battle placed them into three main groups.

In Group "A" a "blitz" technique was employed as soon as the separation of sloughs allowed; the patient was dressed on two or more consecutive days with the powder, vaseline gauze (open mesh), and dry gauze. Pressure with a crêpe bandage was universally applied. On the 3rd day the grafting operation was performed.

In Group "B." In this group dressings were done intermittently with penicillin powder. The treatment was spread over a long period with two or three days between dressings. Two of these patients were also the recipients of a full course of parenteral penicillin—Case 17 for a clinical septicaemia and Case 22 for pneumonia.

In Group "C." In this group of cases no penicillin was applied until the time of grafting. Case Nos. 16, 29, 30, 31, 33, and 25 all came into the theatre for their operation with their arms still encased in oiled silk of their Stannard envelopes. The granulations were cleansed with saline and the powder applied as elsewhere but only at the time of grafting.

The following table represents the result of the three groups:—

Percentage of Take Areas

	Total number Patients	Total number Areas Grafted	90 %	80-90 %	70-80 %	60-70 %
Group "A"	15	19	10	5	2	2
Group "B"	10	19	9	7	2	1
Group "C"	10	17	8	6	2	1
All Groups	35	55	27	18	6	4

In commenting upon these results Battle emphasises that in only four of these cases was the haemolytic streptococcus isolated prior to using penicillin, and that in the absence of controls, his figures are of little value in drawing conclusions.

Anaesthesia

The use of strong solutions of electrolytic sodium hypochlorite requires a general anaesthetic. We have never had to use an anaesthetic for the subsequent irrigations, except in an occasional case of severe burns in which other forms of treatment had been carried out elsewhere, and the patient had a low threshold to pain and had lost his confidence. Some patients do complain of tingling during, or

smarting pain, lasting for a few minutes after the irrigation, and if this is intolerable it can be met by reducing the strength of the solution—if necessary to 1 % electrolytic sodium hypochlorite. Late cases have complained of severe pain during the irrigation and for these we have used the Minnitt Gas and Air machine with complete success. If necessary this method is fortified by the addition of a little trilene, either by the addition of a Rowbotham's bottle, or suspending a piece of gauze by means of adhesive tape over the air inlet and dropping trilene on to this as required. As the general condition improves, the necessity for an anaesthetic disappears.

For the initial dressing or for the skin grafting operation we use cyclopropane for all patients over two years of age; below this age gas and oxygen suffices.

Gordon (1943) uses intravenous novocaine for painful dressings, but we have not yet given this a trial when carrying out an irrigation.

Scar Contracture

Preventative treatment lies in the prevention of oedema, overcoming infection, and grafting as early as possible. In some deep burns of the dorsum of the hand, no method of local treatment will prevent deformity, and in these cases the hand and fingers must be splinted in the position of optimum function. Ectropion is a particularly dangerous result of contracture because of the risk of permanent damage to the cornea, and must have priority when grafting is performed.

The correction of other contractures can usually be performed by excision of the scar and the application of a free graft with the area splinted if possible in an over-corrected position. After the graft has taken, massage with lanoline should be continued and the splints maintained at night for three months.

Keloids

The cause of this complication is unknown, although infection may play some part. It occurs at the junction of graft to normal skin, at the junction of postage stamp grafts to each other, or when areas have been allowed to heal without grafting. Treatment is called for when the irritation and resultant scratching are severe or for cosmetic purposes. In both cases the X-ray treatment recommended by Levitt and Gillies is of great value.

Summary:

1. Burns do not belong to the realm of minor surgery.

2. The tannic acid era has passed.
3. The best First Aid treatment is—sulphanilamide cream, together with a carefully applied pressure dressing.
4. The prominent methods of definitive local therapy are—pressure dressings, saline baths, and the envelope therapy.
5. Great and continued attention must be paid to the general treatment if more lives are to be saved and better results from local measures achieved.
6. Considerable advances have yet to be made in eliminating *B. pyocyaneus*, *B. proteus*, and *B. coli*.
7. Better results are to be expected with improved methods for restoring lost plasma proteins.
8. Promises of great value are made in the experimental work on the use of acids which will remove sloughs in a matter of hours instead of weeks, thus rendering early skin grafting possible.
9. Early skin grafting and skilled attention to the general condition are the secrets of success in the treatment of full-thickness burns.

Illustrative Cases

Case I (L.J., age 3).

Admitted 5th November 1944. The child's nightdress caught fire on 21st October 1944, as a result of which she received full-thickness burns of the right side of the neck, extending over the shoulder to the right wrist, involving a large area of the right chest wall, the right buttock, and a large area of the lateral side of the right thigh. (Estimate, by Berkow (1924) method, was 27 %.)

She was admitted to her local hospital the same day and was given one pint of plasma and two of saline. After a few hours the areas were cleansed under gas, oxygen and ether anaesthesia. Propamide, tulle gras, and bandages were applied as a dressing. This dressing was repeated three times, under a similar anaesthetic in fourteen days. The one smear taken showed the presence of *B. proteus*.

16th Day (T. 101°, P. 130, R. 30). The day following the transfer, all the areas were irrigated under gas, oxygen, and a little ether, with electrolytic sodium hypochlorite (1 in 20): a few sloughs separated, but most were densely adherent. The limbs were oedematous and all areas infected. After drying, 5 % sulphanilamide cream on gauze, wool, and a crêpe bandage were applied. A posterior plaster shell was made. General condition poor.

Blood:

Erythrocytes, 3,580,000 per cu.mm.	Neutrophils 70 %
Haemoglobin, 69 % (10.76 gms.%)	Eosinophils —
Colour Index, 0.9	Basophils —
Halometer, 7.2 u.	Lymphocytes 28 %
Leucocytes, 23,125 per cu.mm.	Monocytes 2 %
Volume of packed cells, 24 mm.	
Mean corpuscular volume, 68 cu.	
Mean corpuscular haemoglobin concentration, 44 %	

Urine:

Albumen absent
Urea 0.7 gm.
An occasional leucocyte present
Rbc's crystals absent
No growth on culture

Bacteriology (all areas):

Pus cells and gram negative bacilli
Culture overgrown by *B. proteus*

17th Day (T. 98.4°, P. 130, R. 28). Plasma protein = 5.7 gms. per 100 m.l. plasma.

18th Day (T. 99°, P. 140, R. 28). Redressed as on the 16th day; blood transfusion started, one pint given at slow drip, followed by half a pint of plasma.

19th Day (T. 101°, P. 120, R. 28). General condition very poor; cyanosis; drowsy.

20th Day (T. 99°, P. 140, R. 28). Has gradually taken better diet since admission; eggs, milk, oranges, marmite.

21st Day (T. 99°, P. 140, R. 28). Cyanosis less marked; plasma protein 5.04 gms. per 100 m.l. plasma; dressing under cyclopropane; all granulations much cleaner; most of slough separated; irrigation as before, but zinc peroxide used instead of sulphanilamide because of cyanosis. Concentrated serum in 200 c.c. water given by slow drip (4 hours).

22nd Day (T. 100°, P. 130, R. 26). Pulse volume stronger than for last few days; taking diet well; looks less toxic. Plasma protein 7 gms. per 100 m.l. plasma. Albumen 3.5 per 100 m.l. plasma. Globulin 3.11 per 100 m.l. plasma. Fibrin 0.39 per 100 m.l. plasma.

24th Day (T. 98.4°, P. 130, R. 26). All smears showed few pus cells but no organisms. Under pentothal, postage stamp Thiersch grafts from left thigh applied to right arm, neck and thorax. Similar grafts from inner aspect right thigh applied to outer aspect right thigh. Stannard envelopes to arm and leg over grafts. Acriflavine emulsion wool to neck and thorax. Pressure dressing of wool and crêpe bandages to all areas. Child placed in posterior plaster shell with 90° abduction right shoulder. Operation time 1½ hours. General condition very poor.

25th Day (T. 101°, P. 160, R. 30). A very poor day.

26th Day. Slowly improving.

27th Day (T. 99°, P. 140, R. 24).

Blood: Erythrocytes 3,210,000
Haemoglobin 69 %
Colour Index 1.0

Under cyclopropane both envelopes irrigated—grafts *in situ*—granulations healthy pink. Chest—grafts *in situ* and granulations pink; irrigation: compresses of electrolytic sodium hypochlorite 1 in 100 applied. Neck—owing to movements of the child 25 % of grafts lost; dressed as for thorax. Pressure dressings to all areas.

28th–34th Day (T. average 101°, P. 140, R. 24). General condition improving. Taking high protein diet well. Dressings every other day under cyclopropane. Very slow spread of grafts.

Blood:

Erythrocytes 2,350,000 per cu.mm.	Neutrophils 48 %
Haemoglobin 49 % (5.67 gms. %)	Eosinophils 1 %
Colour Index 1.07	Basophils —
Halometer 7.5 u.	Lymphocytes 47 %
Leucocytes 10,937 per cu.mm.	Monocytes 4 %
Volume of packed cells 25 mm.	
Mean corpuscular volume 92 cu.	
Mean corpuscular haemoglobin concentration 23 %	
Total protein 6.5 gms. per 100 m.l. of plasma	
albumen 3.18	" "
globulin 2.9	" "
fibrin 0.42	" "

Bacteriology:

Staphylococci overgrown by *B. proteus* in all areas.

35th Day. Blood transfusion—one pint in 5½ hours. Had a rigor during transfusion.

ILLUSTRATIONS ON BURNS

By

R. P. OSBORNE,
F.R.C.S.

CASE I

1.—Before grafting. 21st day following injury.

2.—After grafting. 90th day following injury.

CASE II

1.—Showing limbs in Stannard envelope

2.—Lower limbs in 6th week, showing spread of postage stamp grafts.

CASE IV

1.—Before operation with eyes closed.

2.—Before operation with eyes open.

3.—After operation with eyes closed.

4.—After operation with eyes open.

CASE V

1 and 2.—Before operation.

3 and 4.—After operation.

CASE V

5.—After operation.

1.—Before operation.

6.—After operation.

2.—After operation.

38th Day. Dressing under cyclopropane. Excellent spread of grafts on arm and leg; new envelopes applied; very slow spread of grafts on chest and neck. Small pressure sore on scalp noticed on 33rd day, has suddenly spread and is now 4 in. $\times$ by 2 in. with bone exposed.

From then onwards rapid progress was made; the leg was healed on the 43rd day, and the arm on the 49th—the envelopes being removed on these days. The temperature came down to 99° and the child was much brighter, much easier to nurse, and was eating extremely well.

On the 45th day more postage stamp Thiersch grafts taken from the left upper arm, were applied to the neck and chest, as some of these had failed in the former and were rather sparse in the latter, but those present had spread very well.

From the 52nd day she had irrigations daily without any anaesthetic, and did not complain of any pain.

Blood:

Erythrocytes 3,970,000 per cu.mm.

Haemoglobin 70 %

Colour Index 0.9

Leucocytes 7,812 per cu.mm.

On the 58th day massage with lanoline to all the healed areas and exercises were begun. From then onwards she made steady progress; all areas healing with the exception of a small obstinate area on the tip of the shoulder. She remained an in-patient on the 100th day because her home was situated too far away from any hospital where she could receive the necessary physiotherapy. There is 100 % function of the leg, abduction of the shoulder to 90°, full function of the elbow, and some tendency to contracture of the neck scar.

Comment: An extreme full-thickness loss in a very young patient who responded well to general measures, though many anxious days were experienced. A striking improvement was noticed following the rapid spread of grafts. The reappearance of organisms within the envelopes is explained by the fact that these could not be rendered occlusive at their upper extremities due to the extent of the burns on to the shoulder and buttock. The organisms did not interfere with the spread of the grafts. The pressure sore was a disappointment, for she had excellent nursing attention.

The child's life was saved by the general measures taken.

Case II (E.S., aged 26).

Admitted 17th July, 1944. A Service patient burnt during a tank battle in Italy seven weeks previously, as a result of which he received partial-thickness burns of the whole face and neck, full-thickness burns of the helices, deep partial-thickness burns of most of both forearms, dorsum of hands and fingers, scattered partial-thickness anterior chest wall, full-thickness of both legs from the junction of upper and middle third of the thigh to the junction of middle and lower third lower leg, and full-thickness both heels. (Berkow Chart = 36 %.)

In Italy he received eight pints of plasma, two of concentrated serum, and two of glucose saline within the first week. The local dressings consisted of saline compresses, renewed twice daily, supplemented later by penicillin powder. Sloughs began to separate during the second week, and in the fifth week the limbs were placed in plaster. These plaster dressings were renewed at intervals, and finally removed during the voyage to England because of a profuse discharge of offensive greenish pus and intense pain in the legs.

On admission he was terrified of having any further change of dressing because of the pain he had always had in the past. He was thin, toxic looking, without an

appetite, very thirsty, and would not attempt to move any of the limbs, which he held in extension. Under anaesthesia (pentothal), the granulations were found to be pale, flabby, and exuberant, discharging a copious quantity of greenish pus; the hands and lower limbs were oedematous. A Stannard envelope was applied to the right forearm and hand, with a plaster splint and pressure dressing externally.

5 % sulphanilamide cream and pressure dressings were applied to the lower limbs and ears.

Blood: Erythrocytes 2,850,000 Haemoglobin 50%

Bacteriology:

B. pyocyaneus, *B. proteus* and *B. coli* in all areas.

The right upper limb was irrigated twice daily and was healed six weeks later, with some flexure contracture of the m.p. joints, but with very useful range of movement.

The lower limbs presented a problem because of the continued persistence of the organisms found on admission, and the large amount of pus produced. Homografts were applied as postage stamps to the right leg five days after admission, and these took; thereafter both legs were irrigated every 48 hours in an irrigation bed in which he remained, but so low was his morale that gas and air analgesia (Minnitt apparatus), sometimes supplemented with trilene, had to be used.

Postage stamp grafts taken from the left upper arm were applied to the left leg in the third week, and to the right leg from the right upper arm in the fifth week, both limbs being placed in Stannard envelopes as described by the author (1944). Blood transfusions were given at each operation. An excellent "take" followed except on the posterior aspect of the left lower leg. The bacteriology remained unaltered, and the discharge of pus persisted, but less profuse.

From the fourth week onwards he could tolerate irrigations without anaesthesia, but he would not co-operate with the physiotherapist except in the upper limbs.

He was transferred to another hospital twelve weeks after admission, using both hands very well, 45° flexion of the hips, but only 10° at the knee joints. Some oedema persisted in the right leg. He had gained weight, and was altogether a different man.

Comment: After seven weeks' treatment abroad in the difficult conditions of warfare, oedema, infection, anaemia, and a low morale existed, all of which delayed grafting on reaching England, and placed a great strain on the nursing staff. Plasma protein remained low throughout (between 5 and 6 gms. %). In view of the "take" of homografts it is clear that a more rapid improvement would have resulted if far more of these had been applied. Lack of a dermatone severely restricted the available donor areas, for more of his own skin could have been used despite the continued presence of the infection.

Case III (M.B., aged 32).

Admitted 23rd June 1944, five hours after receiving extensive "blistering" of the left thigh, following scalds by hot water. No first aid measures had been used.

All areas cleansed with electrolytic sodium hypochlorite (1 in 5), under gas and oxygen anaesthesia, followed by the application of a Stannard envelope. Subsequently irrigated twice daily with 1 in 20 solution, and movements encouraged. Envelope removed on fifteenth day with complete healing and 100 % function.

Bacteriology: *Staph. albus* on admission. No growth on sixth day.

Comment: An excellent result because of the early treatment of unbroken blisters, and continued exercises throughout treatment.

Case IV (J.T., aged 47).

Admitted to Stoke Mandeville Hospital 24th July 1942. In January 1942, during an epileptic fit, he fell into the kitchen fire and sustained full-thickness burns of the right cheek, lower eyelid, and deep partial-thickness burns of the upper eyelid and right side of the forehead. He was treated by local dressings (details unknown), and allowed to heal by scar tissue formation with resultant severe ectropion of the right lower eyelid and a marked ectropion of the lower lip. There was a marked right corneal opacity.

The ectropion was corrected by freeing the lower eyelid into an over-corrected position, followed by the application of a Thiersch graft. Further work on the lips and cheeks was postponed for domestic reasons but could not be carried out because death followed a fall into his bath during another fit.

Comment: Corneal opacity resulting from neglected ectropion—a condition which should be prevented by grafting at the earliest possible opportunity.

Case V (T.B., aged 6).

Admitted to Stoke Mandeville Hospital 19th January, 1943. On 7th August, 1942, whilst playing with matches, his clothing caught fire as a result of which he received extensive burns of both upper limbs, right side of face and neck, and the right chest wall. All areas were treated at his local hospital with gentian violet and he was discharged five months later with the severe scar contractures seen in the illustrations.

Of these the neck scar received primary attention because if left it would interfere with the development of the mandible. These scars were divided, all deep scar tissue removed, and the wound opened out to the fullest extent. Free grafts were applied, and the patient placed in a previously prepared posterior plaster shell.

Later the flexure contracture of the elbow and adduction contracture of the axilla were corrected in the same manner, the limb being placed vertically above the head by means of plaster splints.

All these grafts took well, resulting in a full recovery of function in all joints, except for a few degrees limitation of extension of the neck.

Comment: An example of the correction of scar contracture by free grafts. These burns known to be deep at the time of injury should have been treated by grafting within two or three weeks during the original treatment; the contractures could have been foreseen and prevented.

I am deeply indebted to all the members of my team at the County Hospital, Whiston, and particularly to Professor T. Pomfret Kilner for his constant encouragement and helpful criticism.

REFERENCES

- ACKMAN, D. et al (1944), *Annals of Surgery*, **119**, 161.
 ALDRICH, R. H. (1933), *New Eng. Journ. Med.*, **208**, 299.
 ALDRICH, R. H. (1937), *New Eng. Journ. Med.*, **217**, 911.
 ALLEN, H. S. and KOCH, S. L. (1942), *Surg. Gynae and Obstet.*, **74**, 914.
 BATTLE, R. J. V. (1944), Personal Communication.
 BARNES AND TRUETA (1941), *Lancet*, **1**, 623.
 BERKOW, S. E. (1924), *Arch. Surg.*, **8**, 138.
 BLACK, D. A. K. (1940), *Brit. Med. Journ.*, **2**, 693.
 BODENHAM, D. C. (1943), *Lancet*, **2**, 725.
 BROWN, J. B. and McDOWELL, F. (1943), *Skin Grafting of Burns* (J. B. Lippincott).
 BUNYAN, J. (1940), *Proc. Roy. Soc. Med.*, **34**, 23.
 BUNYAN, J. (1941), *Brit. Med. Journ.*, **2**, 1.
 CLOWES, G. H. A. et al (1943), *Annals of Surgery*, **118**, 761.
 COLEBROOK, L. (1944), *Brit. Med. Journ.*, **1**, 342.
 CONNOR, G. J. and HARVEY, S. C. (1944), *Annals of Surgery*, **120**, 362.
 DAVIDSON, E. C. (1925), *Surg. Gynae and Obstet.*, **41**, 202.
 DINGWALL, J. A. (1943), *Annals of Surgery*, **118**, 427.
 DOUGLAS, B. (1936), *Journ. Tennessee M.A.*, **29**, 160.
 DOUGLAS, B. (1939), *Amer. Journ. Surg.*, **43**, 2.
 DOUGLAS, B. (1939), *South Med. Journ.*, **32**, 12.
 DOUGLAS, B. (1944), *Surgery*, **15**, 96.

- ELKINTON, J. R., WOLF, W. A., and LEE, W. E. (1940), *Annals of Surgery*, **112**, 150.
 ERB, I. H. et al (1943), *Annals of Surgery*, **117**, 234.
 GABARRO, P. (1943), *Brit. Med. Journ.*, **1**, 723.
 GOLDBERG, H. M. (1944), *Lancet*, **1**, 371.
 GORDON, R. A. (1943), *Canad. M. Ass. J.*, **49**, 478.
 FLEET, G. A. and ACKMAN, F. D. (1944), *Canad. M. Ass. J.*, **2**, 109.
 HARKINS, H. N. et al (1940), *Surg. Gynae and Obstet.*, **4**, 410.
 HANNAY, J. W. (1941), *Brit. Med. Journ.*, **2**, 4.
 HUDSON, R. V. (1941), *Brit. Med. Journ.*, **2**, 7.
 LEE, W. E., WOLFF, W. A. et al (1942), *Annals of Surgery*, **115**, 1131.
 LEVENSON, S. M., and LUND, C. C. (1943), *Journ. Amer. Med. Assoc.*, **5**, 272.
 LEVITT, W. M., and GILLIES (1942), *Lancet*, **1**, 440.
 McINDOE, A. H. (1940), *Proc. Roy. Soc. Med.*, **34**, 56.
 MITCHENER, P. H. (1933), *Brit. Med. Journ.*, **1**, 447.
 MOWLEM, R. (1941), *Proc. Roy. Soc. Med.*, **34**, 221.
 MULHOLLAND, J. H. et al (1943), *Annals of Surgery*, **118**, 1015.
 OLSON, W. H., and NECHELES (1943), *Amer. Journ. Physiol.*, **139**, 574.
 OSBORNE, R. P. (1944), *Brit. Journ. Surg.*, **125**, 24.
 PADGETT, E. C. (1942), *Skin Grafting* (Baillière, Tindall and Cox).
 PEARSON, B. P. et al (1941), *Brit. Med. Journ.*, **2**, 41.
 RAVDIN, I. S. (1940), *Annals of Surgery*, **112**, 576.
 SILER, V. E., and REID, M. R. (1942), *Annals of Surgery*, **115**, 1106.
 TAYLOR, F. H. L. et al (1943), *Annals of Surgery*, **118**, 215.
 WELLS, D. B., et al (1942), *New Eng. Journ. Med.*, **26**, 629.
 WILSON, W. C. (1929), *Med. Research Council*, No. 141 (H.M. Stationery Office).
 WHIPPLE, A. O. (1943), *Annals of Surgery*, **118**, 187.
 WOLFF, W. A., and LEE, W. E. (1942), *Annals of Surgery*, **115**, 1125.

FRACTURES—PART I PRINCIPLES OF DIAGNOSIS AND TREATMENT

by E. H. HAMBLY, F.R.C.S.

(Surgeon E.M.S., Royal National Orthopaedic Hospital, W.I)

The knowledge and treatment of fractures has progressed within the last decade almost beyond recognition in certain branches of the subject. Side by side with this development on the medical side there is growing in the minds of laymen and patients alike a greater knowledge of the treatment and attainable results. On the other hand, the clinician is greatly helped by the ever-growing variety of radiological and therapeutic methods available.

PRINCIPLES OF CLINICAL DIAGNOSIS

The patient complains of pain, swelling, and tenderness in the neighbourhood of a particular bone or joint.

He frequently gives a clear history of an injury, which he usually describes as a sprain. The history may be misleading, inasmuch that, in an old fracture, the injury may have been forgotten. It is only too easy to accept the patient's history of a sprain at its face value. *Every sprain should be assumed to be a fracture until proved to the contrary by X-rays.*

On examination of the patient the following points should be particularly noted. A general examination should always be made to exclude