ductus in *all* patients treated by ligation has not yet been achieved, and division between clamps may supersede simple ligation.

5. The effects and results of operation are reviewed.
6. Illustrative cases are recorded.

REFERENCES

- ABBOT, MAUDE E. (1936), *Atlas of Congenital Heart Disease*, New York: American Heart Association.
- BARCLAY, A. E., BARCROFT, J., BARRON, D. H., and FRANKLIN, K. J. (1942), *Am. J. Roentgenol.* **47**, 678.
- BOURNE, G., KEELE, K. D., and TUBBS, O. S. (1941), *Lancet* **2**, 444.
- BULLOCK, L. T., JONES, J. C., and DOLLEY, F. S. (1939), *J. Pediat.* **15**, 786.
- EPPINGER, E. C., and BURWELL, C. S. (1940), *J. Amer. Med. Ass.* **115**, 1262.
- GRABIEL, A., STRIEDER, J. W., and BOYER, N. H. (1938), *Amer. Heart J.* **15**, 621.
- GROSS, R. E., and HUBBARD, J. P. (1939), *J. Amer. Med. Ass.* **112**, 729.
- GROSS, R. E. (1944), *Surg. Gynec. and Obst.* **78**, 36.
- HARRINGTON, S. W. (1943), *Proc. Staff Meet. Mayo Clinic.* **18**, 217.
- KENNEDY, J. A., and CLARK, S. L. (1941), *Anat. Record* **79**, 349.
- MUNRO, J. C. (1907), *Ann. Surg.* **46**, 335.
- SHAPIRO, M. J., and KEYS, A. (1941), *New Internat. Clin.* **4**, 148.
- SHAPIRO, M. J., and KEYS, A. (1943), *Am. J. M. Sc.* **206**, 174.
- TOUROFF, A. S. W., and VESSELL, H. (1940), *J. Thoracic Surg.* **10**, 59.
- TOUROFF, A. S. W. (1942), *Am. Heart J.* **23**, 847.
- TUBBS, O. S. (1944), *Brit. J. Surg.* **32**, 1.
- WILSON, M. G., and LUBSCHEZ, R. (1942), *J. Pediat.* **21**, 23.

FRACTURES II

THE UPPER LIMB—DIAGNOSIS AND TREATMENT

By E. H. HAMBLY, F.R.C.S.

(Surgeon E.M.S., Royal National Orthopaedic Hospital, W.I)

It is rarely emphasised in writing that the treatment of fractures generally is very difficult. Nowhere does this apply more than to the common fractures occurring in the upper limb. For example, it is extremely difficult to obtain absolute anatomical and functional correction in many fractures of the elbow and the wrist in adults. Writers only too frequently suggest that such fractures can be easily reduced and full correction obtained. Such statements only depress the reader, who may be having considerable difficulties in his treatment of these fractures. It is only fair to indicate which fractures are difficult to treat and to indicate methods by which the treatment of these difficulties can be rendered more effective.

Nevertheless, the general standard of results still obtained from such fractures as those of the elbow, forearm, and wrist in adults is far too low. This standard can be raised by greater care in treatment, more attention to detail, a desire to obtain better anatomical and functional results

than heretofore, and better after-treatment to the soft tissues and to the joints not immobilised.

Radiographic Diagnosis

More and more the finer points of the diagnosis of fractures rest upon the methods of radiographic procedures. It is still true that clinical examination is pre-eminent, but only too frequently the only clinical sign is tenderness at the site of the fracture.

In injuries to the limbs the whole of a long bone should be X-rayed. Also the joints above or below the fracture must be included in the film. X-rays of both limbs should be taken for comparison. Especially is this necessary in the case of joint injuries. Furthermore, X-rays must be taken in at least two planes at right angles, and preferably in a third oblique plane.

Fractures of the Clavicle

Fractures of the clavicle occur most commonly in the middle and towards the outer extremity. The chief points concerning treatment are the avoidance of stiff fingers and a stiff shoulder. The usual method of treatment by three handkerchiefs or figure of eight bandage is very painful if applied properly, or ineffective if applied comfortably. The reader has only to permit himself to be tied tightly for one night with three handkerchiefs to appreciate this point.

The most efficient and most comfortable method is to place a four-inch square of adhesive felt on the fracture site and a similar square of adhesive felt on the under-surface of the olecranon. Elastoplast is then applied from the axilla on the inner side of the arm, round the under surface of the elbow, up the outer aspect of the arm over the clavicular felt pad, and across the trapezius to the back. Two pieces of elastoplast are so fixed with one circular piece loosely around the arm. This enables the patient to move the shoulder, elbow, and hand from the beginning.

Sayre's method is absolutely contra-indicated as it prevents active movements of all the joints.

Fractures of the Scapula

The scapula may be fractured in the body, neck, coracoid process, or acromium. Although frequently badly splintered by multiple fissure fractures the correct treatment for all these fractures is radiant heat and active exercises from the first day of injury.

Passive exercises should never be given in injuries of the limbs, in fact the writer is convinced that it has no place in bone and joint surgery. The only indication for passive exercises is paralysis associated with a nerve lesion.

Only rarely do severe fractures of the neck of the scapula require traction in abduction for about four weeks.

Fractures of the Humerus

Fractures of the Neck.

Over 95 per cent of fractures of the neck of the humerus are very easy to treat. The remainder however, are the reverse. Such a one is seen in X-ray No. 1.

Fractures of the humeral neck are sub-divided into three groups, namely, crack fractures due to contusion, adduction fractures, which are impacted, and abduction fractures, which may or may not be impacted.

The only treatment necessary for contusion fractures is radiant heat and active exercises daily from the first day. The sling is abandoned after a week.

Adduction fractures are always impacted. The best treatment again is radiant heat and active exercises daily from the day of the injury. Only when the shaft is very adducted on the head is traction in abduction of a frame necessary for four weeks. Even then the deformity frequently fails to correct itself. Nevertheless, functional results are excellent. See X-ray No. 2.

The treatment of abduction fractures depends on whether the shaft is impacted or not. If impacted, treatment consists of radiant heat and active exercises and a sling for one week.

Occasionally, an impacted abduction fracture occurs where the head is dislocated. Again, in the writer's opinion, conservative treatment is preferable after manipulation has been performed.

Open reduction of such a fracture-dislocation only tends to cause aseptic necrosis of the head. If late osteo-arthritis is severe an arthrodesis is indicated. It is incredible what good functional results can be obtained in these cases if radiant heat and active exercises are commenced immediately after the injury, even in the presence of shocking X-rays.

If abduction fractures are un-impacted a serious dislocation results. In these cases an abduction frame clearly makes the deformity worse, unless continuous screw traction is applied. If such traction is applied it must be fixed to strapping on the arm and not to a pin through the olecranon. This causes effusion and stiffness of the elbow joint. The best treatment for such un-impacted fracture-dislocations in abduction is manipulation of the humerus across the chest, at the same time applying pressure outwards. The limb is then placed in a collar and cuff sling for three weeks. Radiant heat and such

active exercises as the sling permits are commenced at once.

Dislocation of the Shoulder

All dislocations of the shoulder, as indeed all suspected injuries to bones and joints, should be X-rayed before reduction to exclude an associated fracture of the humeral head.

Dislocations of the humeral head are usually sub-coracoid. Other positions include sub-glenoid, luxatio erecta, sub-spinous, and between the clavicle and the first rib. The last two are rare.

A deep anaesthetic is essential, e.g. Pentothal

FIG.3.—Fracture of lower one-third of shaft of humerus which cannot be immobilised without an onlay graft.

or Ether. Gas should never be used for reducing fractures and dislocations. Complete relaxation is essential. Reduction is carried out by external rotation of the arm with the elbow by the side. Reduction frequently occurs during this first movement of manipulation. The second movement consists of bringing the elbow across the chest. Finally, the third movement internally rotates the arm.

Five per cent of all dislocations of the shoulder cause injury to the circumflex nerve, with deltoid paralysis. The arm should be held in abduction in a frame until the paralysis recovers. If the paralysis is only partial active exercises are sufficient.

Fracture-Dislocation of the Shoulder

In these cases the head of the humerus may be lying free. It may even be upside down. See X-Ray No. 1. These are very difficult cases. Manipulation should be undertaken with the arm in traction and in abduction or hyper-abduction. Open reduction is contra-indicated on account of aseptic necrosis of the humeral head. The best treatment if the humeral head is still upside down after careful manipulations is to ignore the X-rays and to commence radiant heat and active exercises at once. If osteo-arthritis becomes severe an arthrodesis can be performed.

Fractures of Shaft of Humerus

Fortunately, all fractures of the shaft of the humerus unite well and rapidly with one exception. The latter occur at the junction of the third quarter and the lowest quarter of the humerus (Fig. 3). Here immobilisation without fixation is impossible.

Treatment of fractures of the humeral shaft consists in manipulation and application of a posterior plaster slab from the shoulder to the knuckles. Gravity maintains the shaft in correct alignment provided the wrist is held with a collar and cuff or ordinary sling. Active exercises at the shoulder joint are undertaken from the beginning.

For fractures of the lowest quarter of the shaft see X-ray No. 4, the only treatment, as the writer has found from bitter experience, is an onlay tibial bone graft, which is fixed with four vitallium screws. It is important in this connection to have screws of sufficient length to lodge firmly in the compact bone on the other side of the shaft to the graft. The humerus is then immobilised in a shoulder spica for at least four months. The bone graft is most brittle between the seventh and eleventh weeks.

Fractures of Lower End of Humerus

The two main principles of treatment of injuries to the elbow joint are avoidance of all forcing, stretching, or manipulation, and to avoid the insertion of all metallic foreign bodies.

Supra-Condylar Fractures.

Supra-condylar fractures are extremely common. They are of two very distinct types. The first and commoner consists of *backward displacement* of the lower fragment. See X-ray No 5. This type should be treated by manipulation and fixation in *flexion*. This is done by the application of a posterior plaster slab and a wet cotton bandage. A complete round and round plaster is extremely dangerous. Care should be taken to observe if the radial pulse is still present. Absence of the pulse

after manipulation does not mean Volkmann's ischaemic contracture. It only means that the elbow is too flexed. The writer recently has seen three such cases in one month. The correct treatment when the pulse is absent at the end of manipulation is to extend the elbow to 20 degrees beyond the right angle, and to apply another plaster back slab in that position with an ordinary sling. The clinical sign of true Volkmann's Contracture is complete inability to extend the fingers without severe pain. Treatment for Volkmann's Contracture is to extend the elbow to 20 degrees beyond the right angle. If finger movements do not return after six hours the brachial artery should be explored in the ante-cubital fossa. If the artery is actually damaged, an inch of it should be excised. The writer believes that this operation is never necessary if the elbow is extended to 20 degrees beyond the right angle as soon as the radial pulse is absent, or as soon as extension of the fingers is absent. Clearly, all supra-condylar fractures should be admitted after reduction for observation.

The second type of supra-condylar fracture is that in which there is an *anterior displacement* of the lower fragment. These should be treated by manipulation and fixation with a posterior slab in *full extension*. See X-ray No. 6.

Fracture-Dislocation of Capitellum and Half Trochlea

In adults this is not uncommon. Nearly the whole of the articular surface of the humerus is completely displaced upwards in front of the humerus. See X-ray No. 7. This is a very difficult fracture to treat successfully. The writer has tried open reduction with repair with catgut, attempted closed reduction by extension, and excision of the fragment. All these methods have left a relatively stiff elbow. The best treatment is manipulation and application of a plaster-back slab in full extension. This is left for three weeks. The elbow is then flexed under anaesthesia and a plaster-back slab applied with the elbow at right angles.

Inter-Condylar and Y-shaped Fractures of Humerus

These types of fracture are also very difficult to treat satisfactorily when displacement is marked. Although many forms of treatment have been recommended, the writer favours manipulation and plaster with the arm in full extension for three weeks, after which time it is fixed at right angles under anaesthesia in a back slab. Immediate arthroplasty has been advised, but this gives mobility at the expense of strength.

Anterior Displacement of Whole Epiphysis in Children

This type of fracture should be treated similarly to an anterior type of supra-condylar fracture. The elbow is manipulated and immobilised with a plaster-back slab in full extension.

Displacement of External Condylar Epiphysis

The external condylar epiphysis may be displaced and rotated 90 degrees in both vertical and horizontal axes. This results in cubitus valgus and ulnar paralysis. Treatment consists in manipulation or fixation by open reduction. See X-ray No. 8.

Displacement of Internal Epicondyle

This injury is not rare. The ulnar nerve may be carried with the displaced fragment into the elbow joint. If the detached fragment lies within the joint space open reduction is necessary. The ulnar nerve should always be transplanted at the same time. See X-ray No. 9.

Fractures of the Radius and Ulna

Fracture of Head of the Radius.

Fissure fractures of the radial head only require a sling for four weeks. Active movements are undertaken at once. If fracture of the radial head is suspected clinically it should be treated accordingly, even if the standard three X-ray views, i.e. antero-posterior, lateral, and oblique, are negative.

If there is displacement of the articular surface of a portion of the radial head, the latter should be excised as soon as the arm can be suitably prepared. Delayed excision does not give good results. The radial head should not be excised in children and adolescents on account of Madelung's manus valgus of the wrist developing. The writer's method of excising the head of the radius is by means of the Gigli's saw, after turning back the periosteum. This causes less crushing and late "myositis ossificans" than other methods.

Displacement of Radial Head in Children

In children the whole head of the radius may be displaced laterally. This should be reduced by manipulation and direct pressure on the head. Excision of the head of the radius in children is a disaster. The whole radius shifts upwards, and Madelung's deformity results.

Dislocation of the Elbow

The elbow may dislocate backwards with a fracture of the coronoid process, or forwards with a fracture of the olecranon. Backward disloca-

tions are reduced under deep anaesthesia, and a posterior slab applied with the elbow at right angles. This is removed after four weeks, and active exercises *only* commenced. Anterior dislocations are treated by manipulation and fixation in full extension for six weeks, or, alternatively, the olecranon can be repaired by operation. See X-ray Nos. 10 and 11.

Fracture of the Olecranon

This is a common injury. If the patient is less than forty years of age, the elbow should be plastered in full extension for six weeks. When the limb is taken out of plaster it will be found to be quite stiff. It should be then flexed 10 degrees each week by the use of successive plaster slabs, until the position of right angles is achieved. The plaster slab is then abandoned and active movements only commenced.

Alternatively, and in all patients over forty, the olecranon should be repaired by operation. If the fractured fragment is small, it should be excised and the triceps repaired. In the elderly the fragment should be excised or ignored and active movements commenced.

Monteggia Fracture of Shaft of Ulna with Dislocation of Head of Radius

This type of injury is not uncommon. It is very difficult to treat satisfactorily. See X-ray No. 12. There are two types of Monteggia injuries. In the first type the ulna is angulated forwards and the head of the radius is dislocated anteriorly at the elbow. In the second type the ulnar shaft is angulated backwards, and the head of the radius is dislocated posteriorly. This is the commoner type.

Treatment of the first type in which the ulnar shaft is angulated forwards is to reduce the fracture of the ulna by open operation and fixation by a vitallium plate with four screws. The head of the radius can be replaced by manipulation at the same time.

In the second type of Monteggia injury treatment, where the ulna is angulated backwards, the best treatment is plaster in full extension for four to six weeks depending upon the age of the patient. In many Monteggia injuries the radial head is also fractured. In these cases the head of the radius should be excised early. However, when the head of the radius is not fractured, but only dislocated, excision of the head should be left until late. Otherwise the shaft of the radius rides upwards with manus valgus deformity.

"Myositis Ossificans" of Elbow

This condition is merely calcification of a haema-

toma. It is made worse by passive movements and manipulation. Treatment of this very difficult condition is immobilisation in plaster at right angles for four weeks only. The plaster is then removed, and active movements are undertaken. See X-ray No. 13.

Fractures of Shafts of Radius and Ulna

When both bones are fractured in mid-shafts they should be reduced by manipulation or open operation. The plaster should extend from the knuckles to the axilla. This should be kept on until strong union occurs. Reduction of the plaster to below the elbow is contra-indicated as it fails to immobilise the fracture.

If the mid-shaft of the radius alone is fractured and angulated with dislocation of the lower end of the ulna, the fracture should be reduced by open operation and fixed by one vitallium screw.

"Irreducible" Pronator Quadratus Fracture in Children

This type of injury is shown in X-ray No. 14. It is commonly irreducible by ordinary manipulation. Open reduction is easy, and gives excellent results.

Colles' Fracture-Dislocation of Wrist

Colles' fracture-dislocation of the lower end of the radius and ulnar styloid presents three displacements at once. They are radial deviation, dorsal displacement, and dorsal rotation of the distal fragment. See X-Rays 15 and 16. The hand is thus carried towards the thumb and dorsally. A deep anaesthetic is required. The fracture is reduced by disimpacting the fragments by dorsi-flexion. All three displacements are corrected at the one time. A three-quarter round plaster slab is applied on the radial side of the forearm, from the knuckles to the elbow. The plaster is trimmed to allow free movement of the elbow and fingers. The hand should be in the position of full ulnar deviation and in the same plane as the forearm, i.e. neither dorsi- nor palmar-flexed. Re-X-ray should show the lower articular surface of the radius to be facing slightly towards the palm. The position of this articular surface at right angles to the shaft of the radius is not sufficient.

The plaster should be changed under deep anaesthesia after ten days as all cases tend to relapse at this time. The plaster stays on for five weeks altogether. A sling is worn for twenty-four hours only. Active movements of fingers and elbow must be performed at once. The patient must do his normal work. If the fingers become swollen it is a sign that he is not using the hand

FIG. 16.—Colles fracture-dislocation. Antero-posterior view shows radial displacement of distal fragment.

and fingers enough, provided that the plaster is not too tight in which case it should be changed.

Smith's Fracture-Dislocation of the Wrist

This is the reverse deformity to a Colles. It is associated with an anterior marginal fracture of the lower end of the radius with subluxation of the wrist. The hand is carried palmar-wards. Treatment consists of manipulation under deep anaesthesia, and plastered in full dorsi-flexion. It is a difficult injury to hold in the correct position.

Injuries to the Hand Fracture of the Scaphoid.

A suspected fracture of the scaphoid must be X-rayed in three planes, i.e. antero-posterior, lateral, and oblique. If there is clinical evidence of fracture without X-ray signs the wrist should be plastered as for a scaphoid for three weeks and then re-X-rayed. Treatment consists of plaster with the thumb at right angles to the plane of the palm until the bone is united, i.e. twelve to twenty weeks or more. In late cases, where the diagnosis has been missed, drilling or bone-grafting is very successful.

It may be recalled that simple sprains of the wrist are rarer than fractures of the radius and scaphoid.

Dislocations of the Lunate

This should be reduced under deep anaesthesia by dorsi-flexion and traction, and application of direct pressure to the lunate. If operation has to

be undertaken it is better to excise the lunate than to replace it, as the latter treatment always causes aseptic necrosis of the lunate. The writer has met a surgeon who was able to amputate a leg through the thigh seventeen days after removal of the lunate upon his own wrist, following several attempts at manipulative reduction. Twenty years later he appears to have a normal wrist.

Bennett's Fracture-Dislocation of the Thumb

This is a fracture-dislocation of the base of the first metacarpal. Treatment should consist of traction with a pin through the pulp fixed to cramer wire, which is incorporated in a forearm and wrist plaster. The thumb should be in a plane at right angles to the palm, and extension should last for four weeks.

Fractures of the Metacarpals

Fractures of the shafts of the metacarpals other than that of the thumb should be treated by manipulation followed by active exercises only. The fingers should never be splinted nor immobilised in these fractures. Stiff fingers are the result.

Fractures of the neck of the metacarpals with forward angulation of the distal fragment should be corrected by right-angled flexion and backward pressure, and not by hyper-extension.

Fractures of the Phalanges

These fractures usually show forward angulation and should be treated by extension by strapping on either side of the fingers in flexion on a malleable wire splint on the palmar surface for ten days. Each finger should point to the tubercle of the scaphoid. No finger must ever be splinted in extension. Strapping should never be applied round a finger or toe. It should be applied longitudinally on either side.

All compound fractures of the fingers must be operated upon immediately. They can give more serious after-results than a fracture of the femur in a working man. In operating upon a compound fracture of the finger, the fracture should be reduced. Cut tendons should never be sutured at the original operation. Tendons should be sutured two weeks after healing in the case of a clean wound, and two months after healing in the case of an infected one. The skin should not be sutured at the original operation. Catgut should not be buried. The finger should be splinted in a palmar malleable wire or *plaster in flexion*. If traction is required, it should be done by means of two lateral strips of strapping.

All compound fractures of the fingers should be admitted as emergencies, and should be under the direct supervision of an experienced surgeon. The after-results of these injuries are still very depressing, and only too frequently result in the patient losing his occupation or being down-graded.

THE ART OF ORTHOPAEDICS

PART IV

INJURIES AND DISEASES OF THE SHOULDER AND ELBOW JOINTS

By G. O. TIPPETT, F.R.C.S.

(Hon. Asst. Orth. Surgeon to Queen's Hospital for Children and Croydon Hospital; Med. Off. Physiotherapy Dept. Bermondsey Health Centre)

Shoulder Joint

Owing to the varied uses to which the arms are subjected injuries of the shoulder joint are extremely common, ranging from injuries to the soft parts to dislocations and fractures. As in previous articles, fractures are not being discussed. The traumatic injuries which the examiner must bear in mind are the following:—

Sprains

A sprain may damage by tearing either some of the muscles surrounding a joint or may partially or completely tear the ligaments or capsule of a joint. In injuries to the muscles around the joint the patient usually can tell the examiner what particular movement brought on his trouble and the examiner, by making that group of muscles perform a particular movement, can reproduce the patient's pain. Similarly, if the examiner puts the joint through the opposite range of movement, that is passive movement, pain again will be reproduced. Sometimes, however, all muscles around a joint have been over-stretched, in which case all joint movements will be rather painful. At this stage it is well to bear in mind that what are commonly called the movements of the shoulder joint by the patient are not only the movements between the humerus and the scapula, but also the movements between the scapula and the chest wall, sometimes called the compensatory shoulder movement. The differentiation of these two groups of movement is extremely important in the

SELECTION FROM THE SUBJECTS ALREADY DEALT WITH

Carcinoma of the Lung	February —1943
Childlessness	August —1944
Diseases of Pregnancy	June —1944
Disorders of Micturition	December—1944
Gall-Bladder Diseases	February —1944
Intestinal Obstruction	March —1945
Mass Radiography	May —1944
Meningoencephalitis	March —1944 April —1944
Oesophageal Diseases	October —1944
Renal Tuberculosis	December—1943

CORRIGENDA

In the May issue of the POST-GRADUATE MEDICAL JOURNAL an incorrect caption appeared beneath Fig. 2 in E. H. Hambly's article "Fractures of the Upper Limb," this caption should read "Adduction Fracture of Neck of Humerus."

Owing to an error the April issue of the JOURNAL was numbered incorrectly 234 instead of 233.

H. K. LEWIS & Co. Ltd.*Medical Publishers
and Booksellers***136 GOWER STREET
LONDON, W.C.1***(Adjoining University College and Hospital)*

Telephone: EUSton 4282 (5 lines).

Telegrams: Publicavit, Westcent, London.

Medical Lending Library

ANNUAL SUBSCRIPTION from ONE GUINEA

**FOR THE CONVENIENCE OF POST-GRADUATE
STUDENTS SHORT PERIOD SUBSCRIPTIONS
ARE ARRANGED—FOR THREE OR SIX MONTHS***Detailed Prospectus on application***The Library Catalogue revised to December, 1943, containing classi-
fied index of authors and subjects. To subscribers, 12s. 6d. net;
to non-subscribers, 25s. net, postage 8d.****NEW BOOKS ADDED IMMEDIATELY UPON PUBLICATION**