Gardiner, and both he and many other senior assistants are responsible for much of what I have learnt and for a large part of whatever success I have had in treating patients who are critically ill.

My friend and colleague Mr. Albert Davis, has always put his knowledge and experience at my disposal and any merit which the part of this article dealing with pelvic surgery may have are due to his guidance; the faults are mine.

I have to thank Mr. Mortimer Woolf for permission to reproduce Fig. 5 and Dr. Leonard Symonds for the drawings in Figs. 6 and 7.

Finally, it is a pleasure to thank Sir Gordon Gordon-Taylor for permission to quote his writings. His experience is a sort of National Trust for his colleagues, and this is a welcome opportunity to acknowledge much wise advice given freely over many years.

ADIPOSITY

By RAYMOND GREENE, M.A., D.M., M.R.C.P.

(Hon. Physician Metropolitan Hospital)

The Importance of Adiposity*

In the words of Donald Hunter (1933) "the victims of obesity are . . . to be pitied for many reasons. They must of necessity suffer from a restriction of physical activity, they have a tendency to develop flat-feet and arthritis, they also have an unpleasant tendency to excessive sweating, and they are difficult to anaesthetise, operate on, and nurse. Moreover, they are predisposed to diabetes, bronchitis, and possibly to hyperpiesia." It may be added that obesity is a cause of great unhappiness to many women. Though a lucky few are prone to laugh and grow fat, the majority are sufficiently conscious of the preference for slimmness of the ordinary man to become self-conscious and self-critical, to lose their self-confidence and sometimes to sink into a state of severe depression. The importance of adiposity is well shown by the analysis by Dublin and Lotka (1936) of the life histories of 192,304 men accepted for life insurance.

Table 1

Influence of Weight on Mortality. Deaths per 100,000

Weight	Deaths
Standard	844
Underweight	848
Overweight	1,111
Overweight 5-14 %	1,027
Overweight 15-24 %	1,215
Overweight 25 % and over	1,472

When the age of the "life" is considered, the influence of adiposity becomes still more obvious.

* Adiposity is derived from the Latin *adepts* (fat), obesity from the Latin *ob* and *esus* (having eaten). The former is more generally accurate.

Table 2

Influence of Weight on Mortality. Effect of age, Deaths per 100,000

Weight	Age	
	Under 45	Over 45
Standard ..	463	1,308
Underweight ..	498	1,274
Overweight ..	527	1,824

Table 3

Influence of Overweight on Mortality in Persons aged 45 to 50

Pounds Overweight	Increase in Death Rate over Average
10	8
20	18
30	28
40	45
50	56
60	67
70	81
90	116

In other words, a mere stone of extra weight in middle age, lessens a man's expectation of life by about 10 per cent, and "fifty pounds overweight at age 45 imposes as much extra mortality as valvular heart disease" (Fisk, quoted by Newburgh, 1942).

The Definition of Adiposity

The decision as to whether a patient is too fat or not must obviously be largely a question of taste. Many Oriental races prefer fat women and even in one country fashions change. We must fall back upon averages considered in the light of common sense. The most reliable averages for children are probably those of the Child Health Association and for adults those of Fisk (1923). Anderson's Nomogram is very convenient in practice.

TABLE 4.—Weights of Boys from Birth to School Age

Height, In.	Weight, Lb.											
	1 Mo.	3 Mo.	6 Mo.	9 Mo.	12 Mo.	18 Mo.	24 Mo.	30 Mo.	36 Mo.	48 Mo.	60 Mo.	72 Mo.
20	8	..	..	..	..	..	..	..	..	..	..	..
21	9	10	..	..	..	..	..	..	..	..	..	..
22	10	11	..	..	..	..	..	..	..	..	..	..
23	11	12	13	..	..	..	..	..	..	..	..	..
24	12	13	14	..	..	..	..	..	..	..	..	..
25	13	14	15	16	..	..	..	..	..	..	..	..
26	..	15	17	17	18	..	..	..	..	..	..	..
27	..	16	18	18	19	..	..	..	..	..	..	..
28	..	..	19	19	20	20	..	..	..	..	..	..
29	..	..	20	21	21	21	..	..	..	..	..	..
30	..	..	22	22	22	22	22	..	..	..	..	..
31	..	..	..	23	23	23	24	24	..	..	..	..
32	..	..	..	24	24	24	25	25	..	..	..	..
33	..	..	..	..	26	26	26	26	26	..	..	..
34	..	..	..	..	..	27	27	27	27	..	..	..
35	..	..	..	..	..	29	29	29	29	29	..	..
36	..	..	..	..	..	..	30	31	31	31	..	..
37	..	..	..	..	..	..	32	32	32	32	32	..
38	..	..	..	..	..	..	..	33	33	33	34	..
39	..	..	..	..	..	..	..	35	35	35	35	..
40	..	..	..	..	..	..	..	..	36	36	36	36
41	..	..	..	..	..	..	..	..	..	38	38	38
42	..	..	..	..	..	..	..	..	..	39	39	39
43	..	..	..	..	..	..	..	..	..	41	41	41
44	..	..	..	..	..	..	..	..	..	..	43	43
45	..	..	..	..	..	..	..	..	..	..	45	45
46	..	..	..	..	..	..	..	..	..	..	..	48
47	..	..	..	..	..	..	..	..	..	..	..	50
48	..	..	..	..	..	..	..	..	..	..	..	52
49	..	..	..	..	..	..	..	..	..	..	..	52

TABLE 5.—Weights of Girls from Five to Eighteen Years

Height, In.	Weight, Lb.																	
	5 Yr.	6 Yr.	7 Yr.	8 Yr.	9 Yr.	10 Yr.	11 Yr.	12 Yr.	13 Yr.	14 Yr.	15 Yr.	16 Yr.	17 Yr.	18 Yr.				
38	33	33	..	..	..	..	..	..	..	..	..	..	..	..				
39	34	34	..	..	..	..	..	..	..	..	..	..	..	..				
40	36	36	36	..	..	..	..	..	..	..	..	..	..	..				
41	37	37	37	..	..	..	..	..	..	..	..	..	..	..				
42	39	39	39	..	..	..	..	..	..	..	..	..	..	..				
43	41	41	41	41	..	..	..	..	..	..	..	..	..	..				
44	42	42	42	42	..	..	..	..	..	..	..	..	..	..				
45	45	45	45	45	45	..	..	..	..	..	..	..	..	..				
46	47	47	47	48	48	..	..	..	..	..	..	..	..	..				
47	49	50	50	50	50	50	..	..	..	..	..	..	..	..				
48	..	52	52	52	52	53	53	..	..	..	..	..	..	..				
49	..	54	54	55	55	56	56	..	..	..	..	..	..	..				
50	..	56	56	57	58	59	61	62	..	..	..	..	..	..				
51	..	..	59	60	61	61	63	65	..	..	..	..	..	..				
52	..	..	63	64	64	64	65	67	..	..	..	..	..	..				
53	..	..	66	67	67	68	68	69	71	..	..	..	..	..				
54	..	..	..	69	70	70	71	71	73	..	..	..	..	..				
55	..	..	..	72	74	74	74	75	77	78	..	..	..	..				
56	..	..	..	..	76	78	78	79	81	83	..	..	..	..				
57	..	..	..	..	80	82	82	82	84	88	92	..	..	..				
58	..	..	..	..	..	84	86	86	88	93	96	101	..	..				
59	..	..	..	..	..	87	90	90	92	96	100	103	104	..				
60	..	..	..	..	..	91	95	95	97	101	105	108	109	111				
61	..	..	..	..	..	..	99	100	101	105	108	112	113	116				
62	..	..	..	..	..	..	104	105	106	109	113	115	117	118				
63	..	..	..	..	..	..	..	110	110	112	116	117	119	120				
64	..	..	..	..	..	..	..	114	115	117	119	120	122	123				
65	..	..	..	..	..	..	..	118	120	121	122	123	125	126				
66	..	..	..	..	..	..	..	..	124	124	125	128	129	130				
67	..	..	..	..	..	..	..	..	128	130	131	133	133	135				
68	..	..	..	..	..	..	..	..	131	133	135	136	138	138				
69	..	..	..	..	..	..	..	..	..	135	137	138	140	142				
70	..	..	..	..	..	..	..	..	..	136	138	140	142	144				
71	..	..	..	..	..	..	..	..	..	138	140	142	144	145				

TABLE 6.—Weights of Girls from Birth to School Age

Height, in.	Weight, Lb.											
	1 Mo.	3 Mo.	6 Mo.	9 Mo.	12 Mo.	18 Mo.	24 Mo.	30 Mo.	36 Mo.	48 Mo.	60 Mo.	72 Mo.
20	8	..	..	..	..	..	..	..	..	..	..	..
21	9	10	..	..	..	..	..	..	..	..	..	..
22	10	11	..	..	..	..	..	..	..	..	..	..
23	11	12	13	..	..	..	..	..	..	..	..	..
24	12	13	14	14	..	..	..	..	..	..	..	..
25	13	14	15	15	..	..	..	..	..	..	..	..
26	..	15	16	17	17	..	..	..	..	..	..	..
27	..	16	17	18	18	..	..	..	..	..	..	..
28	..	..	19	19	19	19	..	..	..	..	..	..
29	..	..	19	20	20	20	..	..	..	..	..	..
30	..	..	21	21	21	21	21	..	..	..	..	..
31	..	..	..	22	22	23	23	23	..	..	..	..
32	..	..	..	..	23	24	24	24	25	..	..	..
33	..	..	..	..	..	25	25	25	26	..	..	..
34	..	..	..	..	..	26	26	26	27	..	..	..
35	..	..	..	..	..	29	29	29	29	29	..	..
36	..	..	..	..	..	..	30	30	30	30	31	..
37	..	..	..	..	..	..	31	31	31	31	32	..
38	..	..	..	..	..	..	..	33	33	33	33	..
39	..	..	..	..	..	..	..	34	34	34	34	34
40	..	..	..	..	..	..	..	..	35	36	36	36
41	..	..	..	..	..	..	..	..	..	37	37	37
42	..	..	..	..	..	..	..	..	..	39	39	39
43	..	..	..	..	..	..	..	..	..	40	41	41
44	..	..	..	..	..	..	..	..	..	..	42	42
45	..	..	..	..	..	..	..	..	..	..	..	45
46	..	..	..	..	..	..	..	..	..	..	..	47
47	..	..	..	..	..	..	..	..	..	..	..	50
48	..	..	..	..	..	..	..	..	..	..	..	52

TABLE 7.—Weights of Boys from Five to Nineteen Years

Height, in.	Weight, Lb.																		
	5 Yr.	6 Yr.	7 Yr.	8 Yr.	9 Yr.	10 Yr.	11 Yr.	12 Yr.	13 Yr.	14 Yr.	15 Yr.	16 Yr.	17 Yr.	18 Yr.	19 Yr.				
38	34	34	..	..	..	..	..	..	..	..	..	..	..	..	..				
39	35	35	..	..	..	..	..	..	..	..	..	..	..	..	..				
40	36	36	..	..	..	..	..	..	..	..	..	..	..	..	..				
41	38	38	38	..	..	..	..	..	..	..	..	..	..	..	..				
42	39	39	39	39	..	..	..	..	..	..	..	..	..	..	..				
43	41	41	41	41	..	..	..	..	..	..	..	..	..	..	..				
44	44	44	44	44	..	..	..	..	..	..	..	..	..	..	..				
45	46	46	46	46	46	..	..	..	..	..	..	..	..	..	..				
46	47	48	48	48	48	..	..	..	..	..	..	..	..	..	..				
47	49	50	50	50	50	50	..	..	..	..	..	..	..	..	..				
48	..	52	53	53	53	53	..	..	..	..	..	..	..	..	..				
49	..	55	55	55	55	55	55	..	..	..	..	..	..	..	..				
50	..	57	58	58	58	58	58	58	..	..	..	..	..	..	..				
51	..	..	61	61	61	61	61	61	..	..	..	..	..	..	..				
52	..	..	63	64	64	64	64	64	64	..	..	..	..	..	..				
53	..	..	66	67	67	67	67	68	68	..	..	..	..	..	..				
54	..	..	..	70	70	70	70	71	71	72	..	..	..	..	..				
55	..	..	..	72	72	73	73	74	74	74	..	..	..	..	..				
56	..	..	..	75	76	77	77	77	78	78	80	..	..	..	..				
57	..	..	..	..	79	80	81	81	82	83	83	..	..	..	..				
58	..	..	..	..	83	84	84	85	85	86	87	..	..	..	..				
59	..	..	..	..	..	87	88	89	89	90	90	90	..	..	..				
60	..	..	..	..	..	91	92	92	93	94	95	96	..	..	..				
61	..	..	..	..	..	..	95	96	97	99	100	103	106	..	..				
62	..	..	..	..	..	..	100	101	102	103	104	107	111	116	..				
63	..	..	..	..	..	..	105	106	107	108	110	113	118	123	127				
64	..	..	..	..	..	..	..	109	111	113	115	117	121	126	130				
65	..	..	..	..	..	..	..	114	117	118	120	122	127	131	134				
66	..	..	..	..	..	..	..	..	119	122	125	128	132	136	139				
67	..	..	..	..	..	..	..	..	124	128	130	134	136	139	142				
68	..	..	..	..	..	..	..	..	..	134	134	137	141	143	147				
69	..	..	..	..	..	..	..	..	..	137	139	143	146	149	152				
70	..	..	..	..	..	..	..	..	..	143	144	145	148	151	155				
71	..	..	..	..	..	..	..	..	..	148	150	151	152	154	159				
72	..	..	..	..	..	..	..	..	..	..	153	155	156	158	163				
73	..	..	..	..	..	..	..	..	..	..	157	160	162	164	167				
74	..	..	..	..	..	..	..	..	..	..	160	164	168	170	171				

TABLE 8.—*Ideal Weights of Men and Women **

Men			Women		
Height		Weight	Height		Weight
Ft.	In.	Lb.	Ft.	In.	Lb.
5	..	126	4	8	112
5	1	128	4	9	114
5	2	130	4	10	116
5	3	133	4	11	118
5	4	136	5	..	120
5	5	140	5	1	122
5	6	144	5	2	124
5	7	148	5	3	127
5	8	152	5	4	131
5	9	156	5	5	134
5	10	161	5	6	138
5	11	166	5	7	142
6	..	172	5	8	146
6	1	178	5	9	150
6	2	184	5	10	154
6	3	190	5	11	157
6	4	196	6	..	161
6	5	201			

* Adapted from Fisk.

Table 9 (Nomogram).

The Classification of Adiposity

It has been customary in the past to attempt a differential diagnosis of adiposity on the basis of the distribution of the excessive fat.

A study of my own records and of those of others has convinced me that there is no justification for any such assumption. The distribution appears to be decided in part by racial and in part by individual characteristics. There is, for instance, insufficient justification for previous assertions that "pituitary adiposity" is characterised by a fat body with slim limbs or thyroid adiposity by a general distribution. Either type may be associated with any cause, though the special case of Cushing's Syndrome seems to be most often of the former.

The Causes of Adiposity

Despite much sound research and vague speculation, the cause of adiposity remains obscure. Newburgh (1942) has considered the claims of various hypotheses.

1. Do fat people overeat?

Considerable authorities have gone so far as to say that almost all fat people are fat because they eat too much. That an element of gluttony is common cannot be denied, but experience shows that most fat people can only remain slim by eating much less than the average amount of food and that most slim people can over-eat with impunity. The most that it is justifiable to say of gluttony is that it will inevitably bring out any constitutional tendency to adiposity.

2. Is there a disorder of the use of energy?

We have said that some people who eat well remain thin, whereas others who eat little grow fat. This observation might be explained by a difference in basal metabolic rate, but Boothby

To use the nomogram, the height and weight are measured and a straight line joining the points on the height and weight scales is produced to cut the percentage scale. The point of intersection gives the percentage over ideal weight. If the zero point on the percentage scale is joined to the appropriate point on the height scale, the intersection on the weight scale gives the ideal weight. (By kind permission of A. B. Anderson, B.Sc., Ph.D., M.R.C.S.).

and Sandiford (1922) have shown that such an explanation is untenable. Of 94 adipose patients, 81 had a normal B.M.R. Strouse, Wang and Dye (1924) found no difference between two series of adipose and normal people. Grafe (1933) found only three with a low B.M.R. among 180 grossly adipose patients. A definitely low B.M.R. in an adipose person can probably be explained always by another abnormality, usually myxoedema.

Another possible explanation of adiposity might be found in a reduction in the specific dynamic action of food in fat people, and many investigators have claimed to have established that this in fact can be found. The subject has been considered most critically and investigated afresh by Strang and McClugage (1931) who established beyond doubt that there is no significant difference in this respect between fat and thin people. Lauter (1926) indeed had shown that the normal variations in specific dynamic action are so great that nothing short of its total absence can be considered significant.

The hypothesis of "Luxuskonsumption" has also had a considerable vogue. Specific dynamic action comes to an end within twelve hours, but Grafe and Graham (1911) sought to show that a high intake of food produces an increased metabolism over a much longer period, whereas a low intake reduces the metabolic rate. Such a mechanism would be capable of a very exact control over body weight, and its failure would explain adiposity. The hypothesis appears to be supported by clinical experience, but a more careful scrutiny shows that it is only half true. Wiley and Newburgh (1931) showed that in fact the consumption of excessive food does not stimulate metabolism. That starvation diminishes it is, however, certain (Schondorff, 1912; Grafe and Koch, 1912), and some of the lowest basal metabolic rates are to be found in cases of anorexia nervosa. An explanation is afforded of the failure of starving people to die as soon as might be expected, but not of the phenomenon of adiposity.

3. *Is there a Disorder of Absorption?*

It is clear that if some people absorb more food than others, the more efficient absorbers would tend, other things being equal, to be fatter. This hypothesis was disposed of by Neuenschwander-Lemmer (1936), who showed that the residual calories in the faeces of fat people are not significantly different from those of thin, and that the utilisation of dietary calories, nitrogen and fat is also equal.

4. *Is there a constitutional abnormality of the Adipose Tissue?*

Von Bergmann suggested that there is an

increased affinity for fat in the adipose and Hetenyi (1936) that the adipose tissue of these persons "hung on to the fat" and prevented its use as fuel. These suggestions are not borne out by experiment. Indeed the opposite is apparently the case (Newburgh, 1942).

In his recent Sharpey-Schäfer Lectures before the Royal College of Physicians, Lawrence (1946) postulated the existence of a lipopexic hormone. He described a woman who suffered from complete disappearance of all body fat, hyperglycaemia and glycosuria without ketosis and hyperlipaemia. He suggested that owing to a deficiency of his hypothetical lipopexic hormone, the patient was unable to store fat and that in consequence the fat derived from her dietary carbohydrate was forced to circulate in her blood stream.

We must finally be prepared to admit that we do not know by what process adiposity comes about. An approach to the problem may however be made by considering the rôles played by some of the organs known to be concerned.

The Rôle of the Hypothalamus

It has been recognised for many years that tumours such as craniopharyngiomata and inflammation at the base of the brain may cause adiposity. The best-known examples of obesity due to tumours are the sufferers from Fröhlich's syndrome. Smith (1930) produced a similar condition of dystrophia adiposo-genitalis by injecting chromic acid into the hypophyses of animals by a technique which damaged the hypothalamus. Since removal of the hypophysis does not cause obesity, it may be assumed that the hypothalamic damage was responsible. Ranson and his colleagues (1938) using the Horsley-Clarke apparatus, produced obesity in twelve rats by accurate hypothalamic destruction without damage to the pituitary. According to Hetherington and Ranson (1942) the damage must be extensive and bilateral. Obesity cannot be produced by hypophysectomy in the absence of damage to the hypothalamus (Ranson *et al* 1938) and the obesity produced by hypothalamic damage is unaffected by subsequent hypophysectomy (Hetherington and Ranson, 1942).

In man adiposity is a common sequel to encephalitis (Eaves and Croll, 1930; von Economo, 1931) which may, on the contrary, be followed by emaciation. Eaves and Croll reported that in ten patients examined postmortem (two of whom were fat and four emaciated) the hypothalamic region was more severely affected than any other part of the brain except the substantia nigra. They formed the opinion that less severe lesions were more likely to cause obesity.

The exact site in the hypothalamus which is

concerned is still unknown. There is some surgical evidence that in man the area in question is in the anterior tuberal wall. It is not, at any rate in animals, identical with the water metabolism centre (Hetherington and Ranson, 1942), but it would not be sound to assume that this is true of men, owing to the remarkable vagrancy of cerebral function.

Hypothalamic adiposity may be diagnosed with confidence only when other signs of hypothalamic damage are present. Of these, the most important are diabetes insipidus, oliguria, changes in the sleep rhythm, sexual retardation or atrophy, and gross disorder of the autonomic nervous system. The majority of adipose patients show no such signs or symptoms. Nevertheless a careful search brings to light the fact that a very large proportion have in fact a disorder of their water balance.

Textbooks of medicine are strangely silent about the fact that pitting oedema is almost always present in the lower part of the legs of grossly adipose patients, even in the absence of any demonstrable renal or cardiac lesion. When it is remembered that an increased water content of many pints must be present before pitting can be demonstrated, the fact that oedema is usually (but not always) absent in less extreme cases is not surprising. The importance of water-retention has often been stressed previously, especially by Zondek (1944) and Bauer (1933). Von Grafe (1933) wrote "fatty tissues absorb water like a sponge without circulatory disturbances and with no evidence of oedema," but in fact oedema is often demonstrable. The Volhard water test often shows that fluid-retention occurs in obese patients (Wohl, 1932; Rowntree and Brunsting, 1933). We are brought to conclude that obesity so-called may in certain cases be accounted for largely if not solely by abnormal retention of tissue fluids.

The occasional failure of underfeeding to reduce weight is well illustrated by a case described by Newburgh (1942). The patient was a young woman who on entering hospital weighed 125 kilograms. Her B.M.R. was normal (2,100 calories in 24 hours). Although she was kept in bed, her heat production averaged 2,800 calories a day. She was given a diet yielding 1,600 calories, and it was calculated that in order to provide the excess caloric output it was necessary for her to oxidise about 150 grams of body tissue daily. Her weight should therefore have fallen by this amount daily. In fact an insignificant loss of weight occurred for the first ten days, an actual slight increase occurring for the first two days. A very sudden loss of weight then occurred, the theoretical weight being attained in thirteen days. A study of her water balance showed that the failure to lose weight was accounted for by water

retention and the sudden loss at the end of thirteen days by the excretion of the excessive fluid. This is not a rare and isolated case. In fact, anyone experienced in the treatment of obesity is aware that a delay in weight loss is usual in patients treated with diet alone, and Newburgh (1942) has shown that even in normal people reduction in diet is associated with temporary water retention. In both obese and normal people this delay may be overcome by the exhibition of diuretics.

The Rôle of the Posterior Pituitary

In many of its functions the posterior pituitary must at present be regarded not as an individual structure but as a part of the hypothalamic-pituitary apparatus. This is particularly true of its influence on water metabolism. Diabetes insipidus, for instance, may be due to a lesion of the supraoptic nucleus of the hypothalamus, of the supraopticohypophyseal tract, or of the posterior pituitary. Enough has been said already to make it clear that adiposity and fluid retention are closely related phenomena. The experiments of Verney and his collaborators, although performed on dogs, are of undoubted human significance. They have shown that under the influence of emotion, the posterior pituitary excretes into the bloodstream a hormone which, by its action on the kidney tubules causes a diminution in the excretion of water and its consequent retention in the tissues. That inhibition of normal urinary flow accompanies emotional states such as anxiety and fear, was shown by Bernard (1859), Mackeith and his colleagues (1923), Dobreff (1926), and Klisiecki *et al* (1933). Theobald and Verney (1935) showed that the inhibition takes place after complete denervation of the kidney even after the mildest stimuli such as cutting the hair of the lumbar region and even after the preliminary lathering. They showed, too, that the inhibition is not caused by adrenaline. That the posterior pituitary is responsible is suggested by the exactness with which the effects of anxiety may be mimicked by the injection of posterior pituitary extract. The inhibition may be brought about by as little as 0.0005 of an oxytocic unit. There is the same lag of about ten minutes as after nervous stimuli and albuminuria sometimes occurs whether pituitary extract or nervous stimulation is used. Rydin and Verney (1933) went on to show that the inhibition occurs even after section of the renal nerves, removal of the right adrenal and denervation of the left, the decentralisation of the whole abdominal sympathetic system and the removal of the ganglia from L2 to S1 inclusive. They confirmed that the course of events after emotional stress can be exactly mimicked by the intravenous injection of

as little as 0.25 mg. of posterior pituitary powder, and concluded that emotion stimulated the posterior pituitary by way of the hypothalamus and the supraopticohypophyseal tracts. O'Connor and Verney (1944) finally completed the chain of evidence by showing that the inhibition of urinary flow by emotion is almost entirely destroyed by removal of the posterior pituitary.

It is generally assumed that in man anxiety causes a loss of weight. This is commonly but not invariably true. It apparently has not been pointed out previously that anxiety may cause an *increase* in weight.

A, an intelligent and well-educated married woman of 28, 5 ft. 2 in. in height, weighed 8 stone 7 lbs. until August 1941, when she learned that her husband was missing. She was desperately depressed by the news, sometimes prostrate with grief at the thought of his possible death, at others tortured by the thought of his possible maltreatment by the Japanese. She began immediately to gain weight. In May 1942 she weighed 11 stone. She had complete amenorrhoea during this period and suffered from breathlessness on exertion and palpitations of the heart. On examination in May 1942 she was found to be very obese. There was obvious oedema of the ankles and shins. Nothing else abnormal was found and a complete investigation failed to find any cardiac or renal cause of the oedema. Treatment in hospital with a diet poor in carbohydrates and with urea reduced her weight to 9 stone 13 lb., but when she returned to her occupation as a physical training instructress she rapidly regained her previous weight. She volunteered the information that she always lost weight if she rested a lot. The diuretic effect of recumbency is well known. During 1943 she emerged from her despondency and formed an attachment for a wealthy man who took her daily to the best restaurants. She ceased to follow her diet. Her weight fell rapidly to about 9 stone and menstruation returned to normal.

B, a married woman of 29, 5 ft. 3 in. in height, was fairly slim until June 1945, when she weighed 10 stone 5 lbs. Her only child then got otitis media and afterwards mastoiditis. The mother, who was extremely anxious, began to get fatter. The mastoiditis was followed by a series of complications. B was first seen in March 1946, when her weight was 12 stone 10 lb. The child was still in hospital and the mother so distressed that she was unable to talk of the illness without tears. Nothing abnormal could be found other than obesity and pitting oedema of the ankles and shins. The condition is responding to treatment by diet and urea.

Apart from its effect on water metabolism, the posterior pituitary appears to have an influence on the transport of fat. After the injection of

posterior pituitary extract into rabbits and rats, Coope and Chamberlain (1925) observed fatty infiltration of the liver of such a degree that in some cases the liver was doubled in size.

The Rôle of the Anterior Pituitary

The importance of the anterior pituitary in the production of adiposity has been overstressed since Fröhlich first described his syndrome in 1901. Of this disorder, due to a pituitary tumour, obesity is a cardinal sign. It is now realised that this sign is, however, due not to the pituitary tumour directly, but to its effects upon the neighbouring hypothalamus. Yet the concept of pituitary obesity thus born has survived. It is instructive to observe that even experienced observers like Le Marquand and Tozer (1943), who show between the lines that they have ceased to believe in it, find it difficult to say so in so many words.

Apart from Fröhlich's syndrome, two conditions call attention to the possible influence of the anterior pituitary in producing adiposity; Dercum's disease and Cushing's disease. Dercum's disease is a rare syndrome in which symmetrical nodular deposits of fat occur on the back of the neck, the shoulders and arms, the trunk, hips, thighs and medial aspects of the knees. The nodules are extremely tender on pressure and may be spontaneously painful. There is usually a generalised adiposity as well. As in other examples of extreme obesity, amenorrhoea is a frequent symptom. The patient is usually unduly tired and depressed, symptoms which may well be due to the increased effort of movement and consciousness of her bizarre appearance. The origin of the disorder is obscure and has been laid at the door of the thyroid (by Dercum himself), the anterior pituitary, the ovaries and the nervous system. I have been driven to the conclusion that no sound evidence incriminates any endocrine gland and that there is no essential difference between Dercum's disease and the more common types of obesity. The pain in the nodules appears to be the result of tension. That the tension may be the result of an excessive fluid content is suggested by the facts that the fat deposits in other obese patients are often tender and that in both general and nodular obesity the pain is often relieved by dehydration.

Cushing's original description of his syndrome includes obesity as a cardinal feature, a fact often quoted as evidence for the existence of adiposity of anterior pituitary origin. It is doubtful, however, whether the inclusion of this sign is sound. The typical protuberance of the abdomen may be due to thinning of the bodies of the vertebrae or even to collapse, resulting from the osteoporosis often present. Newburgh (1942) quotes cases of

his own, reviews other publications, and concludes that obesity is not characteristic of Cushing's syndrome. Albright, Parson and Bloomberg (1941) drew attention to this apparent obesity in three patients in whom the bulging abdomen was due not to actual fat but to crushed vertebrae and relaxed abdominal walls. I recently observed a case of a woman with Cushing's syndrome who presented the typical appearance of "fat" body and slim limbs, but who was found at postmortem examination to have no more than a normal subcutaneous deposit of fat. Even supposing that the consideration of a sufficiently large number of cases shows that obesity is more common in Cushing's syndrome than in the general run of people of a similar age group, much more must be known about the aetiology of the disease before the obesity can be laid at the door of the anterior pituitary. The influence of the adrenals, for instance, must be borne in mind.

The Rôle of the Thyroid

The importance of the thyroid has been overestimated with less excuse than has that of the anterior pituitary. It is a safe bet that any student asked to give the cardinal signs of hypothyroidism will include obesity among them. In fact patients with deficient thyroids are fat no more often than those whose thyroids are normal. Adiposity and hypothyroidism are both common at the change of life, a fact which has created a false impression. Hypothyroid patients are often emaciated. Mackay and Sherrill (1941) observed the effects of an equal diet on two groups of rats, one of which consisted of animals whose thyroids had been removed. The normal rats gained 193 gms. in about a year, the thyroidectomised rats only 178 gms. The bodies of the former contained 31.1 per cent of fat, those of the latter only 6.4 per cent. The thyroidectomised animals looked just as plump, presumably because they contained more water. Plummer (1940) studied 200 myxoedematous patients. In 8.5 per cent the weight was more than 50 lbs. above normal, but in 5.5 per cent it was more than 30 lbs. below normal. The higher the B.M.R., the smaller was the excess and vice versa. Those whose B.M.R. was above - 28 per cent had an average excess of weight of 11.5 per cent; those whose B.M.R. was below - 38 per cent weighed an average of 14.7 per cent too little. A few days of treatment with thyroid resulted in the loss of excessive fluid and the average weight of the still hypothyroid group dropped from 10.1 lbs. too much to 2.9 lbs. too little. Once again the importance of water metabolism is accentuated. It must, however, be remembered that the estimation of the B.M.R. is not strictly a measure-

ment of thyroid activity. It is what it claims to be, a measurement of the metabolism of the patient at the time of the test. There are many factors other than thyroid activity which may influence it. It seems probable that the best test for thyroid activity is the measurement of the blood iodine, the technique of which is difficult. Until this or some better method has become generally available, it is important to avoid confusing hypometabolism with hypothyroidism. The B.M.R. must be interpreted in the light of the clinical condition of the patients. This was done in Plummer's series, all the members of which were not merely hypometabolic but myxoedematous.

The Rôle of the Adrenals

Patients with Addison's disease are invariably thin, whereas those with adrenal hypertrophy are often, though not always, very fat. When adiposity is associated with the presence of an adrenal tumour, the excessive weight is rapidly lost after operation. Whereas before operation, these patients are singularly resistant to dietary treatment, loss of weight afterwards occurs with a normal diet.

The experimental facts have been reviewed by Goldzieher (1945). Both normal animals and those whose adrenals have been removed, gain weight when given cortical extract. This may be in part due to better absorption from the small intestine (Verzar and Laszt, 1936); patients with Addison's disease occasionally showing steatorrhoea.

Conversely, adrenalectomised rats lose weight and there is ultimately almost complete loss of body fat in those which survive long enough. Cortical extract has a double effect, upon fat formation and upon the fat storing cells. Much of the body fat is derived from carbohydrate and from protein. In cortical deficiency the production of carbohydrate from protein is reduced, the liver glycogen falls and hypoglycaemia occurs. In such conditions all available sugar is needed for energy production and fat is not formed. In addition, the injection of cortical extract causes a fall in blood lipids, owing to these being taken up by various tissues, especially of the reticulo-endothelial system. This drop is greatly reduced if the reticulo-endothelial system has been blocked previously by the injection of dyes or of Indian ink. The evidence of the blood cholesterol is, however, difficult to interpret, for in Addison's disease it is low and in overactivity of the cortex, it is usually high. Though in the former the low values might be explained by depletion of the cholesterol stores, the high values in the latter appear inconsistent with experimental findings.

The blood lipoids are, in fact, high in most cases of obesity of whatever origin, but whereas in normal people the fat content of the blood remains constant when the caloric value of the diet is reduced, in obese patients it falls by 18 to 43 per cent (Hetenyi, 1936; Beznak, 1938). After the injection of 60 grams of fat, the blood fat of normal people rises on an average by 84 per cent, but of obese patients by only 12 per cent (Zondek, 1944). This finding is explicable on the assumption that under the influence of the cortical hormones, the fat is more rapidly taken up by the tissues of the obese, but direct evidence for such an explanation is lacking.

The action of the adrenal cortex on water metabolism is highly complex and not yet entirely clear. In adrenal deficiency, the capillary permeability is increased to such an extent that circulatory collapse cannot be countered by the injection of fluids, even of whole blood or serum, because in the absence of cortical activity they immediately leak out into the tissues and cause oedema. On the other hand, a great loss of water accompanies the loss of sodium and chloride in adrenal deficiency, and the consequent dehydration may be extreme. At the moment, the relation of these facts to obesity of apparently adrenal origin cannot be clearly seen.

The Rôle of the Gonads

The castration of animals and birds with a view to fattening them has been practised immemorially and the eunuch of the harem is always represented as of enormous corpulence. In man, adiposity is not, however, invariably a sequel to castration. The Skopcecs, a Russian religious sect in which castration before puberty is enforced, are not all obese, and obesity does not as a rule follow when the testicles are removed after puberty. The congenital eunuch is usually tall and thin. Obesity commonly but by no means invariably follows oophorectomy and the menopause. The inconsistency with which adiposity follows gonadectomy suggests that the connection, if it exists, is indirect.

Many attempts have been made to show that the gonads directly effect general metabolism, but the results have been so conflicting that no definite conclusions can be drawn. It seems likely that the adiposity of castrates may be explained as the result of diminished activity in addition to a hereditary tendency to lay down fat. Many elderly people, and perhaps castrates, become more interested in food as their libido diminishes. At the menopause the adjustment of the endocrine system is complicated and in many women there is clinical suggestion that the obesity should be considered not of hypogonadic but of hypedrenal origin.

The Rôle of the Pancreas

Two apparently conflicting facts have led to much discussion of the rôle of the pancreas in adiposity, much of which has been futile because based upon the old "fixation" of interest on the pancreas in the mechanism of diabetes. In the first place a high proportion of adipose people have an abnormally high tolerance for carbohydrates (a low glucose tolerance curve) and may react to a reduction of carbohydrate with hypoglycaemic symptoms. Embleton (1938) found that 27 per cent of fat men and 65 per cent of fat women were in this category. In the second place, the diabetes which develops in later life is commonly associated with adiposity.

The association of obesity with a low glucose tolerance curve was neatly explained by Falta by the hypothesis that the increased sugar tolerance is due to hypernesia* which increases the appetite and so leads to obesity. Unfortunately the more extreme forms of hypoglycaemia are usually found in thin people. Were Falta's hypothesis true, it might be extended to explain the apparently anomalous association of a low sugar tolerance with adiposity by regarding senile diabetes as "the punishment of gluttony," the continued over-eating being due to the over-production of insulin and the ultimate death of the islets to overwork. Alternatively one might suggest that over-eating causes first hypertrophy and ultimately exhaustion of the islets. This is pure theorising. The careful questioning of my own group of fat diabetics suggests that an insignificant proportion of them have been unduly gluttonous. In most cases the diabetes has been discovered within a very few years of the development of the adiposity, which suggests that either the adiposity and the diabetes were coincident or that retribution followed rapidly on gluttony. Against the latter possibility is the fact that none of my cases of adiposity with hypoglycaemia, many of whom have been fat for many years, have so far become diabetic.

In the present state of our ignorance it seems better to make no attempt to relate the hypoglycaemic with the hyperglycaemic cases. The hypoglycaemic cases may perhaps be hypothalamic in origin for there is often a considerable degree of water retention. An attempt has been made to explain the hyperglycaemic cases by the work of Young (1941) on the pituitary diabetogenic factor. His extract causes accelerated growth but no diabetes in puppies, diabetes without growth in adult dogs. This difference is probably due to the presence in the extract of a pancreatrophic as well as a diabetogenic factor. The islets of the

* Hyperinsulinism is an ugly and inaccurate hybrid which should be dropped.

puppy respond sufficiently well to the former to overcome the effects of the latter, and in doing so cause increased growth. On the other hand, the islets of the grown dog respond less well (perhaps because they are relatively fewer in number) and the animal becomes diabetic. It does, however, increase in weight, and there is a temptation to ascribe the adiposity of the senile diabetic to a similar mechanism. This is hardly justified by the facts. The increase in weight is due to the nitrogen retaining action of pituitary extract, perhaps mediated by the islets of the pancreas. There seems to be no evidence of a fat-retaining or water-retaining effect. Indeed Young (1944) states that a study of the composition of the tissues shows that the increase in weight is due to an increase in the protein containing tissues.

It is well known that the fat elderly diabetic, suffering from the "diabète gras" of the old French physicians, does not suffer from ketosis except when he is treated with a slimming diet, when some ketosis would be expected in a patient who is not diabetic at all. He, or more commonly she, is relatively insensitive to insulin, and usually loses all signs and symptoms of diabetes if she can be persuaded to remain on a low carbohydrate diet until she has attained her correct weight. Lawrence (1946) suggests, tentatively but very plausibly, that the disorder of these patients is due to an excess of a lipopexic hormone or a deficiency of a lipocatabolic hormone. The hyperglycaemia is due, he suggests, not to any deficiency of insulin production, but to overfilling of the fat depots to such an extent that they are unable to accept further fat from the ingested carbohydrate, which is thus forced to circulate uselessly until the kidney excretes it.

The Treatment of Adiposity

It has been made clear that we do not yet understand the causes of adiposity. The treatment therefore can be based on no sound rationale. It must be symptomatic. There are, however, two points of view to be kept simultaneously in mind. The patient is suffering from both fat-retention and water-retention. To neglect either is to obtain only partial success.

Diet

Whatever the cause of the abnormal deposit of fat, the law of the conservation of matter applies. Fat must come from food, and whether we believe with some authorities that all fat people eat too much or with others that they do not, it remains certain that if the patient eats less food than he requires for energy, he must lose flesh. Experience

shows that in starvation, fat is lost more rapidly than more useful tissues. Partial starvation is the essence of treatment.

A simple reduction in calories, leaving the proportions of fat, protein and carbohydrate unaffected, is often successful. With a clear-sighted and collaborative patient it may be enough to say "Whenever you help yourself, think how much you would take in the past and then take half as much." With others, the Lawrence Line Diet,* designed for the treatment of diabetics, may be more useful. Each "line" of this diet gives approximately 150 calories, made up of 10 gms. of carbohydrate, $7\frac{1}{2}$ gms. of protein and 9 gms. of fat. It is only necessary to explain the very simple working of the scheme and to prescribe so many lines a day. The patient has a very wide choice of foodstuffs and need not be confined to a fixed menu. In cases in which I have used this method, I have usually prescribed about half the calculated requirements of the patient in question. The ordinary requirement of a sedentary adult is about 2,600 calories. For such a patient I would order about 1,300 calories a day; that is about 9 Lawrence Lines. The loss of weight on such a diet, provided the patient collaborates honestly, is slow but steady. It is necessary to ensure that enough protective foods are consumed. Green vegetables are, in the Lawrence diet, allowed *ad libitum*, and the patient must be encouraged to eat of them a sufficient quantity to ensure an adequate intake of vitamins A and C. In these days of meat shortage it may be as well to prescribe an additional source of the B vitamins and of iron. Provided that green vegetables are eaten in good quantity, hunger is not a serious disadvantage after the first few days.

In those patients, and they are numerous, in whom the element of water-retention is obvious, a delay may occur on such a diet before the loss of weight begins. There is reason to believe that starvation itself may cause temporary water-retention to a degree which may mask the loss of fat. Even if this is not so, the failure to excrete the excess of water may be at first depressing to the patient. This phase is, however, temporary, as Fig. 2 (from Newburgh, 1942) shows. In any case, the phase of water-retention may be eliminated by the use of a diuretic. It is possible that carbohydrates may have a water-retaining action (Földes, 1933). It is certainly true that in cases of gross obesity, in which water-retention is invariably great, the best results are not achieved if attention is focussed entirely on calories and if a diet low in energy value but normal in constitution is prescribed. In such cases better results are

* Obtainable from Messrs. Lewis of Gower Street, London, W.C.

FIG. 2.—The circles represent the body weight corrected for water retention. The retention of 2,900 grams of water in sixteen days explains the failure to lose weight, since 2,900 grams of body tissue was destroyed during the same period.

achieved by the drastic reduction of carbohydrates. Fats at present need not concern us. Experience shows that the amount of fat at present available is harmless.

It is my practice to give to all those patients in whom a great and rapid reduction in weight is necessary a very simple if somewhat onerous diet.

Salt in the diet should be kept as low as possible, for a high salt intake leads to water-retention.

It is surprising how large a proportion of fat people are ready to abide by these instructions for many months. When once a comparatively normal weight has been achieved, it is possible to "slacken off." Obviously a larger intake of food can be allowed when once it is no longer necessary to absorb less than the amount necessary for maintenance. Nevertheless the constitutionally adipose patient can never return to a normal diet without a return of corpulence. The treatment of obesity is purely symptomatic. The constitutional abnormality remains.

This diet makes no claim to scientific accuracy, and allows the consumption of certain carbohydrate foods (e.g. beetroot) and disallows certain foods with little carbohydrate (e.g. turnip). It is, however, easy to remember and therefore easier to follow than a regime more accurately constructed.

TABLE 10

DIET FOR OBESITY

Foods to be avoided:

1. Bread and everything else made with flour.
2. Cereals, including breakfast cereals and milk puddings.
3. Potatoes and all other white root vegetables.
4. Foods containing much sugar.
5. All sweets.
6. Salt. If possible, no salt should be used in the cooking; other members of the family adding it at the table.
7. Fluids should be limited to 2 pints a day.
8. Fats should not be eaten in excess, but the small amount at present available is generally permissible.

You can eat as much as you like of the following foods:

1. Meat, fish, birds.
2. All green vegetables.
3. Eggs, dried or fresh.
4. Cheese.
5. Fruit, if unsweetened or sweetened with saccharin.

You should weigh yourself once a week at the chemist, using the same chemist and wearing the same clothes. On this diet you should lose between one and two pounds a week. If you lose more than two pounds, you are being unnecessarily strict. If you lose less than one pound, you are not being strict enough.

It is found that patients are discouraged by over-detailed instructions. The diet may be made less monotonous by the use of the carbohydrate-free foods manufactured by reliable firms (e.g. Callard & Co.), but it must be remembered that most of these preparations contain a small amount of carbohydrate. The proof of the pudding is in the eating, both with these and ordinary foods. If the patient can eat a certain carbohydrate food and still lose weight, there is no need to adopt a doctrinaire attitude. Alcoholic drinks should be forbidden, not only because they have food value in themselves, but because they increase the appetite.

Thyroid

It has been pointed out that thyroid deficiency is not an important cause of adiposity. Nevertheless, thyroid extract undoubtedly has a use in treatment. Its value probably lies less in its capacity to raise the general metabolism of the body than in its effect on water balance. Volhard's water test, often abnormal in obese people, may be restored to normal by thyroid treatment. The delay in weight reduction which often occurs on a simple dietetic treatment may be prevented by the simultaneous prescription of thyroid extract.

The initial dose of thyroid in adults should be $\frac{1}{2}$ to 1 grain. If the patient does not lose weight satisfactorily, at a rate, that is, of 1 to 2 pounds a

week, the dose may be increased cautiously by $\frac{1}{2}$ to 1 grain at fortnightly intervals. An increase in the resting pulse rate, a loss of weight greater than 2 pounds a week, the occurrence of hyperpnoea on exertion or of palpitations, the appearance of symptoms of sympathetic overactivity such as flushing or sweating or an increased nervousness, are all danger signals. The thyroid should be stopped for a fortnight or until all signs of overdosage have disappeared, and begun again with a smaller dose. It is rarely necessary to prescribe more than 5 grains a day.

Thyroid extract should only be used in patients who can be kept under close observation. Though permanent hyperthyroidism is rarely produced by overdosage with thyroid, it has occurred (Bruun, 1945; Lous, 1945), and the possibility must always be kept in mind.

There is no necessity to use proprietary preparations of thyroid, the ordinary B.P. preparation of thyroideum siccum by any reliable maker satisfying all requirements. There is no point in using mixtures containing anterior pituitary extract, which is inactive by mouth and has no effect on obesity by any route; nor is there any need to give thyroid by injection. It is the only glandular extract which is fully effective when given orally.

Diuretics

Diuretics are of the greatest value, and many of the grosser cases of obesity can only be improved if they are used in conjunction with dietetic measures. Mersalyl has been used often, but the treatment of severe adiposity takes a long time and there are theoretical objections to its continued use. The most satisfactory drug in my hands has been urea, which can be taken harmlessly in large doses for an indefinite time. One heaped teaspoonful of the crystals (equal to approximately 5 grams) is taken three times a day. To many the taste is somewhat objectionable and I have found no effective way of masking it. Fruit juice appears to be the best vehicle.

Benzedrine Sulphate

Amphetamine or benzedrine sulphate is sometimes useful in reducing the appetite of those who find a rigid diet too difficult to follow. It should be given in doses of 5 to 10 mgm. in the early morning, so that sleep is not impaired. It is preferable to obtain the collaboration of the patient by more natural means.

Exercise

Exercise is helpful, but not as helpful as is often supposed. This is partly because a little food

makes up in energy for a lot of exercise, and partly because exercise often increases the appetite disproportionately. A Varsity crew usually puts on weight while in training, though it is true that the increase is in muscle, not in fat. On the other hand, the loss of salt in the sweat may help towards the desired negative water balance, provided that the patient does not drink more than he would if he were not taking exercise and does not replace the salt. In short, exercise is useful when the patient is prepared to stick to instructions, however hungry or thirsty the exercise may make him.

Turkish Baths

In a Turkish bath, rapid loss of weight occurs owing to sweating. The loss of salt is a great advantage if the patient is on a low salt diet, for he has no means of replacing it, and even if he drinks a large volume of fluid after his bath, the kidney excretes it in order to maintain the electrolyte concentration.

Anterior Pituitary Extracts

A so-called whole extract of the anterior pituitary is well spoken of in the advertisements of the less reputable hormone-mongers, but nowhere else. It is inactive when given by mouth, and ineffective when given by injection. There is neither rationale for its use nor evidence of its value.

There is, perhaps, some reason for the injection of the thyrotrophic factor, but it is much cheaper to give thyroid extract by mouth.

Oestrogenic Substances

These are often prescribed for those women who put on weight excessively at the menopause. The metabolic changes at this period have already been discussed. The period is one of general endocrine readjustment, and it is doubtful whether ovarian deficiency is directly the cause of the adiposity. On this insecure rationale, various preparations are prescribed. Ovarian extracts are inert by any route. The so-called "natural" pure oestrogens such as oestradiol, though mildly active by mouth and highly potent when given by injection, appear to have no value in reducing body weight, and their tendency to increase water-retention may even have the opposite effect. The synthetic oestrogens such as stilboestrol, are equally valueless for this purpose.

Dinitrophenol

This very powerful metabolic stimulant should never have been introduced to clinical medicine.

It is unfortunately highly effective and a small spate of deaths followed its first appearance in the medica¹ press.

Summary

The slimming of fat patients is important, not only for aesthetic and psychological reasons, but because adiposity predisposes to many more serious disorders and shortens the span of life.

The causes of adiposity remain obscure. Though many of the ductless glands have probably some effect in determining body weight, it is not at present possible to ascribe the fault accurately in most cases of human obesity. There is, however, good reason to guess that in many the hypothalamus and the posterior pituitary must bear the brunt of the blame. The fact that gross adiposity, and perhaps also milder degrees, show evidence not only of fat retention, but of water-retention, has been stressed too little in the past. When this fact is borne in mind, many apparent anomalies in the reaction to treatment are easily explained, and the results of treatment are far more satisfactory.

The law of the conservation of mass cannot be circumvented. The reduction of the diet below the energy requirements of the patient must eventually cause a loss of fat. That it does not always cause a great loss of weight is due to the disturbed water balance. A fat person is like a wet sponge and much of the excess weight is due to water rather than to fat.

For this reason a diet low in calories and preferably low in carbohydrate, salt and fluid should be supplemented by the use of diuretics, of which urea is the safest and most effective. Thyroid extract has a small place in treatment, and benzedrine may occasionally be helpful. Endocrine products other than thyroid extract are valueless, and other substances which increase the metabolic rate are too dangerous to use.

It is often difficult to persuade the patient that there is no comfortable road to normal proportions. *Il faut souffrir pour être belle.* If, however, co-operation can be obtained, excellent results are almost always to be obtained by dietetic and medical measures.

REFERENCES

- BAUER, J. (1933), *Med. Klin.*, **29**, 1297.
 BERNARD, C. (1895), *Leçons sur les propriétés physiologiques et les altérations pathologiques des liquides de l'organisme*, **1**, 297. Paris.
 BEZNAK, A. de (1938), *Presse Med.*, **46**, 996. W.W.
 BOOTHBY, W. M., and SANDIFORD, I. (1922), *J. Biol. Chem.*, **54**, 783.
 BRUUN, E. (1945), *Acta Med. Scand.*, **122**, 13.
 COOPE, R., and CHAMBERLAIN, E. N. (1925), *J. Physiol.*, **60**, 69.
 DOBREFF, M. (1926), *Pflügers Arch.*, **213**, 511.
 DUBLIN, L. I., and LOTKA, A. J. (1936), "Length of Life," New York.
 EAVES, E. C., and CROLL, M. M. (1930), *Brain*, **53**, 56.
 EMBLETON, D. (1938), *Brit. Med. Jour.*, **2**, 739.
 FALTA, W. (1923), "Endocrine Diseases," London.
 FISK, E. L. (1923), "Health Building and Life Extension."
 FÖLDES, E. (1933), "A New Approach to Dietetic Therapy," Boston.
 FROHLICH, A. (1901), *Wien, klin. Wchnschr.*, **15**, 883.
 GRAFE, E. (1933), "Metabolic Diseases and their Treatment," Philadelphia, New York.
 GRAFE, E., and GRAHAM, D. (1911), *Ztschr. f. physiol. Chem.*, **73**, 1.
 GRAFE, E., and KOCH, R. (1912), *Deutsches. Arch. f. klin. Med.*, **106**, 504.
 GOLDZIEHER, M. A. (1945), "The Adrenal Glands in Health and Disease," Philadelphia.
 HETHERINGTON, A. W., and RANSON, S. W. (1942).
 HETENY, G. (1936), *Deutsches. Arch. f. klin. Med.*, **179** 134.
 HUNTER, D. (1933), *Lancet*, **2**, 994.
 INGLE, E. J. (1943), *J. Clin. Endocrinol.*, **3**, 603.
 KLISIECKI, PICKFORD, ROTHCHILD, and VERNEY, E. n. (1933), *Proc. Royal Soc. B.*, **112**, 496.
 LAUTER, S. (1926), *Deutsches. Arch. f. Klin. Med.*, **150**, 315.
 LE MARQUAND, H. S., and TOZER, F. H. W. (1943).
 LOUS, P. (1945), *Acta Med. Scand.*, **122**, 83.
 LAWRENCE, R. D. (1946), *Lancet*, **2**, 724.
 MACKAY, E. M., and SHERRILL, J. W. (1941), *Endocrinology*, **28**, 518.
 MACKIE, E. M. (1923), *Proc. Royal Soc. B.*, **98**, 413.
 MACLAGEN, N. F. (1937), *J. Physiol.*, **90**, 385.
 NEVENSCHWANDER-LEMMER, N. (1936), *Ztschr. f. d. ges. exper. Med.*, **99**, 395.
 NEWBURGH, L. H. (1942), *Arch. Int. Med.*, **70**, 1033.
 O'CONNOR, W. J., and VERNEY, E. B. (1942), *Quart. J. Exper. Physiol.*, **31**, 393.
 PLUMMER, W. A. (1940), *Trans. Am. Ass.* "Study of Goitre," 88.
 RANSON, S. W., FISHER, C., and INGRAM, W. R. (1938), *Endocr.*, **23**, 175.
 ROWNTREE, L. G., and BRUNSTING, L. A. (1933), *Endocr.*, **17**, 377.
 RYDIN, H., and VERNEY, E. B. (1938), *Quart. J. Exper. Physiol.*, **27**, 243.
 SCHONDORFF, B. (1912), *Arch. f. d. ges. Physiol.*, **67**, 430.
 SMITH, P. E. (1929-30), "Relation of the Activity of the Pituitary and Thyroid Glands," Baltimore.
 STRONG, J. M., and McCLUGAGE, H. B. (1931), *Am. J. M. Sc.*, **182**, 49.
 STROUSE, S., WANG, C. C., and DYE, M. (1924), *Arch. Int. Med.*, **34**, 275.
 THEOBALD, G. W., and VERNEY, E. B. (1935), *J. Physiol.*, **83**, 341.
 VERZAR, F., and LASZT, L. (1936), *Bio. Chem. Ztschr.*, **288**, 356.
 VON ECONOMO, C. (1931), "Encephalitis Lethargica," London.
 VON GRAFE, E. (1933), "Metabolic Diseases and their Treatment," Philadelphia.
 WILEY, F. H., and NEWBURGH, L. H. (1931), *J. Clin. Invest.*, **10**, 733.
 WOHL, M. (1932), *Am. Therap. Soc.*, **31**, 211.
 YOUNG, F. G. (1941), *Brit. Med. J.*, **2**, 897.
 YOUNG, F. G., (1944), *Brit. Med. J.*, **2**, 717.
 ZONDEK, H. (1944), "Diseases of the Endocrine Glands," London.