EDITORIAL

London, February 1949

There are few fields in medicine in which our ideas have not, in recent years, been radically changed. From infant welfare throughout the whole range of medicine to the newly-developing science of geriatrics we have greatly advanced. If the mother's milk fails to make the infant thrive, as is increasingly the case in this country to our great regret, science provides substitutes. Maternal mortality rates show a marked and continuing fall. The treatment of most of the infective diseases has been profoundly modified by chemotherapy and those still resistant are under constant and scientific attack. To mention only two, how seldom now does the surgeon have to make those agonizing decisions as to whether an infected finger with lymphangitis and early signs of septicaemia shall be incised, whilst the role of the gonococcus has been so changed that it is now stated that the commonest form of urethral stricture in this country is that following the use of the indwelling catheter.

The tubercle bacillus we know to be susceptible to chemotherapy, but progress in this field has not been as rapid as had been hoped. Much work is being done on the subject as well as on that of the filterable viruses. For what we know of them we are largely indebted to our American colleagues. In 'Viral and Rickettsial Infections of Man' (Philadelphia; J. B. Lippincott, 1948) Rivers of the Rockefeller Institute for Medical Research has gathered all the known facts and observations on the subject. As in Pasteur's day, the work has also a direct bearing on agriculture, for crops of many kinds, from the humble potato to tobacco, are affected by destructive virus diseases. Electron micrographs of various types of virus are shown, that responsible for vaccinia being barrel-shaped, whilst the influenza virus, by the technique illustrated, appears as a small white ball. Others are filamentous. Such research requires complicated apparatus as well as teams of experts in the use of electron microscopes, ultracentrifuges, tissue cultures of various kinds for virus propagation and animal inoculations in-

volving monkeys as well as simpler types. This field the Americans have made particularly their own; all credit to them.

In the field of cancer a vast amount of work has been done both in the evaluation of known methods of treatment and in experimental study, whilst, in certain types, hormone therapy has given encouraging, though not curative results. Broadly speaking, however, advances in the treatment of cancer have been so far disappointing. We have as yet neither a test for its presence nor good evidence of its susceptibility to any form of chemotherapy.

As medicine proceeds further and further in such divergent lines we see more and more the tendency to specialization. As a result, from his earliest days the medical student is introduced to his life's study in a series of water-tight compartments between which he himself is left to establish his own equilibrium. For this his greatest asset is sound *common sense*. With the present-day keen competition for places in our medical schools, the onus on those responsible for the selection of the candidates to be admitted is very heavy. The boy who passes out top in his Higher Certificate examination may be by no means the most likely to make the best doctor. The very fact that he has passed his examination with such distinction may well indicate a lack of balance in his outlook, in that he may have given up his whole time to his books to the exclusion of other activities healthy for both mind and body. Such a boy may well make the most brilliant research scholar but he may have little background for success, in the best sense of the word, in general or consulting practice. In such a test his record throughout his school career, his masters' reports, his hobbies and above all his character should be given as much weight as his final examination result.

Granted the capacity to learn, we believe that the most important requisites for the doctor of tomorrow will still be the gifts of *sound common sense* and a *friendly understanding of his fellow men*.

ANNOTATION

Angiocardiography

The radiographic examination of the heart and great vessels following the intravenous injection of a radio-opaque substance is known as *angiocardiography*. Castellanos, Pereiras and Garcia first reported the extensive use of the procedure in infants, and Robb and Steinberg in adults. Sussman has performed angiocardiography on almost 1,500 cases without untoward reactions.

The radio-opaque material used is a 70 per cent. solution of diodrast, and quantities from 40-50 cm. may be injected intravenously in adults, with proportionately smaller doses regulated by age and weight in children. The injection must be given as rapidly as possible to ensure good delineation of the heart and great vessels. Following the injection, the patient may experience a feeling of heat, nausea and faintness which passes off within a few minutes in most cases. As a fall in blood pressure may follow the injection of diodrast, adrenalin should be at hand for use if required. Patients who might be adversely affected by a sudden fall in blood pressure should not be submitted to this procedure. Severe kidney or liver disease are thus contraindications, and to eliminate individual idiosyncrasy a small amount of diluted diodrast should be given intravenously as a test before angiocardiography is undertaken. Successive photographs can be taken on 70 mm. films, or separate exposures can be made once or twice per second on X-ray plates. Successful angiocardiography depends on a well-trained team of workers.

Angiocardiography is valuable to the clinician in that it presents a more accurate delineation of the heart and vessels than can be obtained from simple radiography. Following a successful injection, one can see successively the superior vena cava, the right auricle, the right ventricle, the pulmonary artery and branches, the pulmonary veins, the left auricle, left ventricle, the aorta and the branches arising from the aortic arch. It has been reported that the coronary arteries have been visualized in some patients.

Robb and Steinberg have discussed the alteration of the pulmonary vascular pattern in chest disease. They describe the displacement of the hilar vessels by enlarged glands, and have ob-

served these changes in both primary and secondary tumours of the mediastinum and in tuberculosis. In the latter disease, they describe three types of changes in the pulmonary circulation. These are diminished vascularity due to narrowing and obliteration of the vessels, displacement of the pulmonary artery by tuberculous adenitis, and severe degrees of displacement of the heart and great vessels by fibrosis. In some cases of tuberculosis and of chronic suppurative pulmonary conditions, areas of increased vascularity can be seen. Sussman reports the appearances in cases of neoplasia of the lung. Infiltrative tumours may cause partial or complete occlusion of the larger branches of the pulmonary artery, whereas circumscribed tumours, even though malignant, cause displacement and distortion, but not occlusion. Hence, angiocardiography may be of value in cases where bronchogenic carcinoma has been suspected but cannot be proven even by bronchoscopy.

Angiocardiography has proved of considerable value in the surgery of the heart. The site and extent of coarctation of the aorta can be shown, and the presence of a ductus arteriosus may be demonstrated by an alteration in the outline of the descending aorta, but patency is not necessarily proven thereby. If a large shunt is present, revisualization of the pulmonary artery occurs. In Fallot's tetralogy, the features of pulmonary stenosis and right ventricular hypertrophy are shown, and shunting of diodrast from the right ventricle into the overriding aorta and left ventricle can be seen. The procedure is also of use in confirming the clinical diagnosis of septal defects, and may be used to demonstrate acquired aneurysms of the aorta or vessels arising from its arch.

It is obvious that the use of angiocardiography in the diagnosis of congenital heart disease is extremely important. It may be invaluable in the consideration of operative measures. With the co-ordination of accurate history, clinical examination, electrocardiography, angiocardiography and cardiac catheterization, a complete diagnosis can now be made in many cases. With increased skill and improved methods, angiocardiography is likely to be an important diagnostic procedure in many cases of cardiac disease, and it may well prove of value in the diagnosis of pulmonary conditions.

E. ROHR.