- BEVAN, G., EVANS, R., PUGH, I., SUDDS, M. V. N., PARKER, M. T., and SLADDEN, A. F. S. (1948), *Lancet*, **1**, 545-550.
 BIGGER, J. N. (1946), *Lancet*, **1**, 81.
 BRADLEY, W. H. (1949), *Lancet*, **1**, 869.
 DESRANLAU, J. (1948), *Canad. J. Publ. Hlth.*, **39**, 317-319.
 DOUGLAS, A. D. M. (1949), *Lancet*, **2**, 105.
 FELIX, A. (1935), *Lancet*, **1**, 799.
 FRY, R. M., JONES, R. E., MOORE, B., PARKER, M. T., and THOMSON, S. (1948), *Brit. Med. J.*, **2**, 295-296.

- HODGSON, A. E. (1944), *Brit. Med. J.*, **2**, 339.
 MCSWEENEY, C. J. (1946), *Lancet*, **2**, 114-117.
 MCSWEENEY, C. J. (1948), *Ibid.*, **1**, 691-692.
 MURGATROYD, F. (1949), *Brit. Med. J.*, **1**, 851-852.
 PARSONS, C. G. (1948), *Lancet*, **1**, 510-513.
 WOODWARD, T. E., SMADEL, G. E., LEY, H. L., Jun., GREEN, R., and MANKIKAR, D. J. (1948), *Ann. Inter. Med.*, **29**, 1, 131-134.

CORRESPONDENCE

CARDIOSPASM

SIR,

I thoroughly enjoyed reading Mr. Rodney Maingot's article entitled 'The Surgery of Cardiospasm', which was published in the May 1949 number of this journal.

There are one or two observations which I should like to add regarding the treatment of this condition.

In a normal person, the gastric contents are prevented from regurgitating into the oesophagus by two main mechanisms :—

1. The action of the cardiac sphincter.
2. The 'pinchcock' action of the diaphragm.

Any operative procedure which is embarked upon must necessarily interfere with the first of these—but it is of the greatest importance to preserve the second, i.e. the pinchcock action of the diaphragm.

If both of these mechanisms are damaged by surgery, a patulous cardia results. This produces oesophagitis and eventually peptic ulceration of the oesophagus, which is a worse condition and is more difficult to deal with surgically than cardio-

spasm. The operation of oesophago-gastrostomy should only be mentioned in the treatment of cardiospasm to be condemned. It certainly relieves the obstruction at the cardia, but it creates a dangerous anomaly because there is nothing to prevent gastric juices regurgitating into the oesophagus. I know of at least one fatality from haemorrhage following this operation, which came from peptic digestion of the oesophageal wall.

In Leeds, a modified Heller's operation, as described by Mr. Maingot, has been performed on those cases which did not respond to dilatation with the Negus hydrostatic bag. This is a safe operation for it does not interfere with the pinchcock action, and post-operatively there has been no regurgitation into the oesophagus. It has given excellent results both symptomatically and radiologically in the 17 cases in which it has been performed. I agree with Mr. Maingot that it is the operation of choice. Other operative procedures altering the cardiac mechanism may prove to be dangerous.

Yours, etc.,

G. H. WOOLER, F.R.C.S.

Leeds.