HYPOPITUITARISM

By A. A. G. LEWIS, B.Sc., M.D., M.R.C.P.

*Physician to the Connaught Hospital and the Woodford Jubilee Hospital
Saltwell Research Scholar, Royal College of Physicians (From the Professional Medical Unit, Middlesex Hospital)*

Introduction

Many of the clinical features of hypopituitarism were described by Simpson in 1872 in his account of the symptoms associated with post-partum superinvolution of the uterus. The syndrome resulting from the simultaneous failure of several endocrine glands was described by Claude and Gougerot in 1907, but at this time the dominant role of the adenohypophysis in the endocrine system was not appreciated. Douglas Robertson (1951b) has recently drawn attention to the paper by Glinski (1913) describing the association of anterior pituitary necrosis with post-partum haemorrhage. Simmonds published his account of the necropsy of a patient suffering from atrophy of the anterior lobe in 1914. He did not at first realize the importance of the small size of the thyroid and adrenals, in fact he emphasized that their histology was normal. In subsequent papers he developed his conception of pituitary failure, though in the course of this he stressed unduly certain features of his cases such as cachexia and progeria. It was particularly unfortunate that in the title of two of his papers the word 'Kachexie' should have appeared, for this became inseparably associated for many years afterwards with the syndrome of panhypopituitarism.

In 1942 Escamilla and Lissner published an extensive review of the literature, collecting details of 101 cases of this condition. Their four criteria of severe pituitary deficiency (loss of weight, loss of sexual function, asthenia and a B.M.R. below -20 per cent.) ensured that at least one of Simmonds' original misconceptions would be perpetuated. Fraser and Smith in 1941 stated that the nutrition of these patients was usually good. This was further emphasized by Sheehan and Summers (1949) in their detailed description of the syndrome resulting from severe longstanding hypopituitarism. This account was based on 95 cases in which complete or nearly complete destruction of the anterior lobe was found at necropsy, and is therefore the most authoritative

available. It confirms and extends the description of the syndrome of panhypopituitarism arising from post-partum necrosis of the anterior lobe given by Sheehan in a series of papers. This particular condition is now frequently referred to as Sheehan's disease.

In recent years, as the recognition of the essential features of the complete syndrome became clearer, and as biochemical tests and hormone assays for the investigation of suspected cases came to be more widely used, more interest has been taken in the less severe degrees of pituitary failure and several papers have appeared describing such cases (Cooke and Sheehan, 1950; Oelbaum, 1952a). One result of this increased interest has been a clearer understanding of the natural history of the disease, many aspects of which have recently been discussed by Hubble (1952).

The Hormones of the Anterior Pituitary

The adenohypophysis produces at least six hormones: Follicle-stimulating (FSH), luteinising or interstitial-cell-stimulating (LH or ICSH), lactogenic or luteotropic, thyrotropic (TSH), adrenocorticotrophic (ACTH) and growth hormone.

There is increasing evidence that the adrenal cortex secretes more than one hormone and that Selye's original conception that the gland produces both glucocorticoids and mineralocorticoids was correct. Whether his hypothesis that the production of the latter is stimulated by a different pituitary hormone from the former is also true is not known, but it appears very improbable. It is still doubtful whether adrenal androgen production is stimulated by ACTH or by LH.

Anterior pituitary extracts also inhibit the action of hexokinase in catalyzing the transformation of glucose to glucose-6-phosphate, an inhibition which is opposed by insulin. It has been claimed (Li, 1952) that growth hormone antagonizes the action of insulin and that this may partly explain its diabetogenic activity in animals. Although this activity appears to be that of growth

hormone itself, the latter always produces growth in young animals, not diabetes (Young, 1952). The exact relationship between the growth hormone and the glycotropic action of the anterior pituitary is therefore not clearly known, but there is no doubt that the gland plays a direct part in the regulation of the blood sugar by opposing the peripheral action of insulin.

Panhypopituitarism

Pathology. In a large number of cases destruction of the anterior lobe follows a severe post-partum circulatory collapse. Sheehan and Murdoch (1938) thought that the blood flow to the anterior lobe was reduced at delivery, and that circulatory collapse would then favour thrombosis in the small vessels, with infarction and necrosis of much of the secretory tissue. They found a relationship between the degree of collapse and the frequency of necrosis in a series of cases; necrosis occurred in seven of the eight with the most severe degree of collapse. Both collapse and necrosis were found to be more frequent in multiparae. The necrosis may be very extensive, only small islets of healthy tissue remaining.

In other cases there may be scarring, fibrosis or cystic degeneration of the anterior lobe for which no definite cause can be found. Occasionally a head injury appears to have been responsible. Sometimes evidence is found of previous destructive lesions due to tuberculosis, gummatous necrosis, or other granulomata. Syphilis of the pituitary is a rare condition; the subject has recently been reviewed by Oelbaum (1952b). Bleisch and Robbins (1952) collected accounts of 50 cases with granulomata involving the anterior lobe, and added four of their own. In many of these cases there was other evidence of syphilis, tuberculosis or sarcoidosis. Occasionally trauma appeared to have been the cause, and the possibility of primary specific pituitary degeneration could not always be excluded. In some cases the sella turcica seems to be empty at post-mortem examination ('empty sella'). This appearance is probably produced by healing and extreme scarring of a granulomatous condition.

Other causes of hypopituitarism include surgical hypophysectomy and cysts and tumours, either in the sella or in its neighbourhood.

In 76 of the 95 cases collected by Sheehan and Summers, all the other endocrine glands were certainly or probably atrophied. Only in six of the 95 had at least one endocrine gland been definitely normal (endocrine atrophy was present in only half the cases of moderate pituitary damage). The adrenal glands were small; only the medulla and the zona fasciculata of the cortex remained. The thyroid was small and atrophic;

in 15 cases it was only represented by dense fibrous tissue. The female genitalia and ovaries were atrophic, the endometrium being merely a layer of cubical epithelium. No developing follicles or corpora lutea were seen in the ovaries. In males the testes were atrophic and the prostate and penis small. In spite of the fact that the weights of the patients were within the normal range, most of the viscera were small.

Clinical Features

In severe longstanding hypopituitarism all the features described by Sheehan and Summers (1949) are usually present. The most characteristic picture is seen in women who have recovered from post-partum haemorrhage complicated by pituitary necrosis. There is usually a history of very severe collapse following delivery or manual removal of the placenta, which required the transfusion of several pints of blood. Lactation was never established and menstruation and libido never returned. Afterwards the patient became dull and apathetic, mentally slow and sensitive to cold, with no interest in running the home, often retiring to bed or sitting in front of the fire for long periods of the day. Occasionally a psychosis has been suspected. The appearance and speech are characteristic, the patient having a forlorn, round-shouldered attitude and describing her symptoms in a dreary voice. The face is pale and flabby, sometimes with a slight yellow tint from carotinaemia. The mucosae are pale and the eyebrows thin. The body hair has disappeared apart from some remaining on the labia majora. The skin is thin and dry, and depigmentation, which gives it a wax-like appearance, is widespread. Even the areolae are pale. Erythema ab igne frequently occurs on the shins, but the areas are not pigmented. There is no tanning after exposure to sunlight. Pain in the knees and stiffness are common, and with the apathy and physical weakness may cause the patient to lie in bed almost as if suffering from a spastic paraplegia. Nocturia is the rule, and the patient may notice that she does not pass large volumes of urine after drinking during the day. Atrophy of the vagina may lead to severe dyspareunia. Sterility is almost invariable. There is an anaemia, often normochromic. Blood pressure may be normal or low. X-ray of the chest does not reveal the large heart of myxoedema, but the E.C.G. may show low voltage curves, sometimes with changes due to hyperkalaemia.

There has been a great deal of controversy and misunderstanding about the weight of these patients ever since Simmonds emphasized the occurrence of cachexia. Many of them look thin and ill, but they are not cachectic or emaciated in the

strict sense. Sheehan and Summers stressed that the body weight was not usually abnormal. Other authors consider that a moderate loss of weight is common. In reviewing the details of 35 cases of post-partum necrosis Farquharson (1950) found that only seven weighed less than 100 lb., five being described as 'thin or emaciated.' He also stated: 'It would appear that *complete* destruction of the gland is followed *at once* by loss of weight and other symptoms of the disease.' It seems probable that the apathy and loss of interest and of appetite found in the severe cases may lead to considerable weight loss at various times in the course of the illness. The case described by Aitken and Russell (1934), in which the pituitary was destroyed by haemorrhage into a chromophobe adenoma, lost 35 lb. in as many weeks after the haemorrhage, while the case of Farquharson and Graham (1931), with post-partum necrosis, gained 20 lb. when her appetite improved for some years before death. A complication such as tuberculosis, which is not uncommon, may also reduce the weight.

While post-partum necrosis of the gland is the commonest cause of pituitary failure in women, in men the syndrome results from fibrosis, granuloma or tumour. The onset is gradual, usually with loss of libido and of potentia. The external genitalia and prostate become small and the voice may become high-pitched. The patient rarely has to shave.

A number of these patients become comatose, often with little warning. In many cases this is due to hypoglycaemia, which may occur even after comparatively short periods of fasting or after the administration of insulin given to stimulate the appetite or as a diagnostic measure. Other cases are liable to crises similar to those of Addison's disease with diarrhoea, vomiting and collapse. Occasionally coma is associated with disturbances of body temperature, either with fever and tachycardia at the onset of an infection, to which the patient may readily succumb, or, rarely, with severe hypothermia. Sheehan and Summers (1952) have described one case in which the temperature had to be measured with a bacteriological thermometer, being as low as 87° F. The patient was pulseless and the heart beat slow and almost imperceptible. She was revived by immersion in a warm bath.

Laboratory Investigations

(1) *Urinary 17-ketosteroids.* Fraser and Smith (1941) found an excretion of less than 0.5 mg./24 hr. in nine of ten cases, the normal range in women for their laboratory being 5 to 15 mg./24 hr. Venning and Browne (1949) and Escamilla (1949) reported somewhat higher values, but in every case they

were below the normal range, being relatively lower in males.

(2) *Insulin tolerance test.* Details of this test are given by Fraser and Smith (1941). An attempt is made to assess two variables—the sensitivity to insulin and the response to the hypoglycaemia which insulin produces. All cases should be on a high carbohydrate diet for four days previously. After withdrawing blood for fasting blood sugar estimation, insulin is injected intravenously to reduce the blood sugar level by at least 45 per cent. Normally this may be achieved by a dose of 0.1 units/kg., but in suspected panhypopituitarism not more than one-third of this dose should be given as there may be great sensitivity to insulin. It is best to talk to the patient as clouding of consciousness may be the first indication of dangerously low blood sugar levels, in which case the test should be terminated with a glucose drink. Blood is taken for glucose estimations at 20, 30, 45, 60 and 120 minutes after the injection. Normally when the blood sugar level has been halved the response is rapid and the normal level is regained within two hours. There may be considerable delay in myxoedema and in Addison's disease, but in severe hypopituitarism the response may be almost absent, and this feature, together with the great sensitivity to insulin, is characteristic.

The test should not be performed if the fasting blood sugar is much below normal as it may be in cases of hyperpituitarism.

(3) *Gonadotropins.* The urinary excretion of FSH is greatly reduced (Klinefelter, Albright and Griswold, 1943; Escamilla, 1949). The test is based on the original method of Zondek (1930), depending on the enlargement of the uterus of immature mice by extracts of the urine injected after serial dilution. Extracts can be prepared either by alcohol precipitation or by ultracentrifugation.

(4) *Robinson-Power-Kepler test.* This empirical test depends upon the summation of the anomalies of renal function found in adrenal deficiency. These are: (a) Lowered renal blood flow and glomerular filtration rate with consequent lowered urea clearance and raised blood urea. In panhypopituitarism this is partly due to the deficiency of growth hormone which is probably responsible for the microsplanchnia. The average weight of the two kidneys in Sheehan and Summers's cases was 170 g. (normal 300 g.); (b) failure of water diuresis in response to a water load, with nocturia. The cause of this is uncertain, but appears to be due to loss of tubular selectivity, the patient being unable to pass large volumes of dilute urine, or small volumes of very concentrated urine, in response to the appropriate stimuli. Severe cases pass small volumes of urine with a

chloride concentration similar to that of the plasma, irrespective of hydration or dehydration, during the day (Chalmers and Lewis, 1951). At night large volumes of less concentrated urine may be passed (Robertson, 1951a).

In this test a water load of 20 ml./kg. is given to the fasting patient in the morning, after the night urine specimen has been collected. Further specimens are collected at hourly intervals. In the patient with adrenal deficiency none of these specimens will be larger than that collected overnight because of the nocturia and failure of diuresis in this condition, but this anomaly is not diagnostic. In the second part of the test the following index is derived:

$$\frac{\text{Vol. of largest day urine} \times \text{Plasma chloride}}{\text{Vol. of night urine} \times \text{Urinary chloride}} = \frac{\text{Urinary urea}}{\text{Blood urea}}$$

In adrenal deficiency the factors of the numerator tend to be lower than in normals, while those of the denominator are greater, so that the figure derived is a low one. Robertson, Power and Kepler (1941) found that the factor was below 25 in adrenal deficiency and above 30 in normals, though occasionally cases with cachexia or with hepatic or renal failure gave anomalous results.

(5) *B.M.R.* This is abnormally low, usually below -30. The serum cholesterol may be raised, but is not usually in the range found in primary myxoedema.

(6) *Plasma electrolytes.* The sodium and chloride levels may be low and that of potassium high, but the figures are often normal.

(7) *Response to the injection of ACTH.* It may take several injections of the hormone to elicit a full adrenal response, which may therefore be less at first than that found in normals but greater than that in Addison's disease. It has been suggested (McIntosh *et al.*, 1950) that this incomplete response may be valuable in diagnosis and that a clear distinction can be made between the rises in 17-ketosteroid excretion in patients with hypopituitarism and in normals after ACTH. However, this perhaps overestimates the significance of minor variations in the results of a complicated chemical analysis.

Incomplete Lesions

Detailed study of the patients suffering from post-partum necrosis of the anterior lobe has led to an appreciation of the true nature of the syndrome of hypopituitarism. Where haemorrhage and collapse in the puerperium have been followed by the appearance of less complete degrees of deficiency the definite time of onset fixed by the clinical history has enabled the evolution of the disease to be followed from the beginning. Several

cases have been recorded where quite a small surviving remnant of the gland has been found post-mortem and has evidently prevented the occurrence of any of the symptoms of the syndrome. This is analogous to the finding that only a small part of the secreting tissue of the neurohypophysis may prevent the occurrence of diabetes insipidus. In some cases the complete syndrome of panhypopituitarism may only develop after many years. Hubble (1952) describes a case in which the pubic hair disappeared and amenorrhoea developed after a post-partum haemorrhage, but adrenal and thyroid failure were delayed for 25 years. This emphasizes the importance of Sheehan and Summers's description of the complete syndrome, for they were careful to base this only on the study of longstanding severe cases.

In many other cases the pituitary failure may never be more than partial. Sheehan and Murdoch (1938) reviewed the histories of 128 women who had suffered from post-partum haemorrhage and collapse between 1930 and 1936. Absence of any mammary reaction was the commonest sign of pituitary failure and occurred in 44 cases. Three of these were in the group with the least degree of collapse and had no other symptoms. Menstruation was absent or infrequent in 32 cases, while body hair was lost in only 11. Failure of lactation and of sexual function are therefore the commonest symptoms, but the second of these is not invariable. Several cases have been reported in which extensive destruction of the gland has been found post-mortem, but menstruation and even pregnancy had occurred after the onset of the disease. Bishop and de Mowbray (1951) have described a patient who developed symptoms of hypopituitarism with almost complete amenorrhoea after a pregnancy at the age of 22. She twice became pregnant subsequently, going to term on the second occasion and improving considerably during this time, but relapsing after the confinement.

Oelbaum (1952a) emphasizes the variability of endocrine dysfunction in post-partum hypopituitarism. One case showed only a deficiency of gonadotropin excretion; amenorrhoea was complete, but thyroid and adrenocortical function were considered to be normal. The urinary 17-ketosteroid excretion was 7.0 mg./24 hr., and the insulin tolerance test normal. Another case had a normal gonadotropin excretion but was found to have an infantile uterus at a subsequent laparotomy. Cooke and Sheehan (1950) stress that patients may suffer from genital superinvolution, deficiency of thyroid or adrenal cortical function or hypoglycaemia or from any combination of these, and believe that the irregularities depend on the amount of the pituitary destroyed and the

time which has elapsed since destruction. One of their cases menstruated for six years before the full syndrome developed. Another developed amenorrhoea and genital atrophy after her only pregnancy *but breast-fed the baby for three months*.

Paschkis and Cantarow (1951) reported one case with normal 17-ketosteroid excretion but hypoglycaemia unresponsiveness, and four with the converse. These results favour the hypothesis that adrenal androgen production is controlled by the secretion of LH rather than the suggestion, made by the authors, that the adrenal functions may be individually suppressed.

In 1940 Means, Hertz and Lerman described the syndrome of 'pituitary myxoedema.' This is essentially a condition of hypopituitarism in which the predominant features are those of thyroid deficiency. Unless carefully sought, signs of deficiency of other pituitary hormones may not be found and the case may be mistaken for one of primary myxoedema. It must be borne in mind that while hypopituitarism frequently occurs without much loss of weight, any such loss in the course of the disease is strongly against the diagnosis of primary myxoedema. It is important to recognize such cases because they may react in an anomalous manner to thyroid. Occasionally small doses may precipitate an adrenal crisis, presumably by suddenly increasing the demand for adrenal hormones. On the other hand some cases are resistant to thyroid and show little improvement with doses effective in primary myxoedema.

Chromophobe adenomata frequently cause progressive hypopituitarism. Here again in the great majority of cases sexual failure occurs first. Amenorrhoea occurs in women, without menopausal symptoms, and in men there is loss of libido and potentia. Dott, Bailey and Cushing (1925) emphasized the importance of these symptoms early in the course of pituitary destruction by chromophobe tumours. In many cases there is little evidence of other deficiencies for many years, and it is possible for enlargement of the sella and optic atrophy to occur with normal pituitary function.

Pituitary Dwarfism

In this condition, hypopituitarism in childhood manifests itself mainly in failure of growth. The child is small but normally proportioned. Usually sexual maturation never occurs so that a very slow rate of growth may be maintained for many years. The signs of thyroid and adrenal deficiency are not usually evident. Sometimes the cause is a pituitary tumour such as a craniopharyngioma. In other cases there appears to be an isolated deficiency of growth hormone production. Hewer (1944) reported the necropsy of a female dwarf who

had had one child. Of 2,000 cells in the anterior pituitary only 0.2 per cent. were eosinophils.

Occasionally cases occur with isolated deficiencies of gonadotropic or of thyrotropic hormones (Hubble, 1952).

Hypopituitarism Secondary to Nutritional and Psychogenic Factors

In a recent stimulating review of the course of anterior hypopituitarism Hubble (1952) has discussed the failure of sexual function and growth that may occur in such conditions as anorexia nervosa and coeliac disease. While amenorrhoea in anorexia nervosa is frequently psychogenic, there is reason to believe that in some cases it is due to the malnutrition. It is known that in rats a deficient diet may depress sexual function before restricting growth. Banks, Lloyd and Scowen (1952) reported that severe diet restriction in two women after the menopause reduced the excretion of FSH to zero in a few weeks. There is usually no other evidence of pituitary deficiency in anorexia nervosa, though the B.M.R. is usually low. Thorn *et al.* (1950) observed a low 17-ketosteroid excretion and considered that this was evidence of impaired adrenal function, but other laboratory investigations such as the insulin tolerance test and the Robinson-Power-Kepler test are normal.

Dwarfism is usual when coeliac disease persists into adult life, in fact, if skeletal development is retarded by more than a few years it usually persists, at least partially, even when the nutritional deficiencies are completely corrected. One of Hubble's cases was still growing in his thirties, so that production of sex hormones (which ultimately limit growth by causing fusion of the epiphyses) was evidently more depressed than that of growth hormone.

Differential Diagnosis

Anorexia nervosa. This is the condition which is most frequently mistaken for Simmonds' disease, and there is no doubt that descriptions of the latter in the past have been vitiated by reports of cases in which these two conditions have been confused. The patient with anorexia nervosa in an advanced stage is emaciated, but in spite of this her mental and physical activity is often quite remarkable, whereas in panhypopituitarism the patient is dull, apathetic and inert even when the weight is normal. In anorexia nervosa amenorrhoea occurs early but body hair is not lost. Apart from a low B.M.R. the other laboratory investigations are generally normal.

Addison's disease. Body hair is often absent and the urinary excretion of 17-ketosteroids is low in women with Addison's disease, presumably be-

cause of the failure of production of adrenal androgens, but in hypopituitarism the characteristic pigmentation of Addison's disease is not seen and low blood pressure with low serum levels of sodium and chloride and high potassium occur much less often. As Thorn *et al.* (1949), have pointed out, the adrenal deficiency of hypopituitarism appears to lead predominantly to hypoglycaemia rather than to hypochloreaemia. In the insulin tolerance test the patient with Addison's disease may be less rather than more sensitive to insulin than normal (possibly because of increased glycotropic activity of the anterior pituitary), though there is hypoglycaemia unresponsiveness (Fraser, Albright and Smith, 1941).

Myxoedema. The importance of distinguishing between primary myxoedema and so-called pituitary myxoedema has been emphasized above. While the latter type rarely occurs without sexual failure, as Hubble has stressed, reports in the literature of cases in which menstruation and pregnancy have occurred in undoubted hypopituitarism do not make even this distinction completely reliable. In primary myxoedema urinary 17-ketosteroid excretion may be very low and the blood sugar may not return to normal in the insulin tolerance test. A distinction between these two conditions may therefore be very difficult to make. A detailed inquiry into the sexual function, menstrual rhythm and obstetric history, careful examination of the fundi and of the visual fields, radiological examination of the pituitary fossa and investigation of adrenal function may therefore be necessary in a doubtful case. The difficulty of making this distinction again emphasizes the importance of starting thyroid therapy with small doses of the hormone and the dangers of giving thyroid indiscriminately to any patient.

Primary ovarian insufficiency (Albright, Smith and Fraser, 1942). This is sometimes mistaken for hypopituitarism. There may be other stigmata such as webbing of the neck, coarctation of the aorta or cubitus valgus. Body hair is scanty but not absent, and the breasts are small. Sexual functions have never matured and the body proportions are eunuchoid, the span being greater than the height. These patients are not dwarfs but are usually moderately stunted, the height often being about 4 ft. 8 in. Assay of the urinary FSH excretion clinches the diagnosis for it is high in this condition.

Other mistaken diagnoses, commonly made on account of the appearance, are those of anaemia or nephritis, the latter particularly when nocturia is a prominent symptom. 'Post-partum debility,' neurosis and various psychoses may also be diagnosed.

Treatment

Sheehan and Murdoch (1938) reported a case in which improvement occurred during pregnancy following previous post-partum necrosis, and it was thought that some hyperplasia of the remnant of the anterior lobe had occurred. The case of Bishop and de Mowbray improved in pregnancy but relapsed later. The patients reported by Jakob (1923) and Farquharson, Belt and Duff (1938) became worse after pregnancy. It appears, therefore, that the chances of any real improvement occurring as the result of pregnancy, even in the unlikely event of conception taking place after post-partum necrosis, are extremely small.

Pituitary failure is not an indication for surgery in cases with cysts or tumours in the neighbourhood of the sella; in the great majority of cases surgery is undertaken to save the sight. Operation may, in fact, abolish what little endocrine function remains, with a fatal outcome (Grant, 1948). Caughey, James and Macleod (1952) reported four cases in which prolonged stupor developed after operations on pituitary tumours. Recovery occurred in two on treatment with cortisone.

In the great majority of cases the only possible treatment is some form of hormonal replacement therapy. A course of ACTH (25 mg. intramuscularly, six-hourly for a few days) brings about a dramatic improvement in those cases with symptoms of adrenal deficiency. Increased appetite, activity and libido, and euphoria and gain of weight occur, and this improvement may last for weeks after the end of the course (Rolland and Matthews, 1952; Summers and Sheehan, 1951; Hart, 1951). A normal diuretic response to water may be obtained (Chalmers and Lewis, 1951), and the nocturia may lessen. Hypoglycaemia may be prevented. Cortisone (25 mg. a day by mouth) brings about a similar improvement and is the best maintenance therapy. Either ACTH or cortisone should be given whenever surgery becomes necessary. Treatment of the adrenal deficiency with DOCA and extra salt give much less dramatic results than in cases of Addison's disease, though it may be beneficial.

Thyroid should only be given with extreme caution, starting with $\frac{1}{8}$ gr. daily while the patient is under observation. The simultaneous administration of ACTH, DOCA or extra salt reduce the risk of precipitating an adrenal crisis. In some cases the dose of thyroid can be greatly increased and the effects may be striking, particularly on the lethargy and the mental changes.

Injections of growth hormone are ineffective in pituitary dwarfism. Growth may be stimulated by androgen therapy but will cease when this

brings about fusion of the epiphyses; 500-1,000 i.u. of chorionic gonadotropin twice a week may stimulate growth and sexual function in these cases.

Adult females with hypogonadism can be given oestrogens with benefit.

In all cases testosterone therapy is beneficial, often strikingly so. In addition to stimulating a sense of wellbeing and libido, it brings about increased protein anabolism with a positive nitrogen balance and weight gain in both sexes. Muscular development and vigour increase and, in males, the penis enlarges and erections become frequent. It is claimed that spermatogenesis may be stimulated (Hurxthal, Bruns and Musulin, 1949). Body hair is increased. These actions have been discussed by Oelbaum (1952b). Methyl testosterone can be given sublingually, testosterone propionate can be given by injection or, most conveniently, 200 mg. of testosterone can be implanted.

Crisis and coma call for immediate and vigorous treatment. As hypoglycaemia is the commonest single cause large quantities of glucose must be given by mouth or intravenously. ACTH and cortisone can be given simultaneously but their effect will not be felt for some hours, so that diarrhoea and vomiting call for intravenous saline

and cortical extract (preferably lipo-adrenal cortex if this is available, as this contains more glucocorticoid fraction). DOCA (5 or 10 mg.) should be injected intramuscularly at the same time, and as infection is so frequently the cause of the collapse, a large intramuscular dose of penicillin or intravenous aureomycin should be given.

The treatment of pituitary coma has recently been fully discussed by Sheehan and Summers (1952). They observed no improvement following intravenous cortisone, saline or DOCA. They emphasized the importance of recognizing hypothermic coma. This occurred in one of their patients in spite of previous injections of DOCA, recovery occurred after immersion in a warm bath.

One of the dangers of hypopituitarism lies in the fact that the patient may be so apathetic that she fails to realize when her condition has deteriorated and a doctor may not be called until a crisis or coma has developed. Death may follow with appalling suddenness in such cases in spite of all efforts on the part of the physician. The prognosis in any severe case of hypopituitarism is therefore always grave and never exactly predictable. The patient must be kept under regular observation and must be told to report at once if any infection develops.

BIBLIOGRAPHY

- AITKEN, R. S., and RUSSELL, DOROTHY (1934), *Lancet*, ii, 802.
- ALBRIGHT, F., SMITH, P. H., and FRASER, R. W. (1942), *Amer. J. med. Sci.*, 204, 625.
- BANKS, P. J., LLOYD, H. M., and SCOWEN, E. F. (1952), 'Ciba Foundation Colloquia on Endocrinology', Vol. 4, p. 289. London, J. & A. Churchill.
- BISHOP, P. M. F., and DE MOWBRAY, R. R. (1951), *Proc. R. Soc. Med.*, 44, 163.
- BLEISCH, V. R., and ROBBINS, S. L. (1952), *Arch. intern. Med.*, 89, 877.
- CAUGHEY, J. E., JAMES, A., and MACLEOD, E. K. (1952), *Brit. med. J.*, i, 1216.
- CHALMERS, T. M., and LEWIS, A. A. G. (1951), *Lancet*, ii, 1158.
- CLAUDE, H., and GOUGEROT, H. (1907), *C. r. soc. Biol.*, 63, 785.
- COOKE, R. T., and SHEEHAN, H. L. (1950), *Brit. med. J.*, i, 928.
- DOTT, N. M., BAILEY, P., and CUSHING, H. (1925), *Brit. J. Surg.*, 13, 314.
- ESCAMILLA, R. F. (1949), *Ann. intern. Med.*, 30, 249.
- ESCAMILLA, R. F., and LISSER, H. (1942), *J. clin. Endocrinol.*, 2, 65.
- FARQUHARSON, R. F. (1950), 'Simmonds' Disease.' Springfield, C. C. Thomas.
- FARQUHARSON, R. F., and GRAHAM, D. (1931), *Tr. Ass. Amer. Phys.*, 46, 150.
- FARQUHARSON, R. F., BELT, T. H., and DUFF, G. L. (1938), *Tr. Amer. Clin. and Climatol. A.*, 54, 106.
- FRASER, R. W., and SMITH, P. H. (1941), *Quart. J. Med.*, 10, 207.
- FRASER, R. W., ALBRIGHT, F., and SMITH, P. H. (1941), *J. clin. Endocrinol.*, 1, 297.
- GLINSKI, L. K. (1913), *Przegl. Lek.*, 4, 13.
- GRANT, F. (1948), *J. Amer. med. Ass.*, 136, 668.
- HART, F. D. (1951), *Brit. med. J.*, ii, 795.
- HEWER, T. F. (1944), *J. Endocrinol.*, 3, 397.
- HUBBLE, D. (1952), *Lancet*, i, 1123.
- HURXTHAL, L. M., BRUNS, H. J., and MUSULIN, N. (1949), *J. clin. Endocrinol.*, 9, 1245.
- JAKOB, A. (1923-4), *Virchow's Arch.*, 246-7, 151.
- KLINEFELTER, H. F., ALBRIGHT, F., and GRISWOLD, G. C. (1943), *J. clin. Endocrinol.*, 3, 529.
- LI, C. H. (1952), 'Ciba Foundation Colloquia on Endocrinology', Vol. 4, p. 260. London, J. & A. Churchill.
- MCINTOSH, H. W., MCALPINE, H. T., SINGER, B., and HOFFMAN, M. M. (1950), 'Proceedings of the First ACTH Conference', p. 20. London, J. & A. Churchill.
- MEANS, J. H., HERTZ, S., and LERMAN, J. (1940), *Tr. Ass. Amer. Phys.*, 55, 32.
- OELBAUM, M. H. (1952a), *Brit. med. J.*, ii, 110.
- OELBAUM, M. H. (1952b), *Quart. J. Med.*, 21, 249.
- PASCHKIS, K. E., and CANTAROW, A. (1951), *Ann. intern. Med.*, 34, 669.
- ROBERTSON, J. D. (1951a), *Lancet*, ii, 282.
- ROBERTSON, J. D. (1951b), *Brit. med. J.*, i, 921.
- ROBINSON, F. J., POWER, M. H., and KEPLER, A. J. (1941), *Proc. Mayo Clin.*, 16, 577.
- ROLLAND, C. F., and MATTHEWS, J. D. (1952), *Brit. med. J.*, i, 1220.
- SHEEHAN, H. L., and MURDOCH, R. (1938), *J. Obst. Gynaec. Brit. Emp.*, 45, 456.
- SHEEHAN, H. L., and SUMMERS, V. K. (1949), *Quart. J. Med.*, 18, 319.
- SHEEHAN, H. L., and SUMMERS, V. K. (1952), *Brit. med. J.*, i, 1214.
- SIMMONDS, M. (1914), *Dtsch. Med. Wschr.*, 40, 322.
- SIMPSON, J. Y. (1872), 'Clinical Lectures on Diseases of Women.' Edinburgh (quoted by Sheehan and Murdoch).
- SUMMERS, V. K., and SHEEHAN, H. L. (1951), *Brit. med. J.*, ii, 564.
- THORN, G. W., FORSHAM, P. H., FRAWLEY, T. F., HILL, S. R., ROCHE, M., STAEHELIN, D., and WILSON, D. L. (1950), *New Engl. J. Med.*, 242, 783.
- THORN, G. W., FORSHAM, P. H., PRUNTY, F. T. G., BERGNER, G. E., and HILLS, H. E. (1949), *Ann. N.Y. Acad. Sci.*, 50, 646.
- VENNING, E. H., and BROWNE, J. S. L. (1947), *J. clin. Endocrinol.*, 7, 74.
- YOUNG, F. G. (1952), 'Ciba Foundation Colloquia on Endocrinology', Vol. 4, p. 255. London, J. & A. Churchill.
- ZONDEK, B. (1930), *Klin. Wschr.*, 91, 1207.