

in contrast with other antihypertensive drugs

# VASOLASTINE

is free from side-effects.

***It is non-toxic.***

In addition it resolves atheromatous deposits, normalises cholesterol and phospholipid values, and restores elasticity of arteries and capillaries.

# VASOLASTINE

The MODERN CAUSAL ENZYME THERAPY  
for hypertension and cardiovascular diseases



Manufactured by Enzypharm C. V. Holland

Further information and list of publications  
gladly sent on request by the Sole U.K. Distributors:

**F.A.I.R. LABORATORIES LTD.; TWICKENHAM MIDDLESEX**


**FORMULA per ampoule:**

Enzymes of the lipid	
metabolism	30 mg. ≡ 9000 units
Amin-oxidase	20 mg. ≡ 4000 units
Tyrosinase	20 mg. ≡ 4000 units
Stabilizing solvent	to 2 ml.

**Presentation:**

Boxes of 6, 25 and 100 ampoules.

# Salazopyrin<sup>®</sup> the proven therapy in ulcerative colitis now available as a



In distal proctocolitis, stump proctitis, and as an adjuvant to oral treatment in ulcerative colitis with severe manifestations in the rectum and rectosigmoid, the new *suppository form* of Salazopyrin is a logical therapy. Indeed, because of the well-attested dangers of steroid treatment <sup>1</sup> Salazopyrin may often be the *only* logical therapy.

<sup>1</sup> *Lancet* 1965, 1, 188.

Salazopyrin (sulphasalazine) is available in suppository form (0.5G) as well as plain 0.5G tablets and 0.5G EN-tabs (enteric coated tablets).

Suppository dosage: 1 to 2 suppositories in the morning immediately after bowel action, repeated in the evening.

Precautions: Salazopyrin should be administered under constant medical supervision, including periodic blood examinations. If serious symptoms occur, such as, leukopenia or sensitisation, the drug must be discontinued. There is no specific antidote for Salazopyrin. Teratogenic effects have never been reported.

Contra-indications: A history of marked sensitivity to sulphonamides.

Further information is available on request from:

 PHARMACIA  
(GREAT BRITAIN) LTD

Paramount House, 75 Uxbridge Road, Ealing, London, W.5. Tel: 01-579 0102/7