Everything I have said, however, is immediately invalidated if it is obvious clinically and on scientific grounds of respiratory function tests, that the patient's respiratory or cardiac reserve is not adequate to withstand major chest surgery, or that, by the inevitable resection of normally functioning lung tissue during the surgical treatment of lung cancer, the patient will be turned into a respiratory cripple.

Finally, I will leave you with a few statistics. Only about 20% of all lung cancer patients are suitable for surgical assessment. The overall operative

mortality is about 3%. The post-operative survival overall is 34.8% at 3 years, 27.3% at 5 years and 16.6% at 10 years. If the lesion is confined to the lung the 3 years survival after pneumonectomy is 48% and after lobectomy is 65%. If, however, the disease has spread beyond the lung the 3 year survival after pneumonectomy plummets to 17.9% and that after lobectomy is halved—at 32.2%.

Lung cancer is a very difficult disease to treat successfully and from these figures it will be apparent that surgery has a limited part to play in its treatment.

Discussion

SESSION 3

Chairman: DR M. W. McNICOL

(Physician, Central Middlesex Hospital and Willesden Chest Clinic)

DR F. H. SCADDING (Brompton Hospital): One of the hazards of aspiration lung biopsy frequently seen is implantation secondaries in the needle tract. This does not matter very much when there is no question of surgery. What is the risk of aspiration biopsy? I tend to avoid this if the case seems otherwise operable.

MISS SHEPHERD: Yes, I think this is a very real risk if the lesion proves to be a neoplasm. A case comes to mind of a lady of some 40 years who had a peripheral right upper lobe lesion and it was difficult to establish its pathology. She was not a suitable candidate for surgery, and the social circumstances were such that it would be extremely valuable to know whether this was a neoplasm or not. Fortunately for her I did a drill biopsy and it turned out to be tuberculous, and in this particular case I do agree there is a risk of implantation.

DR SCADDING: You are saying it is better not to do it if the case seems otherwise operable?

MISS SHEPHERD: One could certainly put a very strong argument on this line. The other problem, though, in the case of a mesothelioma which is not quite a lung carcinoma, is that again, if it is necessary to make a diagnosis, it is far safer to do a drill biopsy than do an open pleural biopsy. One has to weigh it up.

DR T. DEELEY (Velindre Hospital, Cardiff): I wonder if I might answer Dr Scadding's question also. I have now done a considerable number of drill biopsies of the lung and I have no evidence at all of aspiration seedlings. I have done some 4000 biopsies of malignant tumours and I think only one, a carcinoma of the breast which was just under the skin, grew through the drill biopsy.

DR McNICOL: Seeing this has been raised, has any one else experience of this?

MR J. W. JACKSON (Gerrards Cross, Bucks): I have never had any evidence of recurrence of tumour along the tract of the drill but I have had evidence of tumour in the tract of the biopsy using the Abram's needle and I have also had tubercle coming out of the tract after an Abram's needle biopsy, but I have never had either when using the high speed drill.

MISS SHEPHERD: When Dr Scadding asked the question actually he used the phrase aspiration needle biopsy. When I was speaking of drill biopsy I referred to the high speed drill. I wonder whether this is a significant factor?

MR JACKSON: Several radiologists are using this technique now, and using it more and more, with a simple needle, not the Abram's needle.

MISS SHEPHERD: I was meaning the Steele high speed drill. This is the type of biopsy needle which I would use in this particular instance, and I think Dr Deeley uses it also.

DR DEELEY: I was talking about the high speed drill which is our own drill rather than the Steele drill which is a larger bore.

DR McNICOL: Can we clarify this? We are talking about high speed drill biopsy, and so far no-one appears to have any evidence of implantation along the drill tract. Now, I think you were going to say something about this, Dr Steele, is that right?

DR S. J. STEELE (London Chest Hospital): I have done 300 biopsies, not all tumours it is true, but a good proportion of those have been tumours, and I have never had any spread along the trace using a high speed drill with a larger trephine than the one Dr Deeley uses, and I also know that the man probably with the widest experience of this in aspiration biopsy is Nordenstrom of Sweden. He has done well over 1000 cases and has had no definite spread along the tract. He has published that and also confirmed that more recently when I spoke to him. But I think in actual fact if you search the literature and actual recorded cases which have spread along the tract are very few indeed.

DR McNICOL: Thank you. Any other comments? Miss Shepherd, I was astonished at your figure of operability. Did I hear you say 20% or 30% of cases?

MISS SHEPHERD: Of these cases presented for assessment for surgery only about 20% are proved to be operable. This indicates, taking the disease as a whole, how limited surgery is.

DR McNICOL: Miss Shepherd has pointed out quite rightly how bad are the results of surgery. This must mean to say we are not selecting the right cases. Too many of the patients operated upon in fact have got extra pulmonary spread. Would it be reasonable, therefore, to use mediastinoscopy routinely in any patient in whom you think surgery might be applicable? Would this not improve the results and prevent patients being operated on unnecessarily?

MISS SHEPHERD: I think the first point is that mediastinoscopy is not an exhaustive search of all mediastinal glands. The only way one can do that is to do a block dissection of them all. Therefore, one has perhaps a certain percentage, and I have not got the figures, of false negatives. Now, if one does not operate on the patients, then what? I think perhaps this would be a point worth discussing when the other speakers have spoken later and one can discuss it more generally. Naturally, a surgeon believes that if he can obliterate a tumour then he is doing the right thing. Whatever method of treatment is used I think the figures are not particularly good. It is a very difficult disease to treat whichever way you handle it. That is my opinion. Would you agree with that, Dr Deeley?

DR McNICOL: This is an interesting point. Is there anyone here who has experience of universal use of mediastinoscopy in the screening process? Obviously it is a theoretical possibility. Is any one doing it or is it being done on any of their patients?

DR SCADDING: I know that Mr Belcher has been doing it at the Middlesex for over a year, and the others at the Middlesex, and he was telling me the other day that he has avoided, or been able to avoid, twelve thoracotomies, cases he would otherwise operate on.

DR McNICOL: That is twelve out of what sort of numbers?

DR SCADDING: I don't know how many. They were not all my cases.

DR McNICOL: Miss Shepherd, do you know the figures?

MISS SHEPHERD: I am afraid I don't know his figures, but I feel that you have got to be very careful when you say you have saved a thoracotomy. What I think we need is a continuous series of paratracheal gland involvement proved, followed by thoracotomy and regional gland dissection before we can get some idea of whether we are right not to operate on these people with paratracheal gland involvement. I think it is right provided there is no proximal bronchial tree involvement, and I think it is right to operate on a patient with a sub-corinial bland involvement if you can get the regional nodes blocked out. Provided the bronchial stump is clear I think one is justified in doing it.

DR J. MIKHAIL (Central Middlesex Hospital): Do you agree that mediastinoscopy's value is mainly in removing paratracheal glands and not hilar glands. As you were saying, those with involvement of the hilar glands should still go to thoracotomy. One has only to do a limited mediastinoscopy and perhaps only remove a paratracheal gland because involvement of those glands are the only indication that it is inoperable.

MISS SHEPHERD: This I confess at the present time is how I utilize mediastinoscopy in the assessment of bronchogenic carcinoma.

DR W. GRIFFEL (Banbury): Mediastinoscopy is an old matter brought into use recently and certainly helps to exclude positive cases, with the limitation which you put on it. We do not know whether we are refusing cases which may still be operable, but we must not forget that a poor prognosis is not directly dependent on what we see by a mediastinoscope or X-ray. Undiscovered secondaries in the brain, liver and everywhere else are the things which make the prognosis so bad. Not the resection of the tumour, because those cases where you have succeeded by lobectomy to remove the tumour with no evidence of secondaries, may live for 20-25 years.

MR JACKSON: Just one thing I think we should always remember is that Grey Turner used to say that no tumour should be deemed inoperable until it had been proved so by operation. The ultimate test of inoperability is operation, and I think that whereas mediastinoscopy and these other tests may guide us on our way, there are many patients, and many of them have been doctors, who prefer to have a thoracotomy before they have been deemed inoperable. We must consider it a black mark to have done thoracotomy and found a tumour inoperable, but I would be the first to admit that I have regretted on three occasions this year not having done a mediastinoscopy on a patient.

DR N. MACDONALD (Clare Hall Hospital): Could I ask Miss Shepherd two questions. One is about the place of bronchoscopy in the diagnosis. It seems to me there are far too many unthinking bronchoscopies done on patients with a respiratory infection, the patient brings up a little blood, etc., and there are a good many patients who are subjected to an unnecessary hazard. Do you not think bronchoscopy is done perhaps rather too frequently? Do you want the second question?

DR McNICOL: I think that is so dramatic and sweeping, Dr MacDonald, we had better take them one by one.

MISS SHEPHERD: Yes, I need to think about that one. In any unit where bronchoscopy is commonly done the procedure with skilled anaesthetic assistance and a skilled bronchoscopist, particularly if there is no endobronchial lesions and no biopsy required, the risk is pretty minimal. I think the risks of bronchoscopy start rising quite sharply when a biopsy is required to be taken. Without any question it behoves us not to subject the patients to an unnecessary investigation. This is something which we must all have in the forefront of our minds. It involves an anaesthetic, it involves muscle relaxants, it involves a period of apnoea, and it is certainly not without risk. But I think this is a local application of a very important general medical maxim, is it not?

DR McNICOL: Can we ask you for the second half of your question, Dr MacDonald.

DR MACDONALD: I would like to pursue the first question a little further.

MISS SHEPHERD: Yes, I perhaps have not quite got there.

DR MACDONALD: Many patients are subjected to bronchoscopy far too often. I am not talking about thoracic units, cases are very often put up to the thoracic surgeon and the thoracic surgeon often does not like to refuse.

MISS SHEPHERD: I think we should throw that out to the chest physicians, don't you?

DR STRADLING (Chest Clinic, Hammersmith Hospital): I don't often disagree with Dr MacDonald but I do here very heartily indeed. I would take the completely reverse view and say that I feel bronchoscopy is done far too infrequently and that many is the time when people sit back pompously and discuss whether they should do a bronchoscopy and when they have got to that stage they should have done it. You never know what you are going to find. Very often you find things which are quite unexpected, very helpful indeed, and I would pick you up if I may, Miss Shepherd. You said it implied a period of apnoea, which of course is absolutely incorrect. With modern methods of anaesthesia you can have respiration which is almost normal.

MISS SHEPHERD: Apnoea on the part of the patient.

DR STRADLING: No.

MISS SHEPHERD: On the part of the patient. The patient is not breathing on his own, you have to use positive pressure.

DR STRADLING: Oh well, it is not apnoea. You are keeping your patient's blood gases normal, or better than normal, but this surely reduces the risk very considerably. I am afraid I am very much in favour of doing this as a matter of routine.

DR McNICOL: Dr MacDonald, your second question.

DR MACDONALD: My second question is what about the place of liver scan? Do you think it is at all useful?

MISS SHEPHERD: Yes, I do think it is useful. I did not touch on this side of things but obviously it is in my mind when I come to clinical examination, 'is there any suspicion at all that there could be liver involvement?' I think liver scan does have a useful part to play. One could argue on this point in the same way as mediastinoscopy; should every patient being presented for surgery have a liver scan? I don't know whether we would be too popular on this one but a case could be made out for it, I think.

DR K. H. LIM (Barnet Chest Hospital): When you mentioned liver scans and mediastinoscopy, would it be worth-while, before removing the lung, to do a laparotomy to look at the liver because we know the imperfections of the liver scan.

MISS SHEPHERD: I think that probably this would not be justifiable because you could only assess the situation on the surface of the liver, and there is an awful lot of liver tissue which you cannot see with a laparotomy. I would think that despite the drawbacks of liver scans this would be more valuable than opening a patient's abdomen.

DR MCHARDY-YOUNG (Central Middlesex Hospital): Can I just ask, if somebody is asking about scanning procedures, why choose to scan the liver and not do bone scans? If we are going to do a tremendous amount of pre-operative assessment I would have thought that bone scans were most important.

MISS SHEPHERD: I did not mention bone scans, but, yes, I do bone scans because bone secondaries from lung carcinoma are very often spinal and they are notoriously difficult to spot on straight radiological grounds. And on several occasions in these instances I have had a bone scan done and as categorical a report as can be made indicating the presence of bone secondaries.

DR McNICOL: Miss Shepherd finished a little early and she has already been answering questions for about 20 minutes. I must say that I am getting a little alarmed that her operability rate at the end of this discussion is going to drop to 10% or less by the time the patients have had their laparotomy and other investigations. However, I think if there are no burning questions we should perhaps leave it there. Thank you very much indeed, Miss Shepherd.