Discussion

Chairman: Dr M.W. McNICOL
Central Middlesex Hospital and Willesden Chest Clinic

DR MIKHAIL: I notice when you were giving your figures for radiotherapy you mentioned that the squamous carcinoma had a survival rate of 10%. Is there any figure at all to show what the survival rate would be if you did not treat the patients at all?

DR DEELEY: It would be less than 5% for 5 years survival rate. Somebody might have a different idea.

DR MIKHAIL: The 3 years survival rate you said would be less than 10% with treatment. What would it be if you did not treat?

DR DEELEY: It would be less than 5%.

DR J. H. ANGEL (Watford/Harefield Hospitals): There was a vogue at one time, I don't know if it still going on, for using combined pre-operative radiotherapy followed by surgery, first using conventional 28 day course and then having a rest, then having the operation, and later on having rather shorter courses, and I think a number of clinical trials have been done over the past 10 years. What is the present position on this? Is it out?

DR DEELEY: It is out at the present moment; I do not think it should be. The idea was to irradiate the tumour and hope that you would change the biological pattern in the cells, making it difficult for them to grow or mobilize or metastasize. I think this is a very desirable method. The trials which were reported previously had a radical course of radiotherapy, quite a high dose, followed by radical surgery and there were a high number of blown stump end complications. The doses were too high and this has got to be repeated, probably for a shorter time but certainly with much lower doses.

DR L. LANGE (Central Middlesex Hospital): I wondered in these 25% of patients with solitary cerebral metastases whether the contra-indication to surgery was the presence of a cerebral metastasis or whether surgery was rejected on the basis of the extent of the growth in the lung, and secondly, whether you feel it is a worth-while procedure to remove a solitary cerebral metastasis and then operate on the lung if it is deemed operable.

DR DEELEY: The answer to the first is that all of the series I have reported were patients with cerebral metastases and very inoperable because of their cerebral metastases. This took priority over the condition of the chest. The answer to the second is that of those patients who had cerebral metastases, half of them had multiple metastases in the brain and the other half had a single solitary metastasis. At the present moment there is no way of identifying single from multiple deposits. Radio-active brain scans will pick up single deposits but it will pick up the largest deposit and there may be smaller satellites around. I think at the present moment the surgeons are not very happy because they are not sure whether it is going to be one deposit or multiple deposits. There are reports in the literature of surgical ablation of deposits but it is rather disappointing because a larger number were found to have multiple deposits. We treat with whole brain irradiation because of this.

DR LANGE: I just wondered whether it would not be worth-while if there is no evidence of multiple metastases, in other words, if all the clinical evidence pointed to just one lesion, would it not be worth-while to try and remove this?

DR DELLEY: Yes, but I do not think we have sufficient means of identifying a single solitary metastasis in the brain. But it has been done by the surgeons and there have been some; I have forgotten the actual name of the report but it was done in Oxford and their results were rather disappointing.

DR LANGE: I don't wish to nag about this but I think we have methods for determining whether there is a metastasis. We can't be certain that there are not other metastases as well.

DR DEELEY: We have methods of saying whether there is a cerebral metastasis but it doesn't help us to say whether it is the only metastasis or whether the brain is riddled with smaller metastases.

DR K. M. CITRON (Brompton Hospital): I would like Dr Deeley to comment on a controlled trial which I am sure you have read. The one from Oxford by Riddehalge in which they designed a study to answer the question: 'What would you do with a patient with unresectable bronchial carcinoma who has no symptoms at the time of diagnosis directly consequent upon that carcinoma, for instance no pain or SVC obstruction? They randomized these into four groups: the first group had no radiotherapy, the second group had radical radiotherapy, the third group had nitrogen mustard and the fourth group had both cytotoxic drugs and radiotherapy. In this study the survival rate was uninfluenced by the treatment, the mean survival rate was only 8 months. The second point, and I think this is important, the quality of life did not appear to be improved at all by giving them either chemotherapy or radiotherapy. I think another important point was that of those who had no radiotherapy in the first place, those that did develop symptoms had just as good palliative relief by subsequent radiotherapy as those who had not received that treatment before. Their conclusion was that there is no indication for treating patients who have a confirmed unresectable bronchial carcinoma in the absence of symptoms. Would you agree with those conclusions? If not, why not?

DR DEELEY: I do know this trial very well. Perhaps I can sum up and you will get my impression if I say the longest survivor of carcinoma of the bronchus treated by radiotherapy was 32 months, and their radiotherapy does not compare with the figures I have just given you.

DR K. M. CITRON: They were given 6000 Rads.

DR DEELEY: I think this is too high. This is what I am trying to say. There comes a stage where you can improve by going up, but after about 4000 Rads you are getting too much effect on the normal tissues, and that

is the optimum level for radiation. I think 6000 Rads is too high. Because you must know also that the air in the lung does not stop the radiation and that when your rays are passing through normal tissues—I calculate my dosage on normal tissues assuming normal tissues absorption—but going through the alveoli filled with air there is not that absorption, and the dose in actual fact received by the tumour may be anything up to 30% higher than that estimated. So that my 4000 Rads represents about 5400 Rads, their 6000 represents a third of that again, may be up to 8000 Rads.

DR McNICOL: Could we ask you to speculate on what might have happened if the doses had been, in your terms, rather more appropriate.

DR DEELEY: I don't know, I think that is the next thing to do, but Professor Bleeheh is going to disagree with me, I think.

PROFESSOR BLEEHEH (Middlesex Hospital): I perhaps have an unfair advantage over the two previous speakers in that I do have a copy of the Oxford paper here. To correct a point, they gave a dose of 4000 Rads in 28 days.

DR DEELEY: Minimum dose.

PROFESSOR BLEEHEH: Well, yes, but . . .

DR DEELEY: It says definitely a minimum dose.

PROFESSOR BLEEHEH: Yes, to a planned tumour dose of 4000 Rads. Well, a minimum of 4000 Rads, in one and in the radiotherapy and chemotherapy to a planned tumour dose. Well, accepting that, I think on Dr Deeley's previous study he would claim this was more than the optimal dose. But to answer the other half of your question about the chemotherapy, which perhaps I can anticipate some of my talk, I think the point of interest in their study was that they showed within the limits of their experimental trial that the late treatment was no worse than immediate treatment, and I think this is an interesting point and one that perhaps should be taken into account in the design of further studies.

DR DEELEY: I am sorry to labour this point, but I don't think they can make that conclusion when their results of radiotherapy are not the best. Sorry, perhaps I am being unkind. I don't say mine are the best, but similar results have been reported from other people.

DR G. POOLE (Hammersmith Hospital): You said that you prefer to treat your patients at home, away from hospital infections and that seems very reasonable. However, perhaps you will have read the recent article about the survival of patients, those who have had post-operative empyema and those who had not, and there was a very significant difference. Those who had empyema survived much better than those who had not, and the conclusion was drawn, rightly or wrongly, that this may have stimulated the resistance in such a way that the disease was attacked more vigorously by the host. I wonder if your figures, if one looked at them, may be capable of telling you whether those patients who had infections during the course of their radiation or other treatment may have done better than those who did not, because this might raise another line along which we might pursue therapy in future.

DR DEELEY: I think this is a slightly different thing. These cases with empyema were not empyema with carcinoma were they? These are pure empyema, and remember our patients are in very poor clinical condition.

They are almost on their last legs, and I think the infection on top of that just tips the balance.

DR GRIFFEL: I think you mentioned that fibrosis is really the thing which kills cancer following your dosage. I think that empyema does the same. The best cases in treating with pneumothorax were cases with fibrosis developing. This is pre-drug treatment where the fibrosis and consequent fibrosis really obliterated the lymphatic spread. This may well happen in radiotherapy, when attempts to approach your dosage were done by Jollis of Northampton, who very early realized that the dosage is important: I live near Oxford, and I always disagreed with this high dosage. I think it does produce general symptoms and it is an attempt really to burn it out by impossible means. This applies to chemotherapy as well. Chemotherapy used as a general remedy reaches levels by which you destroy your patient, and I tried to avoid it by preserving the marrow, by squeezing the aorta for the time of the life of the chemotherapeutic agent, by cutting off the supply to the marrow in the long extremities and in the pelvis where there is a high proportion of marrow. Unfortunately, really very few patients survived this technique and I had to give it up, and I cannot forgive myself for doing it as long as I did. But it was an avenue for treating head tumours and tumours of the leg, by injecting directly, and by radiotherapy as a directly localized agent. That is why careful dosage is still the best way for non-operative cases. You mentioned this trial of treating by radiotherapy followed by surgery. I was one of the physicians where this was done. The interesting part was that in some of these cases when they came to the operation there was no evidence of previously histological confirmed cancer of the lung. So there was something in it, but it was given up because of too high high a dosage of radiation.

DR DEELEY: This is my whole policy; to give lower doses.

DR LIM: It seems to me that I have not been hearing about radiation pneumonitis so frequently nowadays. Is this the case or is radiation pneumonitis due to secondary infection which is nowadays treated promptly with antibiotics?

DR DEELEY: I think this is the case. With 250 kv you needed a number of fields—have we got a blackboard? If you don't mind me just drawing it. Let us take an outline of the chest and put in a tumour dose. With 250 kv the maximum dose received on the skin and the tumour received about 20%. Obviously, this was not feasible with one X-ray field so you put another field on which gave you 40, and another field on which gave you 60, 80, 100. Now we are getting an economic position. Now another field gives us 120 and you needed 76 fields around the chest. With super-voltage X-ray therapy your maximum dosage received between 1–2 cm below the skin and your skin dosage is about 10–15%, your tumour dose is about 70% and by putting on two of those fields you get 140%. We don't put two of those fields on because of the spinal cord, we take one field off. You get 140% but look at the difference in the amount of lung which is irradiated. And you are using higher dosage as well.

PROFESSOR B. J. O'LOUGHLIN (California, U.S.A.): I have been interested in smoking and cancer of the lung

for many years. I think Miss Shepherd left us with a properly pessimistic view of the curability of cancer of the lung, and I think that subsequently we have obscured the picture with our use of numbers. It seems to me that when we say we have a percentage of this or that we should say percentage of what? I think the question which was raised as to whether or not treatment was at all efficacious was a very pertinent one. If you see Miss Shepherd's figures and took 100 cases of cancer of the lung and said that 20% of them were operable, and of these 25% were resectable, and of these 15% were curable for 3 or 5 years, you would get one patient, and since her figures, and I think most surgeon's figures, for operable diseases are approximately the same, it is obvious that the patients might just as well be left alone, and I think probably the same thing is generally true for radiotherapy. I suppose these sorts of analyses are

subject to severe criticism from therapists. I would like to say there are two areas obviously unreferred to in these last analyses, and that is the case of prevention and the case of early diagnosis. In these areas there is hope for improvement. This conference should be more concerned with early diagnosis and prevention.

DR DEELEY: I do hope I had made that clear at the beginning. Three ways of managing cancer of the lung are (a) to prevent it, (b) diagnose it at an early stage where the condition is still confined to the chest and (c) finding methods of treatment. I think the results are very disappointing. I do not think the answer is going to be by improving surgery or improving radiotherapy. I think we have got to go back much further, pick up these patients at an early stage before they metastasize, or better still convince the public that smoking is dangerous.