TABLE 4. Mode of death in lung cancer

terminal care (Smithers, 1973), and ‘chat’ in this context means *patient* chat while the doctor listens. Most patients want to talk about their illness, to express their fears and anxiety, to seek reassurance that ‘it will not be too long’, that they will not suffocate or choke to death. For example, Mr C.J.B. (Case History 1) when admitted said he would like to put his head into a gas chamber. This was understandable in view of his condition; alone and virtually bedfast for 6 weeks. A week later, though washed and shaved and symptomatically improved he still wanted to kill himself. After a further week I asked

him again if he still wished to gas himself. He replied, ‘I don’t want to go home’. It then came out that he was living in a flat at the top of his son’s house and the relationship between them was more than strained. On no account did he wish to go back there. He was relieved when I assured him that we would not expect him to go back. For the rest of his admission, he was fully active and enjoyed life in the hospice. He definitely no longer wished to die. On this occasion we failed to patch up the family quarrel. Sometimes, however, we are able to help by bringing the two sides together.

Terminal illness should not be regarded as an interlude or intrusion into life. It is part of life and can be a most positive experience—a time of increasing maturity and deepening relationships. It is our job as doctors to help it be so.

Acknowledgments

The author wishes to thank Dr Joan Haram for encouraging him to study patients with lung cancer apart from those with other forms of malignant disease, and also Miss Susan Cayre for preparing Table 1. The author is in receipt of a research fellowship from the Sir Halley Stewart Trust.

References

SMITHERS, D. (1973) Care of Dying: Where to die. *British Medical Journal*, 1, 34.
WATSON, L. (1966) Calcium Metabolism and Cancer. *Australasian Annals of Medicine*, 15, 359.

Discussion

Chairman: DR J. R. MIKHAIL
(Central Middlesex Hospital)

DR SCADDING: In your patient who recurred after 11 years, was it the same histology or was it a new primary?

DR TWYXCROSS (St Christopher’s Hospice): I meant to say it could well have been a second primary. It is interesting to note that in those 2 years I was only able to put up 185 patients to take those symptoms from. There were eight other patients but they had to be excluded because they all had associated other primary carcinomas and I think that with a common disease such as lung cancer the possibility of a second lung cancer cannot be excluded.

DR K. P. BALL (Central Middlesex Hospital): I was very interested to hear you comment about the importance of chat. I remember Dr Arthur Wilox, a physician that many of us will remember from the Middlesex, used to underline the importance of not hurrying past a dying patient’s bed. It is very easy to do it on a busy round when one has other people to see for whom one could think you can do more. Another problem, of course, is that there are not enough St Christopher’s Hospices about. Many of the kinds of things you have been recommending there are difficult to do in a general

hospital. Obviously we ought to apply many more things in our own wards but we cannot do all of them.

DR TWYXCROSS: I accept there are problems, and I hope I made it clear that I fully understood it, having worked in general hospitals for some time before I took up this research fellowship. I do not know what the answer is except to follow the advice of the physician you referred to, and that of Sir David Smithers. I think it must be difficult for someone who is under the care of a radio-therapist or a chest physician who has been working trying to cure them and then suddenly at some time you have got to face up to the reality that you have failed. Gloom and everything goes rapidly across, not only the medical team but the relatives and the patient. I don’t know whether it is possible to get round this. We start afresh because we take them on and we have a fresh approach. We don’t have a legacy of failure and we don’t look at these patients as having incurable cancer. We tend to think of them as having diseases of pain, weakness, anorexia, etc. We set out to treat those, and when you have clerked a patient we say ‘Well, look. Your problem seems to me (1) you have got pain, (2) you

are off your appetite and (3) you have got this terrible weakness. We can help all these. Now we are going to work hard on the pain first and then we will come back and think about the second one. I am not promising you anything, but we are going to do our best'. So we can forget about the cancer. If they want to talk about it that is fair enough, but we are concentrating on what is their real problem as far as they see it, their symptoms. How you can incorporate this into a setting where you have been fighting to obtain curative results, I don't know.

DR BALL: I think one can because one has to put a positive science or art of symptomatic treatment. We often think of symptomatic treatment as being almost a failure, but I think we should consider it much more carefully.

DR TWYLCROSS: There should never come a time when we say 'there is nothing I can do for you'. I think this has been the mistake some of us have made in the past.

DR CITRON: I would very much like to support what Dr Ball has said, and I am sure that a lot of this should be done in the general hospitals. Dying is often a very lonely process because the patient is embarrassed to talk to his relatives about things he really wants to talk about. He is often likewise embarrassed to talk to the medical staff about his fears. It is very important to stress to all the staff that they should spend time with the patient to let the patient talk and not give the patient the impression that they are too busy. The second point is that I think that electrolyte analysis is a very useful thing in patients with terminal bronchial carcinoma. I think it was Roche that showed rather more than 15% of these patients have an abnormality such as a raised calcium or a low potassium, sometimes a low sodium. These things can be corrected and as a result the patient will feel very much better and stronger. It is well worth looking for this and correcting it where it is found.

DR TWYLCROSS: I am in entire agreement with you, but I work in a situation where biochemistry has officially been frowned on for a number of years, and it takes time to reverse this situation. I do biochemistry on a number of patients but certainly not on all the carcinoma of bronchus and not all the carcinoma of breast which I am sure we should. If you just do routine electrolytes, including calcium, on your auto-analyser you will miss a certain percentage of your hypocalcaemics because it is hyper-ionized calcium which gives you the symptoms. If you are somewhat wasted or you have been ill for some time, your albumin may well have dropped and if your albumin has dropped you will drop your bound calcium and this may keep your total calcium within the normal range and so you look at your figure and say 'Oh, well, he has only got a calcium of 9.8 mg/l; he can't be hypocalcaemic', but in actual fact he might. So you have either got to get the laboratory to do a specific gravity test and correct as they do at UCH or you have got to do ionized calcium levels themselves. You have got to go into it with your eyes open.

DR LIM (Barnet): I heartily agree about the use of corticosteroids in the relief of many conditions such as secondaries in the brain, weakness and hypocalcemia. My only criticism is very often 4 mg t.d.s. is not enough and very often we have to push the dose up to 20/30 mg of even 40 mg t.d.s.

DR TWYLCROSS: For what symptoms? All those, or any in particular?

DR LIM: Quite a lot of these, brain secondaries, muscular weakness and hypocalcemia. We have had to give Prednisolone, I think in one case we went up to 60 mg t.d.s.

DR TWYLCROSS: I agree entirely in the case of cerebral secondaries. This is perhaps why it is traditional to use Dexamethasone which is seven times as potent as Prednisone. If you are using Dexamethasone start with 4 mg q.d.s. if it has not been tried before and tailed off. We need to go into the exact indications for starting this in terminal care. But 4 mg q.d.s. is about 25-30 mg of Prednisone q.d.s. You really are pushing it in. Correct me if I have got my conversion figure wrong. I am sure you have got to consider this dosage. On the other hand the cerebral secondaries patients have an individual dose, and some of them don't need those high doses. When you have got them where they ought to be it is worth trying to drop until you get a re-appearance of one symptom, be it pain or any other symptom. Drop it until you get that and then just bump it up by one step. But I would agree that there are times when you need big doses, but it is not all the time.

DR MACDONALD: I take it all your patients know what is wrong with them by the time they come to you?

DR TWYLCROSS: It is difficult to put a percentage on that. I would have thought that a majority do not when they come to us, but by the time they die the majority do. You are always going to have a hard core of patients who go to the grave denying that there is anything wrong with them. You will always get a group who will always deny. It is rather like people who have to have cigarettes. They cannot face this and they have to keep up this barrage of denial. The majority of the rest, about 75%, or maybe 90%, do adjust at varying stages to the fact that they are dying. They may not realize they have cancer, but they are not interested in the diagnosis, they are interested in prognosis.

DR MIKHAIL: I think I must now draw this meeting to a close. Can I thank you all very much for your support in coming to-day with all the difficulties of the rail strike and I hope you feel the journey has been worth-while. I would particularly like to thank Dr Ball and ASH for their help in arranging the morning programme and to Professor Astrup in particular for undertaking the long journey to be with us to-day. I think this meeting would like to show their appreciation to Lederle Laboratories for sponsoring this symposium. Without their help we should not be here to-day. I am particularly indebted to them.