

- Tracey, J.K., Fong, Y., Hesse, D.G. *et al.* Anti-cachectin/ TNF monoclonal antibodies prevent septic shock during lethal bacteraemia. *Nature* 1987, **330**: 662–664.
- Welsh, C.H., Lien, D., Worthen, G.S. & Weil, J.V. Pentoxifylline decreased endotoxin-induced pulmonary neutrophil sequestration and extravascular protein accumulation in the dog. *Am Rev Respir Dis* 1988, **138**: 1106–1114.
- Schonharting, M.M. & Schade, U.F. The effect of pentoxifylline in septic shock—new pharmacologic aspects of an established drug. *J Med* 1989, **20**: 97–105.
- Streiter, R.M., Remick, D.G., Ward, P.A. *et al.* Cellular and molecular regulation of tumour necrosis factor- α production by pentoxifylline. *Biochem Biophys Res Commun* 1988, **155**: 1230–1236.
- Sullivan, G.W., Carper, H.T., Norick, W.J. & Mandell, G.L. Inhibition of the inflammatory action of interleukin and tumour necrosis factor (α) on neutrophil function by pentoxifylline. *Infect Immun* 1988, **56**: 1722–1729.
- Tighe, D., Moss, R., Hynd, J. *et al.* Pentoxifylline reverses hemodynamic and histological changes associated with peritonitis in pigs. In: Mandell, G.L. & Novick, W.J. Jr (eds) *Pentoxifylline and Leukocyte Function*. HRP Inc., Somerville, NJ, 1988, pp. 184–189.
- Ward, A. & Clissold, S.P. Pentoxifylline a review of its pharmacodynamic and pharmacokinetic properties, and its therapeutic efficacy. *Drugs* 1987, **34**: 50–97.
- Bjorvatn, B., Bjertnaes, L., Fadnes, H.O. *et al.* Meningococcal septicemia treated with combined plasmapheresis and leucapheresis or with blood exchange. *Br Med J* 1984, **288**: 439–441.

asymptomatic arrhythmias can be induced by hypoglycaemia and may suggest a possible mechanism for the phenomenon of sudden death in insulin dependent diabetics.

M.A. Baxter
C. Garewal
R. Jordan
A.D. Wright
M. Nattrass

Department of Medicine,
General Hospital,
Birmingham B15 2TH, UK.

References

- Collier, A., Matthew, D.M., Young, R.J. & Clarke, B.F. Transient atrial fibrillation precipitated by hypoglycaemia: Two case reports. *Postgrad Med J* 1987, **63**: 895–897.
- Gale, E.A.M. Hypoglycemia and human insulin. *Lancet* 1989, **2**: 1264–1266.
- Gale, E.A.M. The frequency of hypoglycemia in insulin-treated diabetic patients. In: Serrano-Rios, M., Lefebvre, P.J. (eds) *Diabetes 1985*. Elsevier, Amsterdam, 1986, pp. 943–947.
- DCCT Research Group Diabetes Control and Complications Trials (DCCT), results of feasibility study. *Diabetes Care* 1987, **10**: 1–19.

Hypoglycaemia and atrial fibrillation

Sir,
We would like to report a further case of hypoglycaemia-induced self remitting atrial fibrillation. Although 2 similar cases have been previously reported in the *Journal*,¹ we feel that the recent reports of sudden death in young insulin dependent diabetics² may make it important to re-emphasize this interesting observation.

A 27 year old male with Type I diabetes of 15 years duration presented to the Accident and Emergency Department following a severe hypoglycaemic reaction. He was taking three daily injections of a Human Velosulin (6 units) and Insulatard (12 units) and claimed good glycaemic control which was supported by a serum fructosamine of 2.95 mmol/l (normal range 1.9–2.7) on admission. He had however been admitted to hospital on two previous occasions following severe nocturnal hypoglycaemia. There was no past medical or family history of cardiac arrhythmias.

At 20.00 h on the day of admission he became weak and confused. His mother recognized his symptoms and recorded a blood glucose of 1.4 mmol/l (BM 1–44 stick read by meter) before giving the patient milk. In the Accident and Emergency Department (21.00 h) he was given a further dose of oral glucose and a subsequent Reflocheck of 14 mmol/l was recorded. Although asymptomatic he was in atrial fibrillation with ventricular rate 100–150 beats/min, confirmed on electrocardiograph. Physical examination was normal and within one hour the rhythm spontaneously reverted to sinus. Subsequent cardiological assessment (auscultation, resting electrocardiogram and echocardiogram), serum potassium and thyroid stimulating hormone levels were all normal.

Hypoglycaemia is common in insulin treated diabetics^{3,4} but the incidence of associated cardiac arrhythmias remains unknown. This case supports the view that

Inappropriate secretion of vasopressin, hypopituitarism and corticosteroid therapy

Sir,
A recent report describes five patients with hypopituitarism and severe hyponatraemia due to the syndrome of inappropriate antidiuretic hormone secretion (SIADH) that improved within a few days after the institution of hydrocortisone therapy, whereas the infusion of normotonic or hypertonic saline had been found to be less effective.¹

We have observed the association between hyponatraemia due to SIADH and hypopituitarism in a 60 year old previously healthy woman who presented with lethargy, weakness and complete absence of axillary and pubic hair. Laboratory features revealed severe hyponatraemia (111 mmol/l), low plasma/urine osmolality ratio (230/298 mosmol/kg) and an increased urinary sodium excretion (50 mmol/l). Chest X-ray, serum creatinine and arterial blood gases were normal. She was treated by restricting fluid intake and with intravenous administration of 5% saline solution that restored serum sodium concentration and plasma osmolality within the first week of admission. A clinical diagnosis of panhypopituitarism was confirmed by means of the LHRH-TRH test as well as both acute and continuous ACTH stimulation performed after a stay of 7 days, at which time hormonal replacement was instituted.

Oelkers emphasize that a response to corticosteroids should be the hallmark of SIADH secondary to hypopituitarism. Although hydrocortisone can restore sodium excretion to normal by inhibiting vasopressin directly³ there is also evidence that it has mineralocorticoid-like properties and produces renal tubular sodium reabsorption in the absence of ADH.⁴ On the other hand, we re-established normal serum sodium levels after conventional therapy without corticosteroids. Probably plasma osmolality improvement itself after saline infusion may