- 5 UK Prospective Study of Therapies of Maturity-onset Diabetes: 1 Effect of diet, sulphonylurea, insulin or biguanide therapy on fasting plasma glucose and body weight over one year. Multi-centre study. *Diabetologia* 1983; 24: 40–11.
- 6 Sterne J, Junien JL. Metformin: pharmacological mechanisms of the anti-diabetic and antilipidic effects and clinical consequences. *R Soc Med Int Congr Symp Ser* 1981; 48: 3–16.
- 7 Hermann LS. Metformin a review of its pharmacological properties and therapeutic use. *Diabetes Metab Rev* 1979; 5: 233–45.
- 8 UK Prospective Diabetes Study 11. Reduction in HbA1c with basal insulin supplement, sulphonylurea or biguanide therapy in maturity onset diabetes. *Diabetes* 1985; 24: 793–8.
- 9 Wales JK. Treatment of the obese diabetic patient. In: Bjorntorp P, Cairelli M, Howard AN, eds. *Recent advances in obesity research*. London: Libby, 1980; pp 184–9.
- 10 Wu MS, Johnston P, Sheu WHH, et al. Effect of metformin on carbohydrate and lipoprotein metabolism in NIDDM patients. *Diabetes Care* 1990; 13: 1–8.
- 11 Chakrabarti R, Hocking ED, Fearnley GR. Fibrinolytic effect of metformin in coronary artery disease. *Lancet* 1965; 2: 256–9.
- 12 DeSilva SR, Shawe JE, Patel H, Cudworth AG. Plasma fibrinogen in diabetes mellitus. *Diabetes Metab* 1979; 5: 201–6.
- 13 Chan JC, Tomlinson B, Crichtley JA, Cockram CS, Walden RJ. Metabolic and haemodynamic effects of metformin and glibenclamide in normotensive NIDDM patients. *Diabetes Care* 1993; 16: 1035–8.
- 14 Vague P, Juhan-Vague I, Alessi MC, Badier C, Valadier J. Metformin decreases the high plasminogen activator inhibition capacity, plasma insulin and triglyceride levels in non-diabetic obese subjects. *Thromb Haemost* 1987; 57: 326–8.
- 15 Robertson DA, Hale PJ, Nattrass M. Macrovascular disease and hyperinsulinaemia. *Clin Endocrinol Metab* 1988; 2: 407–24.
- 16 Bailey CJ. Biguanides and NIDDM. *Diabetes Care* 1992; 15: 755–70.
- 17 Noll F. *Methods of enzymatic analysis*. 2nd edn. HU Bergmeyer, p1475.
- 18 Prager R, Schernthaser G, Graf H. Effect of metformin on peripheral insulin sensitivity in non-insulin dependent diabetes mellitus. *Diabetes Metab* 1986; 12: 346–50.
- 19 Gawler DJ, Milligan G, Houslay MD. Treatment of streptozocin rats with metformin restores the ability of insulin to inhibit adenylate cyclase activity. *Biochem J* 1988; 249: 537–42.
- 20 Nattrass M, Todd PG, Hinks L, Lloyd B, Alberti KGMM. Comparative effects of phenformin, metformin, and glibenclamide on metabolic rhythms in maturity onset diabetes. *Diabetologia* 1977; 13: 145–52.
- 21 Noel M. Kinetic study of normal and sustained release dosage forms of metformin in normal subjects. *Res Clin Forums* 1979; 1: 35–50.
- 22 Pentikainen PJ, Neuvonen PJ, Penttila A. Pharmacokinetics of metformin after intravenous and oral administration to man. *Eur J Clin Pharmacol* 1979; 16: 195–202.
- 23 Tucker GT, Casey C, Phillips PJ, Connor H, Ward JD, Woods HF. Metformin kinetics in healthy subjects and in patients with diabetes mellitus. *Br J Clin Pharmacol* 1981; 12: 235–46.
- 24 Wiholm BE, Myrhed M. Metformin associated lactic acidosis in Sweden 1977–1991. *Eur J Clin Pharmacol* 1993; 44: 589–91.
- 25 Langan SJ, Deary IJ, Hepburn DA, Frier BM. Cumulative cognitive impairment following recurrent severe hypoglycaemia in adult patients with insulin-treated diabetes mellitus. *Diabetologia* 1991; 34: 337–44.
- 26 Ferner RE, Neil HAW. Sulphonylureas and hypoglycaemia. *BMJ* 1988; 296: 949–50.

Medical Anniversary

JAMES YOUNG SIMPSON, 7 JUNE 1811

(Sir) James Young Simpson (1811–70) was born in Bathgate, Scotland, the seventh son of a baker. He was educated in Edinburgh, where he qualified in medicine (1830), became an MD (1832), and was appointed professor of obstetrics (1839). He introduced chloroform for the management of labour (1847), and this was used by Queen Victoria (1853).

— DG James