Management of shoulder pain in patients with stroke

K Walsh

Abstract

Shoulder pain affects from 16% to 72% of patients after a cerebrovascular accident. Hemiplegic shoulder pain causes considerable distress and reduced activity and can markedly hinder rehabilitation. The aetiology of hemiplegic shoulder pain is probably multifactorial. The ideal management of hemiplegic stroke pain is prevention. For prophylaxis to be effective, it must begin immediately after the stroke. Awareness of potential injuries to the shoulder joint reduces the frequency of shoulder pain after stroke. The multidisciplinary team, patients, and carers should be provided with instructions on how to avoid injuries to the affected limb. Foam supports or shoulder strapping may be used to prevent shoulder pain. Overarm slings should be avoided. Treatment of shoulder pain after stroke should start with simple analgesics. If shoulder pain persists, treatment should include high intensity transcutaneous electrical nerve stimulation or functional electrical stimulation. Intra-articular steroid injections may be used in resistant cases.

(*Postgrad Med J* 2001;77:645–649)

Keywords: shoulder pain; stroke

Shoulder pain is a common complication after a cerebrovascular accident. From 16% to 72% of stroke patients develop hemiplegic shoulder pain.^{1–3} It may occur in up to 80% of stroke patients who have little or no voluntary movement of the affected upper limb.⁴

Hemiplegic shoulder pain has been shown to affect stroke outcome in a negative way.⁵ It interferes with recovery after a stroke: it can cause considerable distress and reduced activity and can markedly hinder rehabilitation.^{6–8} Roy *et al* demonstrated that the presence of hemiplegic shoulder pain is strongly associated with prolonged hospital stay and poor recovery of arm function in the first 12 weeks after stroke.⁹

The cause of hemiplegic shoulder pain is the subject of considerable controversy. The following processes have all been postulated as causes of a painful hemiplegic shoulder: glenohumeral subluxation, spasticity of shoulder muscles, impingement, soft tissue trauma, rotator cuff tears, glenohumeral capsulitis, bicipital tendinitis, and shoulder hand syndrome.^{1–2 10–14} Traction neuropathy of the brachial plexus may also play a part.⁶ Unusual patterns of motor recovery or spasticity or unusually severe focal atrophy may suggest

brachial plexus injury. Poor handling of a hemiplegic limb may exacerbate a pre-existing condition such as osteoarthritis. Thus, pre-morbid disease of the shoulder may predispose to hemiplegic shoulder pain.⁶ Stroke patients may suffer from pain that is caused by the stroke itself (central post-stroke pain).⁶ The role of central post-stroke pain in the aetiology of hemiplegic shoulder pain is uncertain. Abnormal tone (both spasticity and flaccidity) has been suggested as an aetiological factor in hemiplegic shoulder pain. However, clinical observations suggest that shoulder pain does not occur until spasticity develops. Most authorities agree that the aetiology of hemiplegic shoulder pain is probably multifactorial.⁶

Prevention

The ideal management of hemiplegic stroke pain is to prevent it happening in the first place. Various strategies have been employed in the prophylaxis of hemiplegic shoulder pain. For prophylaxis to be effective, it must be begun immediately after the stroke. Once the patient has pain, resultant anxiety and overprotection will follow.¹²

HANDLING

Poor handling and positioning of the affected upper limb in stroke patients contribute toward shoulder pain.^{7 11 12}

The mobility of the recovering stroke patient is dependent on the assistance of nurses, therapists, doctors, other ancillary staff, and family members. It is also dependent on his/her own efforts. Handling, positioning, and transferring on a day-to-day basis can exert great stress on the vulnerable shoulder. The problem may be exacerbated by the patient's sensory and perceptual deficits. There has been concern that trauma to the constituent components of the shoulder joint may be caused by poor handling of the patient's affected arm.¹⁵

Wanklyn *et al* studied the prevalence of hemiplegic shoulder pain and associated factors in patients with stroke.¹⁶ Sixty three per cent of the patients developed hemiplegic shoulder pain in the first six months after their stroke. Patients who needed help with transfers were more likely to develop hemiplegic shoulder pain. Certainly, patients with markedly decreased voluntary movement after a cerebrovascular accident frequently experience shoulder joint malalignment or subluxation in the early stages of recovery.¹²

Careful positioning and handling of the limb are thought to prevent hemiplegic shoulder pain, but there is a range of opinions about how correct limb positioning is best achieved.^{2 9 17}

Colchester General
Hospital, Colchester,
UK

Correspondence to:
Dr Kieran Walsh, Basildon
Hospital, Nethermayne,
Basildon, Essex SS16 5NL,
UK

Submitted 21 March 2001
Accepted 15 May 2001

Braus *et al* investigated the efficacy of an information and education programme in the prevention of hemiplegic shoulder pain.¹⁸ All members of the diagnostic and therapeutic team as well as patients and their family were provided with instructions on how to avoid injuries to the affected limb. The investigators found that awareness of potential injuries to the structures of the shoulder joint reduced the frequency of shoulder pain from 27% to 8%.

Fitzgerald-Finch *et al* advocated the use of the Australian lift when handling these patients: they felt it to be of value as the weight of the patient is taken on the shoulders of the carer and the patient's shoulder is protected.¹⁹

STRAPPING

Glenohumeral joint subluxation may be a contributing factor in the development of shoulder pain in this group of patients.²⁻²⁰ Shai *et al* hypothesised that earlier radiological diagnosis of subluxation might enable more effective prevention than if it is delayed.²¹ However, this has not been proved. Despite this, a variety of slings have been designed to try to correct subluxation and pain in stroke patients with hemiplegia. Not all such devices have been successful: supportive devices developed by Buccholtz Moodie *et al* and Williams *et al* were not proved to be effective in correcting subluxation of the shoulder.²²⁻²³

Physiotherapists have employed various forms of strapping designed for shoulder pain or subluxation after a cerebrovascular accident. Unfortunately, the effectiveness of many of these strapping methods remains largely unproved. Ancliffe undertook a pilot study to determine the effectiveness of a strapping technique to prevent shoulder pain after a stroke.²⁴ The pilot study demonstrated that strapping the hemiplegic shoulder delayed the onset of shoulder pain. In patients with subluxation and shoulder pain, use of a Varney brace has been reported to be successful: patients become asymptomatic within seven days.²⁵

External support can be discontinued when muscle tone around the glenohumeral joint is sufficient to prevent subluxation.²⁶ An exercise programme should always accompany the use of a sling.²⁷

However, a number of authors have reported that slings may hold the limb in a poor position that is likely to cause soft tissue contracture and have an adverse effect on symmetry, balance, and body image.²⁸⁻³⁰

PHYSIOTHERAPY

Some studies have noted that passive abduction of the hemiplegic arm can result in rotator cuff injury: this in turn causes shoulder pain.¹²⁻¹³⁻³¹

However, therapeutic range of motion exercises done by the patients can involve passive abduction of the arm. Kumar *et al* analysed the occurrence of pain in patients receiving three different rehabilitation exercise programs: range of motion by the therapist, use of a skateboard, and use of an overhead pulley.³²

They found that patients who used the overhead pulley had the highest risk of developing shoulder pain and concluded that use of the pulley should be avoided during stroke rehabilitation.

If impingement during range of motion exercises is determined to be the cause of hemiplegic shoulder pain, the amplitude of passive movement should be kept within the pain-free range. Caldwell *et al* reported that pain subsided in 43% of patients with hemiplegic shoulder pain when the amplitude of passive range of motion was reduced.³³

Wanklyn *et al* reported an increase in the prevalence of shoulder pain in the first weeks after discharge in patients who did not continue to exercise properly.¹⁶

Treatment

Radiological investigations should exclude dislocation or fracture of the shoulder before further management is instigated. Various treatments have been suggested as being beneficial in shoulder pain after stroke: these include physiotherapy, localised cooling, infrared, ultrasound, and intra-articular injections of steroids and local anaesthetics.³⁴⁻³⁵ Until recently there has been a shortage of prospective controlled clinical trials.

PHYSIOTHERAPY

Physiotherapy has been used in the treatment of hemiplegic shoulder pain.¹²⁻³⁶ There are two major approaches to therapy in this field: those that focus on the problem as a localised mechanical one; and those that view the problem as a neurological one. Local treatments used have included heat and cold therapy.³⁷⁻³⁹ Slings and shoulder supports have also been used.⁴⁰⁻⁴¹ Positioning is also considered important by many authors.⁷⁻⁴² Other physiotherapy approaches include those of Bobath, Brunnstrom, and proprioceptive neuromuscular facilitation. Until recently, the evidence for the effectiveness of these methods of physiotherapy has been poor.³⁷ Partridge examined the effectiveness of two methods of physiotherapy in the treatment of hemiplegic shoulder pain: cryotherapy or the Bobath approach.⁴³ The cryotherapy approach involved the application of ice to the affected shoulder. The Bobath approach is a neurologically based holistic approach that is frequently used in the UK.⁴⁴

There were no significant differences between the two treatments in terms of severity of pain at rest or on movement or for reported distress. However the proportion of patients who reported no pain after treatment was greater in those who received the Bobath approach.

DRUG TREATMENT

Analgesic, anti-inflammatory, and antispastic drugs have all been used to treat hemiplegic shoulder pain.²⁻¹²⁻³⁷ Simple analgesics and non-steroidal anti-inflammatory drugs should be tried first. Antispasmodic medication may be helpful in spasticity of cerebral origin. Antispasmodic agents may supplement inhibition

and relaxation techniques in physiotherapy.² These agents have a modest effect on post-stroke hypertonicity but their cognitive side effects may limit their usefulness.

Braus *et al* conducted a trial on 36 stroke patients with hemiplegic shoulder pain.¹⁸ In a placebo controlled, non-blinded trial, they found that 31 of the 36 patients became almost symptom-free within 10 days of treatment with low dose oral corticosteroids (methylprednisolone 32 mg daily). A two week course of treatment was given followed by a two week taper. There were no major adverse events due to the drugs. Davis *et al* reported complete resolution of symptoms in 68 patients with hemiplegic shoulder pain using oral steroids and an intensive rehabilitation programme.⁴⁵

Dekker *et al* investigated the effect of intra-articular triamcinolone acetonide on pain and passive range of motion in seven patients with a painful hemiplegic shoulder.⁴⁶ Four out of the seven patients had significant reduction in pain. There was improvement in range of motion but it did not reach statistical significance.

TRANSCUTANEOUS ELECTRICAL NERVE STIMULATION

Leandri *et al* evaluated the effectiveness of high intensity versus low intensity transcutaneous electrical nerve stimulation (TENS) versus placebo for patients with hemiplegic shoulder pain.⁴⁷ Low intensity TENS involves electrical stimulation just above the level of the skin sensory threshold. High intensity TENS is sufficient to elicit muscle contraction and an almost painful sensation. The investigators found that patients who received high intensity TENS had significant improvements in passive range of motion for flexion, extension, abduction, and external rotation at the shoulder. The patients who received high intensity TENS also reported very satisfactory pain relief.

FUNCTIONAL ELECTRICAL STIMULATION

There have been a number of studies of the effectiveness of functional electrical stimulation (FES) in shoulder pain after stroke.

Faghri *et al* studied the effects of a FES treatment programme designed to prevent glenohumeral joint stretching and subsequent subluxation and shoulder pain in stroke patients.⁴⁸ They demonstrated a beneficial effect on subluxation and improvement in other parameters such as pain, range of motion, and arm function.

Chantraine *et al* completed a long term controlled study of the use of FES in hemiplegic stroke patients diagnosed with a subluxed and painful shoulder.⁴⁹ They found that a 24 month FES programme was effective in reducing the severity of subluxation and pain and may have facilitated recovery of shoulder function in these patients.

As an extremely mobile joint, the shoulder sacrifices stability for mobility.¹² Basmajian determined through electromyographic studies that the supraspinatus, and to a lesser extent the posterior deltoid muscles, played a key part in maintaining glenohumeral alignment.⁵⁰

Learning points

- From 16% to 72% of stroke patients develop hemiplegic shoulder pain.
- Hemiplegic shoulder pain has been shown to affect stroke outcome in a negative way.
- Most authorities agree that the aetiology of hemiplegic shoulder pain is probably multifactorial.
- Poor handling and positioning of the affected upper limb in stroke patients contribute toward shoulder pain.
- High intensity TENS can relieve pain and improve range of motion in hemiplegic shoulder pain.
- Functional electrical stimulation has been shown to improve parameters such as pain, range of motion, and arm function in patients with hemiplegic shoulder pain.

Chaco and Wolf also demonstrated the importance of the supraspinatus muscle in preventing downward subluxation of the humerus.⁵¹ Two studies have investigated the application of electrical stimulation to the supraspinatus and posterior deltoid muscles: Baker and Parker demonstrated the beneficial effects of FES in stroke patients with a chronic shoulder dislocation.⁵¹ However, patients deteriorated after withdrawal from treatment though not back to pre-treatment levels. Linn *et al* carried out a prospective randomised study to determine the efficacy of FES in the prevention of shoulder subluxation in stroke patients.⁵² They found that FES does prevent shoulder subluxation, but this effect was not maintained after the withdrawal of treatment.

BOTULINUM TOXIN

Bhakta *et al* evaluated the impact of botulinum toxin on disability caused by upper limb spasticity after stroke.⁵³ They demonstrated improvement in shoulder pain in six of nine patients treated with complete resolution in two patients.

SURGERY

Various authorities in the treatment of hemiplegic shoulder pain have recommended surgery on the structures of the shoulder. There is a lack of trial evidence in this area. Surgery may be of use if conservative methods have failed and the shoulder has become very painful and stiff. Recent improvements in rehabilitation techniques have reduced the need for surgical intervention.⁵⁴

The following procedures have been used to treat hemiplegic shoulder pain: operative surgery on the contractures in muscle tendons, surgical repair of a rotator cuff tear, stellate ganglion block, and scapular mobilisation.^{33 55 56} Braun *et al* reported the use of surgical release of spastic internal rotator muscles in the treatment of hemiplegic shoulder pain; all 13 patients who underwent this operation

Questions (answers at end of paper)

1. What effect does shoulder pain have on the recovery of a patient with stroke?
2. What causes hemiplegic shoulder pain?
3. What individuals play an important part in the handling of stroke patients?
4. What outcomes should be used in the assessment of treatments in hemiplegic shoulder pain?
5. What are the indications for surgery in patients with hemiplegic shoulder pain?

became pain-free in contrast to a control group whose pain remained.⁵⁷

Indications for surgery in various series have included range of motion limitation to the point of functional impairment, pain of such intensity that it interferes with skin hygiene or prevents participation in rehabilitation.³³⁻⁵⁵ Surgery is usually delayed until at least six months after the stroke to allow as much spontaneous functional improvement as possible.⁵⁵

Future

There are a number of obstacles to clinical research in this area. It is often very difficult to ascertain the patient's pre-morbid shoulder function or a history of trauma to the shoulder after the stroke. Most of the work done in this area has been in the form of case series; randomised double blind trials have thus far been sparse. However, further trials are needed to establish the optimal management of patients with hemiplegic shoulder pain.

- 1 Hanukah A, Sashimi H, Ohkawa T, *et al.* Arthrographic findings in hemiplegic shoulders. *Arch Phys Med Rehabil* 1984;65:706-11.
- 2 Van Ouwenaller C, Laplace P, Chantaine A. Painful shoulder in hemiplegia. *Arch Phys Med Rehabil* 1986;46:23-6.
- 3 Williams J. Electromyographic feedback and the painful hemiplegic shoulder. In: Michel T, ed. *Pain*. Edinburgh: Churchill Livingstone, 1985: 183-206.
- 4 Ancliffe J. Shoulder pain in hemiplegia: incidence and influence on movement and recovery of function. *Proceedings 3rd International Physiotherapy Congress*. Hong Kong, 1990: 187-92.
- 5 Roy CW, Sands MR, Hill LD, *et al.* The effect of shoulder pain on outcome of acute hemiplegia. *Clin Rehabil* 1995;9:21-7.
- 6 Griffin JW. Hemiplegic shoulder pain. *Phys Ther* 1986;12: 1884-93.
- 7 Anderson L. Shoulder pain in hemiplegia. *Am J Occup Ther* 1985;39:11-19.
- 8 Mulley GP. *Practical management of stroke*. London: Croom Helm, 1985.
- 9 Roy CW, Sands MR, Hill LD, *et al.* Shoulder pain in acutely admitted hemiplegics. *Clin Rehabil* 1994;8:334-40.
- 10 Van Langenberghe HVK, Hogan BM. Degree of pain and grade of subluxation in the painful hemiplegic shoulder. *Scand J Rehabil Med* 1986;20:161-6.
- 11 Jensen E. The hemiplegic shoulder. *Scand J Rehabil Med* 1980;7(suppl):113-19.
- 12 Cailliet R. *The shoulder in hemiplegia*. Philadelphia: FA Davis, 1980.
- 13 Najenson T, Yacubovic E, Pikenley S. Rotator cuff injuries in shoulder joints of hemiplegic patients. *Scand J Rehabil Med* 1971;3:131-7.
- 14 Rizk TE, Christofer RP, Pinals RS, *et al.* Arthrographic studies in painful hemiplegic shoulders. *Arch Phys Med Rehabil* 1984;65:254-6.
- 15 Cailliet R. *Shoulder pain*. Philadelphia: FA Davis, 1966.
- 16 Wanklyn P, Forster A, Young J. Hemiplegic shoulder pain (HSP): natural history and investigation of associated features. *Disabil Rehabil* 1996;18:497-501.
- 17 Jespersen HF, Jorgensen HS, Nakayama H, *et al.* Shoulder pain after stroke. *Int J Rehabil Res* 1995;18:273-6.
- 18 Braus DF, Krauss JK, Strobel J. The shoulder-hand syndrome after stroke: a prospective clinical trial. *Ann Neurol* 1994;36:728-733.
- 19 Fitzgerald-Finch OP, Gibson JM. Subluxation of the shoulder in hemiplegia. *Age Ageing* 1975;4:10-15.
- 20 Grossen-Sills J, Schenkman M. An analysis of shoulder pain, range of motion and subluxation in patients with hemiplegia. *Phys Ther* 1985;65:731.
- 21 Shai G, Ring H, Costoff H, *et al.* Glenohumeral joint malalignment in the hemiplegic shoulder: an early radiological sign. *Scand J Rehabil Med* 1984;16:133-6.
- 22 Buchholtz Moodie N, Brisbin J, Morgan A. Subluxation of the glenohumeral joint in hemiplegia: evaluation of supportive devices. *Physiotherapy Canada* 1986;38:151-7.
- 23 Williams R, Taffs L, Minuk T. Evaluation of two support methods for the subluxated shoulder of hemiplegic patients. *Phys Ther* 1988;68:1209-14.
- 24 Ancliffe J. Strapping the shoulder in patients following a cerebrovascular accident (CVA): a pilot study. *Australian Journal of Physiotherapy* 1992;38:37-41.
- 25 Krempen JF, Silver RA, Hadley J, *et al.* The use of the Varney brace for subluxating shoulders in stroke and upper motor neuron injuries. *Clin Orthop* 1977;122:204-6.
- 26 McCollough NC. The role of the orthopaedic surgeon in the treatment of stroke. *Orthop Clin North Am* 1978;9:305-24.
- 27 Wilson DJ, Caldwell CB. Central control insufficiency: III. Disturbed motor control and sensation: a treatment approach emphasising upper extremity orthoses. *Phys Ther* 1978;58:313-20.
- 28 Brook MM, De Lateur BJ, Diana-Rigby GC, *et al.* Shoulder subluxation in hemiplegia: effects of three different supports. *Arch Phys Med Rehabil* 1991;72:582-6.
- 29 Carr JH, Shepherd RB. *A motor learning programme for stroke*. 2nd Ed. London: Butterworth-Heinemann Medical Books, 1987: 106-7.
- 30 Davis PM. *Steps to follow*. New York: Springer-Verlag, 1985: 206-35.
- 31 Chaco J, Wolf E. Subluxation of the glenohumeral joint in hemiplegia. *Am J Phys Med* 1971;50:139-43.
- 32 Kumar R, Metter EJ, Mehta AJ. Shoulder pain in hemiplegia: the role of exercise. *Am J Phys Med Rehabil* 1990;69:205-8.
- 33 Caldwell CB, Wilson DJ, Braum RM. Evaluation of treatment of upper extremity in the hemiplegic stroke patient. *Clin Orthop* 1969;63:69-93.
- 34 Wood C. Shoulder pain in stroke patients. *Nursing Times* 1989;85(2):32-4.
- 35 Snels IA, Beckerman H, Lankhorst GJ, *et al.* Treatment of hemiplegic shoulder pain in the Netherlands: results of a national survey. *Clin Rehabil* 2000;14:20-7.
- 36 Griffin J, Reddin G. Shoulder pain in patients with hemiplegia. *Phys Ther* 1981;61:1041.
- 37 Roy CW. Shoulder pain in hemiplegia: a literature review. *Clin Rehabil* 1988;2:35-44.
- 38 Inaba MK, Piorkowski M. Ultrasound in treatment of painful shoulders with hemiplegia. *Phys Ther* 1980;52:737-41.
- 39 Lee JM, Warren MP. *Cold therapy in rehabilitation*. London: Bell & Hyman, 1978.
- 40 Brudny J. New orthosis for the treatment of hemiplegic shoulder subluxation. *Orthotics and Prosthetics* 1985;39:14-20.
- 41 Sodring KM. Upper extremity for stroke patients. *Int J Rehabil Res* 1980;3:33-8.
- 42 Ingenito R, Priestley L. *Hemiplegia: current approaches to patient positioning*. Washington, DC: Rehabilitation Research and Training Center, George Washington Medical Center, 1979.
- 43 Partridge CJ, Edwards SM, Mee R, *et al.* Hemiplegic shoulder pain: a study of two methods of physiotherapy treatment. *Clin Rehabil* 1990;4:43-9.
- 44 Bobath B. *Adult hemiplegia: evaluation and treatment*. London: William Heinemann, 1978.
- 45 Davis SW, Petrillo CR, Eichberg RD, *et al.* Shoulder-hand syndrome in a hemiplegic population: a five year retrospective study. *Arch Phys Med Rehabil* 1977;58:353-6.
- 46 Dekker JHM, Wagenaar RC, Lankhorst GJ, *et al.* The painful hemiplegic shoulder: effects of intra-articular triamcinolone acetonide. *Am J Phys Med Rehabil* 1997;76:43-8.
- 47 Leandri M, Parodi CI, Corrieri N, *et al.* Comparison of TENS treatments in hemiplegic shoulder pain. *Scand J Rehabil Med* 1990;22:69-72.
- 48 Faghri PD, Rodger MM, Glaser RM, *et al.* The effects of functional electrical stimulation on shoulder subluxation, arm function recovery and shoulder pain in hemiplegic stroke patients. *Arch Phys Med Rehabil* 1994;5:73-9.
- 49 Chantaine A, Baribeault A, Uebelhart D, *et al.* Shoulder pain and dysfunction in hemiplegia: effects of functional electrical stimulation. *Arch Phys Med Rehabil* 1999;80:328-31.
- 50 Basmajian JV. *Muscles alive: their functions revealed by electromyography*. Baltimore: Williams & Wilkins, 1978.
- 51 Baker LL, Parker K. Neuromuscular electrical stimulation of the muscles surrounding the shoulder. *Phys Ther* 1986;66:1930-7.
- 52 Linn SL, Granat MH, Lees KR. Prevention of shoulder dislocation after stroke with electrical stimulation. *Stroke* 1999; 30:963-8.
- 53 Bhakta BB, Cozens JA, Bamford JM, *et al.* Use of botulinum toxin in stroke patients with severe upper limb spasticity. *J Neurol Neurosurg Psychiatry* 1996;61:30-5.

- 54 Ryerson SD. Hemiplegia resulting from vascular insult or disease. In: Umphred DA, ed. *Neurological rehabilitation*. St Louis: CV Mosby, 1985: 474–514.
- 55 Jordan C, Walters RL. Stroke. In: VL Nickel, ed. *Orthopaedic rehabilitation*. New York: Churchill Livingstone, 1982: 277–91.
- 56 Black-Schaffer RM, Kirshtein AE, Harvey RL. Stroke rehabilitation. 2. Co-morbidities and complications. *Arch Phys Med Rehabil* 1999;80:S8–S16.
- 57 Braun RM, West F, Mooney V, et al. Surgical treatment of the painful shoulder contracture in the stroke patient. *J Bone Joint Surg Am* 1971;53:1307–12.

Answers

1. Hemiplegic shoulder pain affects stroke outcome in a negative way. It interferes with recovery after a stroke: it can cause considerable distress and reduced activity and can markedly hinder rehabilitation. Hemiplegic shoulder pain has been associated with prolonged hospital stay and poor recovery of arm function in the first 12 weeks after stroke.
2. The cause of hemiplegic shoulder pain is the subject of considerable controversy. The following processes have all been postulated as causes of a painful hemiplegic shoulder: glenohumeral subluxation, spasticity of shoulder muscles, impingement, soft tissue trauma, rotator cuff tears, glenohumeral capsulitis, bicipital tendinitis, and shoulder hand syndrome. The aetiology of hemiplegic shoulder pain is probably multifactorial.
3. The mobility of the recovering stroke patient is dependent on the assistance of nurses, therapists, doctors, other ancillary staff, and family members. It is also dependent on the patient's own efforts. Poor handling and positioning of the affected upper limb in stroke patients contribute toward shoulder pain.
4. Important outcomes include pain relief, improved passive and active range of motion and arm function.
5. Indications for surgery in patients with hemiplegic shoulder pain include limitation of range of motion to the point of functional impairment, pain of such intensity that it interferes with skin hygiene or prevents participation in rehabilitation.