

# PostScript

## LETTERS

### Cost effective strategy for a safe diagnosis of deep vein thrombosis at a district general hospital

We agree with the view of Walsh *et al* that it is not acceptable to miss a deep vein thrombosis (DVT) in any patient,<sup>1</sup> and the negative predictive value of any test (or combination of tests) must approach 100%. The authors mentioned impedance plethysmography, but did not elaborate the utility of such an investigation, yet importantly they discussed the cost issues at the end of their discussion section for a planned management of DVT at a district general hospital.

We propose that the impedance plethysmography together with pre-test probability scoring and a modern D-dimer test would be an ideal choice for a district general hospital, in order to reduce the cost of investigations for DVT, without jeopardising the safety of the patients.

Flanagan *et al* recently reported that computed strain gauge plethysmography can be used as a satisfactory first line investigation for the diagnosis of DVT, with a negative predictive value of 97%.<sup>2</sup> In a recent study a cheaper version known as digital photoplethysmography (D-PPG) has also shown a negative predictive value of 100% when used on its own.<sup>3</sup> We believe that these techniques, together with pre-test probability scoring and a modern D-dimer test, would approach a negative predictive value of 100%, fulfilling the authors' assumption of good practice.

We carried out a trial using combined D-PPG and D-dimer testing to investigate DVT in the outpatient department and compared the results against "gold standard" ultrasonography. In a six month trial (from September 2001 to February 2002) of 134 patients we were able to pick up 18 cases of positive DVT (all Doppler positive), 77 cases of negative DVT (all Doppler negative), and 39 cases with equivocal/unable results (all Doppler negative). This confirmed that 77 cases would not have required a Doppler study, 18 cases would definitely have required a Doppler study, and 39 cases would have required further examinations. A summary of the results are shown in the table 1.

All the recent studies<sup>1-4</sup> have major implications for practice because they show that the combination of a low pre-test probability score, derived from a formal scoring system, and a negative D-dimer test safely excluded DVT in outpatients, obviating further investigation in 40% of patients.<sup>4</sup> The very low incidence of DVT (0.6%) during follow up among those with a low pre-test probability score and negative D-dimer test is reassuring<sup>4</sup> when compared with an incidence of about 1% at three months among untreated patients with suspected DVT and a negative venogram.<sup>5</sup> However, the occurrence of DVT in up to 20% of patients with a high pre-test probability score and negative D-dimer test emphasises the point that the D-dimer test cannot be used in isolation.<sup>4</sup>

Therefore, it is possible that an objective of combining a formally derived pre-test prob-

Variable	No
Clinical	
Positive	18
Negative	116
D-dimer	
Positive	25
Negative	109
D-PPG	
Positive	18
Negative	77*
Equivocal	27*
Unable	12*
Doppler	
Positive	18
Negative	116

\*Doppler negative.  
D-PPG, digital photoplethysmography.

ability score, the modern D-dimer test, and either computed strain gauge plethysmography or D-PPG can safely exclude DVT in outpatients and will be the most cost effective approach at a district general hospital and should now be the initial diagnostic step.<sup>5</sup>

R Sinharay

Tameside General Hospital, Ashton-under-Lyne, UK; ranjitsinharay@hotmail.com

G Strang, D Bird

Royal Glamorgan Hospital, Llantrisant, Mid Glamorgan

#### References

- Walsh K, Kelaher N, Long K, *et al*. An algorithm for the investigation and management of patients with suspected deep venous thrombosis at a district general hospital. *Postgrad Med J* 2002;78:742-5.
- Flanagan DEH, Creasy T, Thomas P, *et al*. Computed assisted venous occlusion plethysmography in the diagnosis of acute deep vein thrombosis. *Q J Med* 2000;93:277-82.
- Tan YK, da Silva AF. Digital photoplethysmography in the diagnosis of suspected lower limb DVT: is it useful? *Eur J Vasc Endovasc Surg* 1999;18:71-9.
- Wells PS, Anderson DR, Rodger M, *et al*. Excluding pulmonary embolism at the bedside without diagnostic imaging: management of patients with suspected pulmonary embolism presenting to the emergency department by using a simple clinical model and D-dimer. *Ann Intern Med* 2001;135:98-107.
- Sinharay R. Pleuritic chest pain and hypoxia—a diagnostic dilemma. *Br J Cardiol* 2002;9:589.

### Dorsalis pedis arterial pulse: palpation using a bony landmark

We read with interest the article by Mowlavi *et al* and agree that the palpation of pedal pulses is a useful clinical tool.<sup>1</sup> Therefore any attempt to standardise the technique of palpation must be applauded. Using the "navicular" method, the authors palpated 78% of pulses in a group of patients under the care of the surgical team. Unfortunately they do not describe the past medical history of this cohort. If their patients have no past medical history

of peripheral vascular disease then their method of palpation is less sensitive than previous studies which detect the pulse in 86.2%–96.9% of healthy patients.<sup>2,3</sup> It would have been useful to know the proportion of arteriopathies in their study. We routinely palpate the dorsalis pedis pulse on a line joining the midpoint of the malleoli to the first webspace. Although this involves soft tissue landmarks, it is nevertheless easy to identify and palpate the pulse on this line. The navicular, in contrast, is not an easy bone to palpate, especially in the oedematous or deformed foot. While they demonstrate a new landmark to readily palpate the dorsalis pedis pulse, we do not feel that they have demonstrated any improvement in either accuracy or ease of use compared with a more traditional landmark.

J Hobson, C Bicknell, N Cheshire

St Mary's Hospital, London, UK; jchobson@doctors.org.uk

#### References

- Mowlavi A, Whiteman J, Wilhelmi BJ, *et al*. Dorsalis pedis arterial pulse: palpation using a bony landmark. *Postgrad Med J* 2002;78:746-7.
- Meade TW, Gardner MJ, Cannon P, *et al*. Observer variability in recording the peripheral pulses. *Br Heart J* 1968;30:661-5.
- Ludbrook J, Clarke AM, McKenzie JK. Significance of absent ankle pulse. *BMJ* 1962;i:1724-6.

## BOOK REVIEWS

The reviewers have been asked to rate these books in terms of four items: readability, how up to date they are, accuracy and reliability, and value for money, using simple four point scales. From their opinions we have derived an overall "star" rating: \* = poor, \*\* = reasonable, \*\*\* = good, \*\*\*\* = excellent.

### Get through MRCP Part 1: 1000 MCQs and Best of Fives.

By Una Coales. (Pp 310; £19.95.) Royal Society of Medicine Press, 2002. ISBN 1-85315-526-8.\*\*\*

The MRCP examination is hard and daunting to get through but the better prepared the candidate then the chance of success early on in either the first or second attempt is obviously much higher. Part 1 is more difficult than part 2 because it is largely theoretically based and revision is tedious and dependent on reading books whereas the part 2, even the written paper, requires clinical knowledge and clinical skills. However, part 1 can be made much easier to prepare for if candidates base their revision around practice questions. There is no point just sitting in the library reading endless textbooks from cover to cover. The MCQ books on the market have been able to give candidates the opportunity to test their knowledge and then to "read around" the subject areas at which they do badly.

With so many books available, one might ask "why produce another?" This is a very pertinent question at the moment as the Royal College of Physicians has decided to drop MCQs from the examination. There will still be two papers, one of which will continue

to be the recently introduced “best of five” and the other will be “n from many”.

Una Coales's book contains six complete mock examinations (each with an MCQ paper and a “best of five” paper). The MCQs in this and other books will not be wasted as they still provide an ideal opportunity to “test one's knowledge” on a syllabus that will largely remain unchanged.

This book offers a fairly comprehensive range of both theoretical and practical questions that are marginally easier than the examination questions—good in the early days of revision when confidence building is essential and reassurance needed that it is possible to pass! It is the first book we have seen to address questions on ethics. However, there are some questions which are outside the MRCP syllabus—a fact that may relate to the background of the author. Each question is criterion referenced with the star ratings giving an indication of the degree of difficulty and what percentage of candidates are expected to get it right. The book's suggested pass marks of 68%–76% for MCQs and 76%–78% for “best of five” reflects the abolition of the negative marking.

The best books available have answers that come complete with explanations; here they are not always present and those that are could be longer and more comprehensive. The MRCP examination as a whole has undergone radical changes to ensure equity and fairness and the candidates of the future need as much practice as possible. All these books, this one included, are there to help.

**E A Freeman, H Nair, J Kitson**  
St Woolas Hospital, Newport, UK

## Review of Psychology for the MRCPsych. 2nd Ed.

By Marcus Munafo. (Pp 198; £18.99.) Arnold Publishers Ltd, 2002. ISBN 0-340-80911-6.\*\*\*

Any book which can increase the chances of passing membership examinations would be a welcome addition to a potential candidate's reading list. Marcus Munafo's *Psychology for the MRCPsych* seems, by its mere title alone, to be explicit in doing this. However, potential readers would do well to heed Munafo's own words of caution in his introduction to this second edition.

The breadth and depth of knowledge required for the written examination can be a significant hurdle, especially if that knowledge is required in specialist areas which may not have previously been taught to any great degree at medical school. Psychology is one of these areas which takes up a significant amount of the college's exam curriculum. Potential future members of the college can therefore expect to be asked a large number of psychology and human development questions and a grounding of knowledge in these areas is a basic necessity.

Munafo's book sets about addressing this task with some aplomb. It seems to assume no, or at least very little, prior knowledge of the subject, and all major areas are covered adequately enough for the reader to feel confident that they have a grasp of the breadth of the subject. The book's close adherence to the curriculum makes it easy to follow for revision purposes. It is clearly and accurately written about the major concepts involved and its

conciseness is to be commended (time can seem in very short supply in the run-up to examinations!). There are frequent summaries throughout the text. However this brevity means that some readers may struggle to fully understand some of the ideas without referring to a more discursive, though lengthier text.

To give the author credit, he identifies that his book is not a replacement for a more traditional psychology text or lecture course, but a bridge between the requirements of the exam and the textbooks that currently exist on this topic and that the candidate will have to do a substantial amount of independent work. It is particularly helpful for revision.

The expanded chapter on neuropsychology helps keep the book up to date, and the inclusion of examples of individual statement questions is in keeping with the recent changes to the exam. These should help give the candidate a flavour of what may be asked in the examination but should by no means be seen as exhaustive.

Overall the book scores highly as a basic introduction and a quick reference guide. It should prove very useful as a revision tool for any part 1 candidate, though those wishing to ensure high marks in this section of the paper would do well to listen to the author's own words and look at additional material elsewhere.

**D Mackintosh**

Specialist Registrar, Leicester General Hospital,  
Leicester

**T Friedman**

Consultant Liaison Psychiatrist, Leicester General  
Hospital, Leicester

- 19 Zwarardemaker H. Der Verlaustau hohen Tönen mit zunehmendem Alter. Ein neues Gesetz. *Arch Ohr Nas und Kehlkopfheilkunde* 1874;32:53–6.
- 20 Kiessling L, Pichora-Fuller MK, Gatehouse S, et al. Candidature for and delivery of audiological services: special needs of older people. *Int J Audiol* 2003;42:2S92–101.
- 21 Arnesen AR. Presbycusis—loss of neurons in the human cochlear nuclei. *J Laryngol Otol* 1982;96:503–11.
- 22 Pichora-Fuller MK, Souza PE. Effects of aging on auditory processing of speech. *Int J Audiol* 2003;42:2S11–16.
- 23 Luxon LM. The anatomy and pathology of the central auditory pathways. *Br J Audiol* 1981;15:31–40.
- 24 Jennings CR, Jones NS. Presbycusis. *J Laryngol Otol* 2001;115:171–8.
- 25 Gates GA, Myers RH. Genetic associations in age-related hearing thresholds. *Arch Otolaryngol Head Neck Surg* 1999;125:654–9.
- 26 Ensink RJ, Camp GV, Cremers CW. Mitochondrial inherited hearing loss. *Clin Otolaryngol* 1998;23:3–8.
- 27 Wright A, Forge A, Kotcheva B. Ototoxicity. In: Booth JB, eds. *Scott Browns otology*. Vol 3. *Otology*. Oxford: Butterworth Heinemann, 1997:3/20/1–2/20/31.
- 28 Lim DP, Stephens SDG. Clinical investigation of hearing loss in the elderly. *Clin Otolaryngol* 1991;16:288–93.
- 29 Schacht J. Aminoglycoside ototoxicity: prevention in sight? *Otolaryngol Head Neck Surg* 1998;118:674–7.
- 30 British Medical Association and Royal Pharmaceutical Society of Great Britain. *British National Formulary* 50, Sep 2005. London: British Medical Association and Royal Pharmaceutical Society of Great Britain, 2005.
- 31 Brors D, Bodmer D. New aspects of inner ear research. *Br J Hosp Med* 2004;65:392–5.
- 32 Gates GA, Cobb JL, D'Agostino RB, et al. The relation of hearing in the elderly to the presence of cardiovascular risk factors. *Arch Otolaryngol Head Neck Surg* 1993;119:156–61.
- 33 Brown CG, Gatehouse S, Lowe GD. Blood viscosity as a factor in sensorineural hearing impairment. *Lancet* 1986;i:121–3.
- 34 Rosen S, Olin P. Hearing loss and coronary heart disease. *Arch Otolaryngol Head Neck Surg* 1965;82:236–43.
- 35 Rosen S, Olin P, Rosen HV. Dietary prevention of hearing loss. *Acta Otolaryngol* 1970;70:242–7.
- 36 Yamazaki T, Ogwa K, Imoto T, et al. Senile deafness and metabolic bone disease. *Am J Otol* 1988;9:376–82.
- 37 Tay HL, Ray N, Ohri R, et al. Diabetes mellitus and hearing loss. *Clin Otolaryngol* 1995;20:130–4.
- 38 Hall SJ, Kerr AG, Varghese M, et al. Deafness in hypothyroidism. (Abstract). *Clin Otolaryngol* 1985;10:292.
- 39 Gates GA, Karzon RK, Garcia P, et al. Auditory dysfunction in aging and senile dementia of the Alzheimer's type. *Arch Neurol* 1995;52:626–34.
- 40 Anonymous. Hearing problems in elderly people: implications for services. *Lancet* 1987;i:1181–2.
- 41 Weinstein BE. Geriatric hearing loss: myths, realities, resources for physicians. *Geriatrics* 1989;44:42–60.
- 42 Pichora-Fuller MK. Cognitive aging and auditory information processing. *Int J Audiol* 2003;42:2S26–32.
- 43 Mulrow CD, Aguilar C, Endicott JE, et al. Quality of life changes and hearing impairment: a randomised trial. *Ann Intern Med* 1990;113:118–94.
- 44 Keller BK, Morton JL, Thomas VS, et al. The effect of visual and hearing impairments on functional status. *J Am Geriatr Soc* 1999;47:1319–25.
- 45 Arlinger S. Negative consequences of uncorrected hearing loss—a review. *Int J Audiol* 2003;42:2S17–20.
- 46 Sekuler R, Blake R. Sensory underload. *Psychology Today* 1987;21:48–53.
- 47 Gennis V, Garry PJ, Haaland KY, et al. Hearing and cognition in the elderly. *Arch Intern Med* 1991;151:2259–64.
- 48 Fisch U. Management of sudden deafness. *Otolaryngol Head Neck Surg* 1983;91:3–8.
- 49 Kveton JF. Evaluation and management of acoustic neuroma. *Current Opinion in Otolaryngology and Head and Neck Surgery* 1993;1:53–63.
- 50 Snik AFM, Mylanus EAM, Cremers CWRG. Implantable hearing devices for sensorineural hearing loss: a review of the audiometric data. *Clin Otolaryngol* 1998;23:414–19.
- 51 Izumikawa M, Minoda R, Kawamoto K, et al. Auditory hair cell replacement and hearing improvement by Atoh1 gene therapy in deaf mammals. *Nature Medicine*. 2005;11: 3, 271–6.
- 52 Ito J, Kojima K, Kawaguchi S. Survival of neural stem cells in the cochlea. *Acta Otolaryngol* 2001;121:140–2.

## CORRECTION

doi: 10.1136/pgmj.2003.006577corr1

Following the decision of the General Medical Council's Fitness to Practise Panel,<sup>1</sup> the publisher is issuing a correction that affects the authorship of a letter published in the *Postgraduate Medical Journal*.<sup>2</sup> Following the panel's decision, this paper should now always be cited as: Bird AD. Cost effective strategy for a safe diagnosis of deep vein thrombosis at a district general hospital. *Postgrad Med J* 2003;79:363.

Ms Bird acknowledges the help of Dr J I G Strang, Dr R Winter, and Mr R Hibberd.

1 [http://www.gmc-uk.org/concerns/decisions/search\\_database/ftp\\_panel\\_sinharay\\_20051122.asp](http://www.gmc-uk.org/concerns/decisions/search_database/ftp_panel_sinharay_20051122.asp)

2 Sinharay R, Strang G, Bird D. Cost effective strategy for a safe diagnosis of DVT at a district general hospital. *Postgrad Med J* 2003;79:363.