seen in the rooms overlooking the courtyard; by a slot device the whole window may be made to disappear into the wall thus giving the patient the maximum amount of fresh air. Unfortunately this device was not permitted by the authorities in those rooms facing the street. By a clever appliance each bed can be raised on to trolley wheels; the nurse is thus enabled to move the patient to any part of the room without undue disturbance. A telephone is installed in each room, and it is left to the doctor to decide as to whether it is to be available to the patient, and if so, whether both out-going and incoming calls may be made.

To avoid the usual noise of bells, a system of coloured lights has been installed to attract the attention of the nurses or attendants.

There are some luxurious suites, comprising patient's bedroom, sitting-room, nurse's bedroom, a bathroom, and in some cases a kitchen.

While I was walking through the building I was interested to note the name of one of the visiting doctors being flashed on a board on every landing. I learnt that this arrangement was to notify the doctor concerned that he was required in some other part of the building, thereby eliminating the constant ringing of bells on each floor. As the number of visiting men increases, I understand that each will be given an identifying number instead of the name appearing on the frame. This arrangement impressed me very much.

An interesting innovation is a fully equipped dispensary. There is a pharmacopœia which has been carefully selected by a group of medical men and which covers most of the requirements of the prescriber. Medicines ordered from this pharmacopœia are free to the patient, and this even includes all hypodermic injections, narcotics, &c.

To turn to the domestic side, there is a magnificent kitchen, equipped with almost every known device for cooking and for the cleansing of utensils, under the supervision of a well-known chef from Paris. Electric lifts carry the food to the different floors. There is quite a nice dining-room at the disposal of the visiting staff, where both lunch and dinner may be obtained at a reasonable cost, and in which, I believe, on certain evenings interesting papers are read.

I concluded my visit by a glimpse into the boiler rooms which give some idea of the immensity of the heating system, sterilization plant, &c., in this building.

I think it is well worth while for our readers to pay a visit to this clinic, where I have no doubt the authorities will be delighted to show them over. While each doctor will form his own opinion as to the suitability of this type of home for his patients, I think there can be no doubt that he will think it exemplifies a new era in the history of nursing homes, embodying as it does the advantages of a well-equipped hospital which hitherto have been more or less available only to the public hospital patient.—[ED.]

Correspondence.

To the Honorary Editor.

SIR,—The note in your current number on the subject of the treatment of liver abscess by the old-fashioned open method seems to me to require some sort of criticism.

In the first place, the suggestion that this method provided the best means of economizing time so as to ensure that the patient could travel by a certain date is open to very serious question. The open method has had, even in skilful hands, a

very serious mortality, and time to recover from the operation is all too short if only three weeks is allowed.

The suggestion that the condition should be treated by drainage at all is an indication that the operator and author of the paper are of opinion that the liver abscess is a condition comparable to a pyogenic abscess, whereas, in fact, there is no resemblance whatsoever to this latter condition. The pathology of the liver abscess is exactly comparable to that of the gumma and of the cold abscess, and in these days, no one would think of

draining a cold abscess as a method of treatment.

When one looks at a liver abscess one finds that there is towards the circumference a very considerable hyperæmia; internal to this there will be a layer of amœbæ, internal again there will be tissue in all stages of destruction; and, finally, the contents consist of debris, of dead cells, &c., and are—and this is a very important point—absolutely sterile. Taking the analogy of a pleural effusion which nowadays is allowed to be absorbed, there is no harm whatever to the individual in allowing this material, the contents of the so-called abscess, to be absorbed. The absorption can be procured by the use of emetine, which is a drug that can only be described in its action as miraculous. Its *modus operandi* is that by killing off the amœbæ the hyperæmia, which has been caused by the attempt of Nature to localize and fight the invading organism, disappears with great rapidity and then the way is open for the absorption of the dead contents of the abscess cavity. The time taken for an abscess to disappear under treatment by emetine is very short, as I saw in the first case I treated wholly by injections of emetine. This patient's abscess was pointing below the right costal margin and in three days the area, which could not be touched owing to extreme tenderness, could be palpated and pressed as freely as though no abnormality had ever existed. Another case reported in the *Lancet* of December 20, 1924, was discharged from hospital in a week. My last case was a man who was shown at two Medical Societies, and who came under my care as a patient in St. Thomas's

Hospital. He was given a grain of emetine on a Friday afternoon, and a grain on the following two days, and was seen once more on the following Friday, when he made the remark that, after the first dose of emetine, travelling in a bus no longer caused pain or inconvenience, and, further, that so quickly did his condition improve that he did not lose any time from his work (which was sedentary) and which he had been on the point of giving up when he came for treatment to the hospital. Apart from these cases I have had several others, all of which have recovered without surgical interference of any sort or kind, and as I have, in my comparatively small experience, 100 per cent. of success, I urge again the necessity for treating amœbic hepatic abscess with emetine, and emetine only as a modern, a safe and satisfactory method. To understand the astounding result that one grain of emetine can produce I would draw attention to cases of liver abscess which had ruptured through the lung and which, before coming under treatment, had coughed up liver abscess pus for a considerable period. One of these cases certainly ceased to cough up liver abscess pus within four hours of an injection of one grain of emetine. Several other cases ceased very rapidly, but I have no definite record as to the lapse of time between the injection and the cessation of the sputum.

In conclusion I would say that the record you publish demonstrates the necessity for an accurate knowledge of the pathology of disease before treatment is undertaken.

Yours faithfully,

London, W.I.

V. S. HODSON.

Extracts from Letters from Post-Graduates.

"I have returned from the Course at Bath. It was most instructive and admirably organized.

"I have only two adverse comments:—

"(1) With the exception of two lectures, it was not possible for all those in the room to hear what was said unless seated close to the lecturers.

"(2) There was scarcely an oppor-

tunity of seeing or examining early cases with rheumatic manifestations. Nearly every case had advanced arthritic changes.

"Probably early material was not available in the hospital, but it might perhaps have been obtainable from outside sources. I greatly enjoyed and profited by the Course."

"First of all, the men at Bath are to be congratulated on the excellent manner in