these reasons it is advisable to give it in the following manner, and to limit the dosage strictly. Atrophan should be used in doses of $7\frac{1}{2}$ gr. three times a day, on three days a week—say Monday, Wednesday and Friday, and should never be given for more than three weeks in all, in any period of three months. Used in this way it is devoid of risk and will produce good results. The mode of action is roughly that the dose on Monday causes all the uric acid present in the blood to be excreted so that some of the uric acid deposited in the tissues is mobilized or forced into solution in the blood during Tuesday. This quantity is again excreted during Wednesday, so that a further quantity of uric acid is mobilized on Thursday and so on. The effect of atophan is augmented by increasing the intake of fluid. Salicylates act in a similar manner but are much less effectual and not nearly so toxic. They may be given over a long period continuously, but probably the best salicylate to use for this purpose is diuretin.

There are many other medicines with reputations as diuretics, but they do not seem to require special mention, because they do not possess any special attributes which render them more useful than those which have been discussed.

ANÆMIA.

By H. V. MORLOCK, M.D.

THE investigation of a patient suffering from a grave anæmia demands a careful history, physical, and pathological examination.

Whatever may be the cause of the anæmia, the symptoms common to all severe degrees of anæmia will be present, i.e., weakness, lassitude, lack of concentration, dyspnœa, palpitations, œdema, often dyspeptic symptoms of varying severity, nevertheless a careful history may elicit other symptoms which may suggest the cause of the anæmia.

Marked cardiac symptoms particularly if fever is present may suggest the possibility of an infective endocarditis, a productive cough, and loss of weight. Phthisis or new growth of the lung, or if the sputum is foul, pulmonary suppuration.

Marked gastro-intestinal symptoms, the possibility of a bleeding peptic or malignant ulcer.

Marked urinary symptoms, the possibility of a chronic nephritis; symptoms referable to the central nervous system would suggest that the anæmia was one which is often associated with central nervous systemic lesions, namely pernicious anæmia.

In female patients the symptoms of pregnancy would demand special attention.

The past history may reveal a recent acute infection such as typhoid, rheumatic fever, or other fevers which are known to cause anæmia.

A recent large acute hæmorrhage, or repeated small chronic hæmorrhages (piles) may be found in the past history. The story of a recent faint may arouse the suspicion of an intestinal or gastric hæmorrhage which was passed down the alimentary tract.

Past attack of anæmia from which recovery was made.

Past attacks of jaundice.

Past residence in tropical or subtropical countries may raise the possibilities of sprue, malaria.

Past venereal disease.

Family history may reveal that relations suffer from anæmia ; pernicious anæmia does occur in family. Acholuric jaundice is usually a familial disease.

As important as the history is the clinical examination of the patient.

The colour of the patient may be suggestive of a definite disease. The lemon tint of pernicious anæmia. The café-au-lait of malignant endocarditis. The pale and pigmented patient of Addison's disease.

The pale and puffy appearance of chronic nephritis. The pale emaciated appearance of malignant disease of the alimentary tract. The biscuit colour of the patient with simple hypochromic microcytic anæmia.

The earthy colour of the advanced phthisis. The typical chlorotic appearance. The pale slight jaundiced appearance of the acute or chronic hæmolytic anæmias.

Starting the **systematic examination** of the patient, the cranial nerves will reveal nothing. The examination of the retina may show hæmorrhage, which may occur in any anæmia but are more commonly found in pernicious anæmia, but may be seen in leukæmia or endocarditis. Subconjunctival hæmorrhages may occur in the same condition.

The **tongue** may reveal a superficial glossitis which may be associated with pernicious anæmia. Sprue or hypochromic microcytic anæmia. An ulcerating angina may be present in acute leukæmia, occasionally in Hodgkin's disease.

Enlarged glands in the cervical region and/or in the axilla and/or groins may suggest lymphatic leukæmia or Hodgkin's disease. A hard gland in the left supra-clavicular fossa malignant disease of the stomach.

The finger-nails may show the spoon-shaped nail associated with hypochromic microcytic anæmia, or the fine clubbing of malignant endocarditis, or the narrow clubbed nail of phthisis, or the broad coarse clubbing of pulmonary suppuration.

In any severe anæmia the **heart** may be found to be dilated, a soft systolic and often a diastolic murmur will be heard about the mitral area, and these signs may give rise to difficulty with regard to the differential diagnosis from malignant endocarditis. The murmurs in this later condition are often harsher than the simple hæmic murmurs, and the occurrence of an aortic diastolic would definitely point to an endocarditis.

The examination of the **lungs** may reveal signs of phthisis, pulmonary suppuration or new growth. If these diseases are responsible for a severe anæmia they will be advanced enough to give very marked signs in the lungs.

Abdominal examination may reveal an enlarged spleen, with or without hepatic enlargement, thus suggesting the possibility of pernicious anæmia, the leukæmias, Hodgkin's disease, acholuric jaundice, Gaucher disease, splenic anæmia, infective endocarditis, chronic malaria, sprue, syphilis or tuberculosis. Hepatic enlargement alone should raise the suspicion of secondary malignant disease. Occasionally a tumour may be found associated with the stomach, colon or kidneys. The presence of ascites might be associated with severe anæmia or might be caused by an underlying malignant condition, or nephritis. The pregnant uterus would be obviously sought for if the history suggested pregnancy.

Rectal, vaginal and testicular examination may give the clue to a severe anæmia. Before leaving the clinical examination of the patient, the whole of the skin should be carefully inspected for purpuric spots which can occur in any blood disease. Occasionally leukæmic infiltration will be found in the skin as bluish flat papules. The urine should be examined for evidence of nephritis and bilirubin.

The **temperature** will seldom give much help, as any severe anæmia may run a low grade of fever, while the acute hæmolytic may run a high temperature, as will the one condition which it may be compared with, namely, infective endocarditis. The leukæmia will also often show marked pyrexia. Very occasionally the Hodgkin's disease will give rise to the typical Pel Ebstein fever.

In those cases **where the anæmia is secondary** to disease of organs other than the blood-forming organs, a careful history and clinical examination will almost always point to the organ which is at fault, and this organ must be investigated by the means at our disposal. The one notable exception to this fact is malignant disease of the alimentary tract, particularly the stomach, which occasionally gives rise to no other symptoms than a severe anæmia. Occasionally because of the cardiac signs and fever which accompany a hæmolytic anæmia, difficulty arose in the differentiation of this condition from an infective endocarditis. The differential diagnosis of these conditions will be dealt with later.

There is now left the investigations which are necessary to differentiate the anæmias which are produced by the diseases of the blood-forming organs themselves.

In these cases the first investigation is a **complete blood-count**. Consideration of the findings as regards the white cell-count and differential count will confirm or rule out a typical myelogenous or lymphatic leukæmia as a cause of the anæmia.

Consideration of the red cell-count and hæmoglobin percentage will show the degree of anæmia which is present and also which of these components of the blood is most reduced, but this investigation alone will not help to differentiate the anæmias unless it is studied in conjunction with the findings on the size of the red cells. The size of the red cells can be measured by the Price Jones curve or the hæmocrit, or the halometer. On this size of the red cells, Wintrobe has based his classification of anæmias.

(1) The **macrocytic anæmias** which are characterized by an increase in the mean size of the red cell and also an increase in its hæmoglobin content; such macrocytic anæmias are pernicious anæmia, sprue, and the anæmia of pregnancy.

(2) The **normocytic anæmias** which include cases of acute blood loss, anæmias of the aplastic or semi-aplastic type and which are characterized by a reduction in the red cell-count without any change, or at most only a slight increase in the size and hæmoglobin content of the red corpuscles.

(3) The **simple microcytic anæmias** which include the most commonly observed form of anæmia, the anæmia associated with chronic infection and intoxications and carcinomata without bleeding. There is a moderate reduction in the corpuscular volume and hæmoglobin, but at no time is this disease as great as the reduction in the number of red cells.

(4) The **microcystic hypochromic anæmia**, which includes the anæmia of chronic hæmorrhage and simple achlorhydric and allied anæmias, and which is distinguished by a marked reduction in the volume of the red corpuscles and a decrease in the hæmoglobin content, which is even more marked than the decrease in size.

From a study of this classification it will be seen that once the complete blood-count and a Price Jones curve or a hæmocritic measurement has been made, one is able to classify the anæmia and thus reduce the differential diagnosis.

Thus, if the case under consideration should fall into the classification of

macrocytic anæmia, the differential diagnosis would then be narrowed to a pernicious anæmia, sprue, the anæmia of pregnancy, and occasionally the anæmia associated with malignant disease of the stomach.

The history will be of definite help in differentiating sprue and the anæmia of pregnancy from the other two conditions, but may give very little help in the diagnosis between pernicious anæmia and a carcinoma of the stomach. Additional pathological investigation : The fractional test meal. This will show an achlorhydria in practically 100 per cent. of cases of pernicious anæmia. Achlorhydria is usually present in sprue but not always ; it may or may not be present in the anæmia of pregnancy ; it is usually, but not necessarily, present in a carcinoma of the stomach.

The **Van den Bergh test** will give an indirect positive in pernicious anæmia, sprue, and the anæmia of pregnancy, but not usually in the carcinoma of the stomach.

The **reticulocyte count** will be low in under-treated cases of sprue and pernicious anæmia, but will usually be high in the anæmia of pregnancy.

The carcinoma of the stomach will almost certainly give radiological evidence of its presence when examined by a barium meal. Further, in this condition, the test for occult blood in the stools will often be repeatedly positive, but this may also occur in pernicious anæmia if there is bleeding into the alimentary tract.

The findings of signs of subacute degeneration of the cord may occur in pernicious anæmia and sprue, but not in the other two conditions.

The therapeutic test of liver treatment.—Pernicious anæmia, and sprue, will recover on this treatment, and the reticulocytes will produce their critical rise about the eighth day. The anæmia of carcinoma of the stomach may show some improvement for a while ; the anæmia of pregnancy will not improve.

If the anæmia falls within the **normocytic anæmias**, one has to consider acute blood loss, malaria, and aplastic anæmias.

In the majority of cases of acute blood loss the history is sufficient to give the diagnosis as to the cause of the anæmia, but occasionally a bleed into the intestinal tract will give rise to no other symptom than an acute anæmia, and the occult test in the stools and radiological examination of the alimentary tract will have to be done before the cause is found.

Thrombocystopænic purpura may give rise to an acute blood loss, and so produce the picture of a normocytic anæmia. In these cases the history of previous bleeding may be obtained, also the presence of marked purpura would give some confirmation. A platelet count would reveal a great reduction in their number, 10,000 to 50,000 ; the normal being 300,000 to 500,000.

Hæmophilia might be responsible for an acute blood loss. In this case the family history would be diagnostic, also the examination of the coagulation time would show a very marked increase.

Malaria would be suggested by the history and would be confirmed by the finding of the parasite in the blood-film.

The **aplastic anæmia** would be found in a child or adolescent ; the anæmia would be marked and progressive in spite of treatment. The reticulocytic count would be permanently very low.

If the anæmia falls into the simple microcytic group, then the carefully taken history and physical examination will reveal the cause of the anæmia.

If the anæmia falls into the **microcytic hypochromic** group, the condition to be considered would be repeated chronic hæmorrhages, and simple achlorhydric anæmia.

The causes of the chronic hæmorrhage may be due to a carcinoma of the alimentary tract; a peptic ulcer, or piles, and these cases must be investigated by a complete radiological examination of the alimentary tract, and if this reveals nothing, a sigmoidoscopic examination, and examination of the stools for occult blood.

Other causes of repeated bleedings may be thrombocystopænia, or hæmophilia.

The simple **achlorhydric anæmia** is usually found much more commonly in females than males, between 30 to 60 years of age. They may present a typical biscuit colour. A superficial glossitis is always present, occasionally also some dysphagia. The finger-nails are sometimes spoon-shaped, signs of involvement of the central nervous system may be present; an achlorhydria is a constant finding. Treatment with large doses of iron is almost a diagnostic test.

Some American authors maintain that chlorosis is closely related if not identical with simple achlorhydric anæmia.

There is still another group of anæmias to be considered, that is, the **hæmolytic anæmias**. Witts has divided these into two groups: the chronic acquired hæmolytic anæmias, and the acute hæmolytic anæmias.

The anæmias are characterized by the presence of jaundice due to the increased hæmolysis. The blood-picture will vary from time to time according to the extent to which the hæmolysis is taking place at the time that the blood-count is done. The reticulocytic count is high, and if an acute hæmolysis is taking place the white cell-count may be raised and even immature white cells may appear in the blood-stream and thus give rise to the suspicion of a lukæmic condition, but the high reticulocytic count in the hæmolytic anæmia will help in the diagnosis.

In the acute form, the clinical picture is that of an acute infection with anæmia, the anæmia producing cardiac signs which may be suggestive of an infective endocarditis. Repeated negative blood-cultures give some assistance in ruling out this later condition, while a positive blood-culture would confirm it.

Acholic family jaundice is characterized by the family history, the repeated attacks of jaundice, the enlarged spleen, and the evidence of increased fragility of the red cells when tested with hypotonic saline.

If an anæmia is present and glandular enlargement is also present, in order to arrive at a diagnosis, it is often necessary, particularly in Hodgkin's disease, to remove one of the glands for biopsy.

A Wassermann reaction should always be taken in a case of severe anæmia, and if no other cause for the anæmia is found, and other signs of syphilis are present, anti-syphilitic treatment should be given; but it must be remembered that a positive Wassermann reaction may be present although the patient is suffering from anæmia due to some other cause.