

Supplementary data

Odds of testing positive for IgM.

Variables	OR	95% CI	P-value
PCR-positive	5.519	1.708-17.828	0.004
Recovery	0.289	0.037-2.241	0.235
Symptomatic	13.368	1.601-111.642	0.017

CI, confidence interval; OR, odds ratio; PCR, polymerase chain reaction

Odds of testing positive for IgG.

Variables	OR	95% CI	P-value
PCR-positive	2.227	0.858-5.780	0.100
Recovery	4.063	2.501-6.601	<0.001
Symptomatic	1.081	0.110-10.595	0.947

CI, confidence interval; OR, odds ratio; PCR, polymerase chain reaction

Figure 1. The comparison of antibody level in PCR-positive, symptomatic and recovered subjects between players and officials.

Variables	Groups	IgM level	P-Value	IgG level	P-Value
PCR-positive (n=29)	Player (n=24)	0.61 (1.4)	0.953	0.95 (1.9)	0.595
	Official (n=5)	0.67 (0.8)		4.94 (8.0)	
Recovered (n=112)	Player (n=92)	0.25 (0.2)	0.006	1.53 (3.0)	0.025
	Official (n=20)	0.45 (0.4)		3.43 (4.7)	
Symptomatic (n=7)	Player (n=6)	0.93 (0.8)	0.317	1.15 (1.6)	0.614
	Official (n=1)	2.10 (-)		1.10 (-)	

* Data are presented as: mean (SD)