OPEN ACCESS

Diffusely increased ^{18}F -FDG thyroid radionuclide uptake is not always hyperthyroidism

Lijuan Sun ¹, Hui Jen Goh, ¹ Nan Xin Wang ^{2,3}, Sanjay K Verma, ⁴ S. Sendhil Velan, ⁴ Melvin Khee Shing Leow ^{1,5,6}

¹Human Development, Singapore Institute for Clinical Sciences, Singapore

²Clinical Nutrition Research Centre, Agency for Science, Technology and Research (A*STAR), Singapore

³Department of Human Nutrition, University of Otago, Otago, New Zealand

⁴Laboratory of Molecular Imaging, Singapore Bioimaging Consortium, Agency for Science Technology and Research (A*STAR), Singapore

⁵Endocrinology, Tan Tock Seng Hospital, Singapore

⁶Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore

Correspondence to

Dr Melvin Khee Shing Leow, Singapore Institute for Clinical Sciences, Singapore 117609, Singapore; melvin_leow@sics.a-star.edu.sg

Received 21 December 2020

Revised 18 March 2021

Accepted 1 April 2021

INTRODUCTION

In the routine evaluation of the thyroid, imaging is often a useful modality, which gives anatomical and functional characteristics that can aid diagnosis. Thyroid ultrasonography is the principal imaging modality in this regard, whereas CT, MRI and PET scanning are much less frequently ordered for thyroid assessment through thyroid lesions are often discovered via such advanced imaging modalities as incidental findings.¹ Unlike radiologists, most physicians tend to be unfamiliar with radiological features of thyroid disorders detected via these latter forms of scanning.²

CASE DESCRIPTION

A 25-year-old woman who was otherwise overtly healthy and asymptomatic without any history of medical illnesses or chronic medications had volunteered for a research study on brown fat. She underwent whole-body ^{18}F -FDG PET/MR imaging according to a standard protocol. This unexpectedly revealed abnormally increased FDG uptake (SUVmax=6.84) over the anterior neck strangely reminiscent of typical radionuclide ($^{99\text{m}}\text{Tc}$ or ^{131}I) thyroid scans of hyperthyroidism (figure 1A,B). What is the diagnosis, and how would you confirm it?

Answer: A prescan screening thyroid function test revealed a serum-free thyroxine (FT4) of 10.0 pmol/L (RI: 8–20) and serum thyroid-stimulating hormone (TSH) of 4.28 mIU/L (RI: 0.45–4.50). An invaluable clue came from a thyroid function test performed 7 months earlier: FT4=13.2 pmol/L (RI: 8–20) and TSH=2.43 (RI: 0.45–4.50). This implied a progressive decline in FT4 associated with a reciprocal increase in TSH. Thyroid ultrasonography was done next, which revealed a general heterogeneous echotexture with parenchymal hypoechogenicity consistent with thyroiditis (figure 2). Corroborating the ultrasound scan, the nonenhanced T2-weighted MR images that were simultaneously acquired during the PET-MRI fusion scan revealed high and inhomogeneous signal intensities within the thyroid parenchyma suggestive of a diagnosis of Hashimoto thyroiditis (figure 3). Serum antithyroid peroxidase autoantibody exceeded 1000 IU/mL (normal <50 IU/mL), while serum antithyroglobulin autoantibody was 33.55 IU/mL (normal <4.1 IU/mL). Despite the relatively ‘normal’ thyroid function tests, the evidence was supportive of an actively evolving thyroid autoimmunity consistent with early Hashimoto’s thyroiditis. Short ACTH stimulation test excluded concomitant hypocortisolism, a

Figure 1 ^{18}F -FDG-PET scans showing diffusely increased radionuclide uptake by the thyroid gland as shown in the coronal section (A) and the sagittal section (B).

crucial step to avoid precipitating Addisonian crisis during L-thyroxine initiation in those with unsuspected adrenalitis from autoimmune polyglandular syndrome. She was prescribed L-thyroxine at 25 µg/day. The dose was doubled 2 years later, which optimised her energy level and body weight. At a 4-year follow-up visit, she remained euthyroid with serum FT4 of 12.5 pmol/L (RI: 8–16), FT3 of 4.7 pmol/L (RI: 3.5–6.0) and TSH of 1.54 mIU/L (RI: 0.45–4.50).

CONCLUSION

The rising prevalence of autoimmune thyroiditis coupled with an increasing use of ^{18}F -FDG-PET scanning for various indications will continue to generate such incidental findings as highlighted by our case.^{3,4} Although diffusely increased ^{18}F -FDG thyroid uptake is observed in hyperthyroidism such as Graves’ disease, physicians should recognise that diffusely increased ^{18}F -FDG thyroid uptake can also be due to Hashimoto thyroiditis,⁵ which contrasts starkly with diminished $^{99\text{m}}\text{Tc}/^{131}\text{I}$ uptake typical of subacute thyroiditis⁶ and that must be differentiated

Figure 2 Thyroid ultrasound scan revealing a gland with heterogeneous echotexture with diffuse hypoechogenicity in bilateral lobes consistent with thyroiditis.

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Sun L, Goh HJ, Wang NX, et al. *Postgrad Med J* Epub ahead of print: [please include Day Month Year]. doi:10.1136/postgradmedj-2020-139643

Figure 3 MRI of thyroid of the same patient, showing non-enhanced T2-weighted coronal (A) and sagittal (B) images with inhomogeneous high signal intensities within the thyroid parenchyma supportive of Hashimoto thyroiditis.

from analogous scintigraphic features of Graves' disease characterised by diffusely accentuated $^{99m}\text{Tc}/^{131}\text{I}$ and ^{18}F -FDG uptake.

Contributors All the listed authors contributed to the manuscript preparation. LS wrote the first draft of the manuscript and submitted the manuscript. LS, HJG, SKV and SSV collected the data. NXW contributed the data and assisted with the editing of the manuscript. ML was the overall principal investigator and physician scientist in charge who analysed and interpreted the data, managed the patient and contributed to the writing, editing and critical revision of the manuscript. All authors approved the final version.

Funding The study was funded by the Singapore Ministry of Health's National Medical Research Council (NMRC) Clinician Scientist Award (grant ID NMRC/CSA-INV/0003/2015) awarded to MKSL.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Lijuan Sun <http://orcid.org/0000-0002-7650-3400>

Nan Xin Wang <http://orcid.org/0000-0003-1744-5930>

REFERENCES

- 1 Farrá JC, Picado O, Liu S, *et al*. Clinically significant cancer rates in incidentally discovered thyroid nodules by routine imaging. *J Surg Res* 2017;219:341–6.
- 2 Soto GD, Halperin I, Squarcia M, *et al*. Update in thyroid imaging. The expanding world of thyroid imaging and its translation to clinical practice. *Hormones* 2010;9:287–98.
- 3 Agrawal K, Weaver J, Ngu R, *et al*. Clinical significance of patterns of incidental thyroid uptake at (18)F-FDG PET/CT. *Clin Radiol* 2015;70:536–43.
- 4 Agrawal K, Weaver J, Ul-Hassan F, *et al*. Incidence and Significance of Incidental Focal Thyroid Uptake on (18)F-FDG PET Study in a Large Patient Cohort: Retrospective Single-Centre Experience in the United Kingdom. *Eur Thyroid J* 2015;4:115–22.
- 5 Nakamoto Y, Tatsumi M, Hammoud D, *et al*. Normal FDG distribution patterns in the head and neck: PET/CT evaluation. *Radiology* 2005;234:879–85.
- 6 Intenzo CM, dePapp AE, Jabbour S, *et al*. Scintigraphic manifestations of thyrotoxicosis. *Radiographics* 2003;23:857–69.