



OPEN ACCESS

Knowledge and practice of home blood pressure monitoring 6 months after the risk and assessment management programme: does health literacy matter?

Sau Nga FU ,¹ Man Chi Dao ,¹ Carlos K H Wong ,^{2,3}
Bernard M Y Cheung ⁴

► Additional online supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/postgradmedj-2020-139329>).

¹Department of Family Medicine and Primary Health Care, Hospital Authority Kowloon West Cluster, New Territories, Hong Kong

²Department of Family Medicine and Primary Care, Li Ka Shing Faculty of Medicine, University of Hong Kong, Hong Kong SAR, China

³Department of Pharmacology and Pharmacy, Li Ka Shing Faculty of Medicine, University of Hong Kong, Hong Kong SAR, China

⁴Department of Medicine, Li Ka Shing Faculty of Medicine, University of Hong Kong, Hong Kong SAR, China

Correspondence to

Dr Sau Nga FU, Family Medicine & Primary Health Care, Hospital Authority Kowloon West Cluster, N.T., Hong Kong; fsn299@ha.org.hk

Received 5 November 2020

Revised 8 April 2021

Accepted 21 April 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: FU SN, Dao MC, Wong CKH, *et al.* *Postgrad Med J* Epub ahead of print: [please include Day Month Year]. doi:10.1136/postgradmedj-2020-139329

ABSTRACT

Background Little is known whether patients with lower health literacy could retain the practice and knowledge of home blood pressure monitoring (HBPM) after an educational programme.

Methods A cluster randomised controlled trial in five primary care clinics recruited participants with uncontrolled hypertension. Clinics were randomised either to a HBPM group education (Risk Assessment and Management Programme (RAMP-group), or individual counselling of self-management (RAMP-individual). Health literacy was assessed by the Chinese Health Literacy Scale for Chronic Care. Practice and knowledge of HBPM were surveyed by a 10-item HBPM knowledge checklist and patient record review 6 months after interventions. Predictors for regular HBPM and good HBPM knowledge were assessed by multivariate logistic regression models.

Results 287 participants (RAMP-group: 151; RAMP-individual: 136) were follow-up for 6 months. 272 participants completed the knowledge questionnaires (response rate 94.8%). 67.8% of the participants performed HBPM regularly, and there was no statistical difference between both interventions. Age more than 65 (adjusted odds ratios (aOR) 2.58, 95% CI 1.37 to 4.86, $p=0.003$), not working (aOR 2.34, 95% CI 1.10 to 4.97, $p=0.027$) and adequate health literacy (aOR 2.25, 95% CI 1.28 to 3.95, $p=0.005$) predicted regular HBPM. Participants in RAMP-group demonstrated a significant lower body weight than those in RAMP-individual (-0.3 ± 2.0 kg vs $+0.7\pm 1.7$ kg, $p<0.001$). The RAMP-group participants were eight times more likely to have full HBPM knowledge score than the RAMP-individual participants (aOR 8.46, 95% CI 4.68 to 15.28, $p<0.001$).

Conclusion Patients could retain HBPM knowledge better after RAMP-group than RAMP-individual. Older, retired and patients with adequate health literacy were more likely to continue weekly HBPM 6 months after education.

Trial registration number NCT02551393.

BACKGROUND

Several national and international guidelines recommended home blood pressure monitoring (HBPM) in the management of hypertension (HT), particularly for those with suspected discrepancies of blood pressure (BP) reading in office and other

casual situations.^{1 2} Despite the worldwide unsatisfactory BP control in different populations, only around half of the patients with HT perform HBPM regularly.^{3 4} Different meta-analysis studies demonstrated a reduction in patients' BP after interventions to facilitate HBPM.^{5 6} However, HBPM has potential pitfalls in itself, namely inaccurate HBPM devices,⁷ the need for coaching on proper HBPM techniques,⁸ selectively or erroneously report BP values by patients,⁹ and the cost of equipment and maintenance. These may hinder patients from using HBPM.¹⁰ Therefore, some of the clinicians were hesitated to promote HBPM in their practice.^{11 12}

High-quality HBPM could be achieved by education intervention and increasing patient-clinician contact time.¹³ Given the multiple potential sources of inaccuracy in HBPM measurement,¹⁴ the sustainability of high-quality HBPM after an educational intervention may be questionable. Knowledge of HBPM and practice of regular HBPM may be affected by patients' age, sex, educational level and health literacy (HL).¹³

This study performed an interim patient outcome assessment of a cluster randomised controlled trial. The study aims to compare whether patients can retain HBPM knowledge and regular HBPM practice 6 months after two types of multidisciplinary education programmes. The sociodemographic predictors for desirable HBPM outcomes were also assessed.

We hypothesise that more patients who attended group education could retain better HBPM knowledge and regular HBPM than those attended conventional individual counselling 6 months post-interventions. Patients with better HL or higher educational level would have higher compliance with regular HBPM and better HBPM knowledge.

METHOD

Written informed consents were obtained before the patients participated in the study.

Study design

This study is an interim outcome analysis of a cluster randomised controlled trial in five government primary care clinics. The methodology and sample size calculation was reported elsewhere.¹⁵ We included clinics with more than three family physicians, and their number of registered

hypertensive patients is more than 6000 in the past 12 months. Three clinics were allocated to the intervention groups, while two clinics were assigned to the control group by simple randomisation. Patients were recruited if they have uncontrolled HT. Uncontrolled HT was defined as clinic SBP more than 140 mmHg (for those aged <80), more than 150 mm Hg (for those aged ≥80) or DBP more than 90 mm Hg in their most recent two clinic visits. Higher cut-off for uncontrolled HT was adopted for older adults (age ≥80) according to the recommendation of treatment initiation in the Eighth Joint National Committee guideline and the National Institute for Health and Clinical Excellence 2014 guideline.^{1,2}

Control clusters (Risk Assessment and Management Programme-individual, usual care)

According to study clinics' HT management protocol, all patients with uncontrolled HT are invited to the Risk Assessment and Management Programme (RAMP) annually.^{3,4} RAMP consists of an individual 15 min face to face nurse clinic. Clinic nurses provide patients' clinical parameters results, assess patients' overall cardiovascular disease (CVD) risk, counsel on lifestyle modification and refer patients to appropriate allied health services such as physiotherapy, dietitian or occupational therapist. Patients already on HBPM are advised to bring their own HBPM devices for checking machine functions and patients' HBPM technique.

Intervention clusters (RAMP-group)

Patients and their family members were invited to the group educational sessions after the first RAMP nurse clinic. The RAMP-group for 10–15 participants lasted for 2 hours offered the day group (9:00–17:00 hours) or the evening group (18:00–22:00 hours). Every participant received an automatic branchial type HBPM device and a self-administered BP record booklet.^{16,17} Second author MD, who is a family doctor, gave a 30 min seminar about basic knowledge of HT, antihypertensive drugs, CVD risk factors and understanding of home BP and office BP.¹⁸ The participants then watched a 10 min video introducing HBPM procedures. Our clinic nurses then demonstrate how to use the HBPM devices, and all participants practised on their device. Lastly, all participants were divided into groups of 3–4 to practise HBPM with their facilitators, might be doctors, nurses or clinic assistances. The facilitators went through a 10 items checklist of HBPM procedures until the participants got all items correct.¹⁹

Six weeks later, all participants returned their BP devices and attended an individual face-to-face nurse follow-up. Their healthy lifestyle, drug compliance and HBPM practice were reviewed. They were encouraged to buy their own validated HBPM devices and communicate their HBPM records with clinicians.

Baseline assessment

Clinic nurses identified patients with uncontrolled HT. They then asked for their written consents of this study. Trained research assistants then assessed participants' sociodemographic data, HT drug prescription and HL by the Chinese Health Literacy Scale for Chronic Care (CHLSCC). All biochemical markers were retrieved from the computerised patient records. Biochemical outcomes included body weight, fasting blood sugar and low-density lipoprotein (LDL).

Table 1 10-item HBPM knowledge checklist

No		BHS	AHS	RACGP
1.	What should be avoided before taking measurements at least 30 min before measurement? Answer: Do not exercise, smoke or consume foods or drinks containing caffeine (such as tea or coffee)	✓	✓	✓
2.	How long should you rest and relax without any distractions Answer: at least 5 min (time between 5 min and 30 min is acceptable)	✓	✓	✓
3.	What is the dressing requirement during measurement? Answer: Wear loose-fitting and comfortable clothes, Roll up the sleeve to expose your upper arm and wrap the cuff around it.	✓		
4.	What are the appropriate characteristics of table and chair used for HBPM? Answer: Choose a stable table and chair of appropriate height (at heart level) with armrest. How should your back and feet be placed? Answer: Sit comfortably and relaxed with your back supported, keep your feet on the floor and do not cross the legs	✓	✓	✓
5.	Show me your HBPM cuff (Check if the cuff size is appropriate)		✓	✓
6.	Could you show me how you put on your cuff? (Observe: Place cuff in an appropriate position: The lower border of the cuff should be 2 cm above the pit of elbow, which is approximately two finger-breadths. Make sure the tubing is placed at the centre of your arm facing the front and that the sensor is correctly placed.)	✓	✓	✓
7.	Check cuff with appropriate tightness: Pull the end of the cuff so that it is wrapped evenly and firmly around your arm. Check that the tightness of the cuff is proper: you should be able to just slip two fingertips beneath the cuff, near its edge at the top end	✓	✓	✓
8.	Please start to measure now (Observe: Press start button, relax during measurement (no talking/movement))	✓	✓	✓
9.	How many times do you measure? When? Answer: Repeat measurement at least 1 min after complete release of the cuff	✓	✓	✓
10.	Can you write down the readings on this booklet? Record all readings appropriately	✓	✓	✓
Total score (full mark 10)				

AHS, American Heart Association²⁰; BHS, British Hypertension Society²¹; BP, blood pressure; HBPM, home blood pressure monitoring; RACGP, Royal Australian College of General Practitioner²².

Outcomes assessment

Apart from repeating baseline biochemical assessment at 6 months after intake, clinic nurses also asked the 10 items HBPM knowledge questionnaire and the practice of HBPM. HBPM practice was verified by doctors' and nurses' consultation records.

Chinese Health Literacy Scale for Chronic Care

The CHLSCC is a locally validated tool to assess HL among Chinese patients with chronic illness. It displayed a good internal reliability (Cronbach's $\alpha=0.91$).⁶ It assesses remembering, understanding, applying and analysing ability. Participants scored 36 out of 48 or above indicated adequate HL; otherwise, they had inadequate HL.

10-item HBPM knowledge checklist

Table 1 shows the 10-item HBPM knowledge questionnaire. It was developed according to the American Heart

Table 2 Patient's sociodemographic and clinical parameters at baseline

Characteristics	RAMP-individual (N=137)	RAMP-Group (N=152)	Total (N=289)	P value*
Age (year; mean±SD, median)	67.9±10.2, 67	66.1±9.7, 65	67.0±9.9, 66	0.528
Gender (%)				
▶ Male	58 (42.3%)	55 (36.2%)	113 (39.1%)	0.285
▶ Female	79 (57.7%)	97 (63.8%)	176 (60.9%)	
Education (%)				
▶ Primary or below	81 (59.1%)	97 (63.8%)	178 (61.6%)	0.352
▶ Secondary school	48 (35.0%)	51 (33.6%)	99 (34.3%)	
▶ University or above or unknown	8 (5.8%)	4 (2.6%)	12 (4.2%)	
Occupation (%)				
▶ Not employed/housewife	105 (76.6%)	120 (78.9%)	225 (77.9%)	0.637
▶ Working	32 (23.4%)	32 (21.1%)	64 (22.1%)	
CHLSCC				
▶ Total 0–48: mean±SD, median	28.11±13.73, 30.5	35.00±12.76, 39.0	31.70±13.65, 35	<0.001†
▶ Adequate (≥36) (%)	51 (37.2%)	94 (61.8%)	145 (50.2%)	<0.001†
▶ Inadequate (<36) (%)	86 (62.8%)	58 (38.2%)	144 (49.8%)	
Smoking status				
▶ Non-smoker	92 (67.2%)	121 (79.6%)	213 (73.7%)	0.056
▶ Current smoker	7 (5.1%)	5 (3.3%)	12 (4.2%)	
▶ Ex-smoker	38 (27.7%)	26 (17.1%)	64 (22.1%)	
Clinic BP				
▶ Systolic BP (mm Hg, mean±SD)	152±9.81	152.0±10.4	152.1±10.1	0.775
▶ Diastolic BP (mm Hg, mean±SD)	78.4±10.6	80.76±11.9	79.7±11.4	0.087
BMI (kg/m ² ; mean±SD)	26.49±4.31	26.31±4.28	26.39±4.27	0.774
▶ Underweight (<18.5)	4 (2.9%)	3 (2.0%)	7 (2.4%)	0.937
▶ Normal (≥18.50 -<23)	22 (16.1%)	24 (15.8%)	46 (15.9%)	
▶ Overweight (≥23 -<25)	28 (20.4%)	29 (19.1%)	57 (19.7%)	
▶ Obese (≥25)	83 (60.6%)	96 (63.2%)	179 (61.9%)	
Duration of hypertension since diagnosis (years±SD)	10.92±9.32	8.43±8.04	9.62±8.75	0.015†
Uncomplicated hypertension (%)	81 (59.1%)	103 (67.8%)	184 (63.7%)	0.127
Complicated hypertension (%)	56 (40.9%)	49 (32.2%)	105 (36.3%)	

*P value of proportions by Pearson χ^2 tests; continuous variables by Student's t-tests.

†P value <0.05 are bold

BMI, body mass index; BP, blood pressure; CHLSCC, Chinese Health Literacy Scale for Chronic Care; RAMP, Risk Assessment and Management Programme.

Association recommendations,²⁰ the British Hypertension Society²¹ and Royal Australian College of General Practitioners patient education material.²² The first to third items are about preparing themselves, including clothing, rest and avoidance of stimulants before measurement. The fourth to the seventh item is about body positioning and cuff application. The eighth to ninth items is about the frequency of HBPM measurement. The last item is about correct HBPM reading recording. The results are presented as scoring over 10 (full mark). The checklist is administered by face-to-face interview, supplemented by patient's consultation records and telephone interview if patients could not be contacted 6 months later.

HBPM practice questionnaire

Participants reported if they practised HBPM regularly (Yes/No), if they owned their HBPM devices (yes/no), how frequent they measured their BP (daily, weekly, monthly, less than monthly), how many times they measured in a day (once, twice, three times, four times or more). Their answers were verified by authors via the patients' consultation record. HBPM practice was routinely documented by doctors and nurses during patients' quarterly follow-up.

Participants were defined as performing HBPM regularly if they measured more than once weekly.

Statistical analysis

The participants' descriptive statistics were reported as means and SD, or frequencies and percentages as appropriate. This study endpoint was 6 months after the interventions. The primary outcome was the differences in the proportion between participants attained RAMP-individual and RAMP-group having good HBPM knowledge using χ^2 tests. The secondary outcomes included the differences in the proportion between the participants attained RAMP-individual and RAMP-group measuring home BP regularly using χ^2 tests. Subgroup analysis of the participants' outcomes with adequate and inadequate HL was performed because of their baseline difference. The unadjusted OR (aOR) was calculated for the predictors for regular HBPM and for having good HBPM knowledge. The aOR was calculated from the multivariate logistic regression model. The ORs were adjusted by age, sex, educational level, HL level, occupation and whether the participant attended RAMP-group or RAMP-individual. A $p < 0.05$ was considered as

Table 3 Study outcomes at 6 months postinterventions

Outcomes	Adequate HL (total 145)			Inadequate HL (total 144)			P value	P value*	RAMP-Group Total	RAMP-Individual Total	RAMP-Group Total	P value
	RAMP-Individual (n=51)	RAMP-Group (n=94)	P value	RAMP-Individual (n=86)	RAMP-Group (n=58)	P value						
Any change of HT drugs	24 (47.1%)	40 (42.6%)	0.602	29 (33.7%)	23 (39.7%)	0.530		53 (38.7%)	63 (41.4%)	0.633		
▲ Yes												
▲ No	27 (52.9%)	54 (57.4%)	0.213	57 (66.3%)	35 (60.3%)	<0.001		84 (61.3%)	89 (58.6%)	<0.001		
Change of body weight (kg, mean±SD)	0.2±1.7	-0.3±2.0	0.213	1.0±1.7	-0.2±1.9	0.378		0.7±1.7	-0.3±2.0	0.604		
Fasting blood sugar (mmol/l, mean±SD)	5.9±1.1	6.1±1.1	0.205	6.5±1.8	6.2±1.4	0.107		6.2±1.6	6.2±1.2	0.608		
Low-density lipoprotein (mmol/l, mean±SD)	3.1±1.0	2.9±0.8	0.330	2.8±0.9	3.1±0.9			2.9±0.9	3.0±0.8			
HBPM practice												
Own their BP devices (%) yes	44 (86.3%)	74 (78.7%)	0.265	52 (60.4%)	38 (65.5%)	0.539		96 (70.1%)	112 (73.7%)	0.495		
▲ No	7 (13.7%)	20 (21.3%)	0.755	34 (39.5%)	20 (34.5%)	0.845		41 (29.9%)	40 (26.3%)	0.215		
Measure BP at home in the past 6 months (%) Yes	39 (76.5%)	74 (78.7%)	0.216	49 (57.0%)	34 (58.6%)	0.784		88 (64.2%)	108 (71.1%)	0.153		
▲ No	12 (23.5%)	20 (21.3%)		37 (43.0%)	24 (41.3%)			49 (35.8%)	44 (28.9%)			
Frequency of HBPM												
▲ More than once weekly	33 (64.7%)	70 (74.5%)	0.216	45 (52.3%)	29 (50.0%)			78 (56.9%)	99 (65.1%)			
▲ Less than once weekly	18 (35.3%)	24 (25.5%)		41 (47.7%)	29 (50.0%)			59 (43.1%)	53 (34.9%)			
HBPM knowledge												
HBPM Assessment Score (%) (N=272)†	12 (24.1%)	63 (71.6%)	<0.001	15 (19.0%)	35 (63.6%)	<0.001		27 (20.9%)	98 (68.5%)	<0.001		
▲ Full Mark (all correct)												
▲ At least one item incorrect	38 (76.0%)	25 (28.4%)		64 (81.0%)	20 (36.4%)			102 (79.1%)	45 (31.5%)			

* P value of proportions by Pearson χ^2 tests; continuous variables by Student's t-tests.

†17 participants refuse to answer knowledge score 6 month post-intervention

BP, blood pressure; HBPM, home blood pressure monitoring; HL, health literacy; HT, hypertension; RAMP, Risk Assessment and Management Programme.

significant. All analyses were performed using SPSS V.25 (IBM).

RESULTS

A total of 289 participants were recruited and follow-up until 6 months. Two participants from the RAMP-individual passed away before study endpoint, therefore, excluded from the analysis. A total of 272 participants completed the 6 months questionnaires (response rate of 94.8%) except 17 of them refused to answer the HBPM knowledge score. Online supplemental table S showed the details of missing data. Table 2 shows the participants' characteristics. Their mean age was 67.0 (SD 9.9); almost two-thirds were women. Their educational level was low. Around 60% of them attended primary school, and another one-third of them attended secondary school. There were significantly more participants with adequate HL in RAMP-group than those in RAMP-individual. Their baseline clinic BP, body weight, waist circumference, proportions of complicated HT and drug compliance were similar.

Table 3 demonstrates the participants' outcomes at 6 months postintervention. Due to the baseline difference in HL, subgroup analysis of participants with adequate and inadequate HL were performed. Among the participants with adequate HL, there was statistically insignificant difference in change of body weight (BW) after intervention. However, among the participants with inadequate HL, the RAMP-group had mean weight loss (-0.2 ± 1.9 kg), when compared with the RAMP-individual had mean weight gain ($+1.0 \pm 1.7$ kg). Overall, the RAMP-group had statistically more favourable weight change than the RAMP-individual. There was no statistically significant difference in the change in HT drugs, fasting blood sugar level and LDL level.

Most of the participants from the RAMP-group (68.5%) could score full marks in the knowledge score questionnaire, while only 20.9% of the participants from the RAMP-individual could achieve the same score ($p < 0.001$). More than 60% of participants with both adequate and inadequate HL could score full marks in RAMP-group. Only around 20% of the RAMP-individual scored full mark no matter they had adequate or inadequate HL.

Concerning about HBPM practice, there was no difference in proportions of participants in possession of HBPM device, regular measurement of HBPM and measuring at least once weekly between RAMP-group and RAMP-individual. There were more than two-thirds of participants owned their HBPM devices. 71.1% of the RAMP-group participants continued to measure HBPM more than once weekly. The proportions were the highest among the RAMP-group with adequate HL (78.7%) compared with the lowest in the RAMP-individual with inadequate HL (57.0%).

The predictors for regular HBPM and good HBPM knowledge were illustrated in table 4 using binary logistic regression models. In the multivariate model, patients older than 65 (adjusted odds ratio (aOR) 2.58, 95% CI 1.37 to 4.86, $p = 0.003$), who were not working (aOR 2.34, 95% CI 1.10 to 4.97, $p = 0.027$) and with adequate HL (aOR 2.25, 95% CI 1.28 to 3.95, $p = 0.005$) predicted regular HBPM. The only predictor for getting full-score in HBPM knowledge was attended RAMP-group instead of RAMP-individual (aOR=8.46, 95% CI 4.68 to 15.28, $p < 0.001$). Age, sex, educational level, HL, HT duration, and being obese did not predict good HBPM knowledge.

Table 4 Logistic regression of predictors for regular HBPM and good HBPM knowledge

Predictors	Regular HBPM*						Good HBPM knowledge†					
	Reference group	OR	(95% CI)	P values‡	AOR‡	95% CI	P values‡	(95% CI)	AOR‡	95% CI	P values‡	
Age	≥65	1.47	(0.91 to 2.38)	0.116	2.58	(1.37 to 4.86)	0.003	(0.48 to 1.29)	0.88	(0.45 to 1.72)	0.341	
Sex	Male	1.06	(0.65 to 1.73)	0.818	0.78	(0.44 to 1.39)	0.401	(0.68 to 1.81)	1.28	(0.69 to 2.38)	0.676	
Working	Not working	1.36	(0.76 to 2.45)	0.304	2.34	(1.10 to 4.97)	0.027	(0.44 to 1.42)	0.62	(0.29 to 1.34)	0.431	
Education	>Secondary School	1.25	(0.76 to 2.04)	0.377	1.19	(0.64 to 2.20)	0.583	(0.72 to 1.93)	1.37	(0.71 to 2.63)	0.515	
Health literacy	Adequate	2.25	(1.38 to 3.67)	0.001	2.25	(1.28 to 3.95)	0.005	(1.23 to 3.25)	1.13	(0.62 to 2.06)	0.005	
Duration of hypertension	<5 years	1.39	(0.85 to 2.28)	0.188	1.66	(0.95 to 2.90)	0.074	(0.59 to 1.60)	0.84	(0.47 to 1.50)	0.917	
BMI at intake	BMI ≥25 kg/m ²	1.05	(0.65 to 1.72)	0.834	0.98	(0.56 to 1.72)	0.955	(0.75 to 2.02)	1.27	(0.70 to 2.29)	0.403	
RAMP-Group or RAMP-individual	RAMP-individual	0.71	(0.44 to 1.14)	0.154	1.01	(0.55 to 1.87)	0.963	(4.74 to 14.29)	8.46	(4.68 to 15.28)	<0.001	
Good HBPM Knowledge	Full Mark	1.61	(0.98 to 2.65)	0.063	1.62	(0.89 to 2.95)	0.115		1.62	(0.89 to 2.95)	0.063	
Regular HBPM	>Weekly										0.12	

P<0.05 as statistically significant.

* Regular HBPM was defined as equal to or more than weekly HBPM.

† Good HBPM knowledge was defined as having full mark in knowledge score.

‡ OR adjusted of age, sex, work, education, health literacy, duration of hypertension, and having obesity or not.

§ P<0.05 are bold

aOR, adjusted OR; BMI, body mass index; HBPM, Home Blood Pressure Monitoring; RAMP, Risk Assessment and Management Programme.

DISCUSSION

After two different educational interventions: group education focusing on HBPM and individual counselling on HT self-management, more than 70% of the participants owned their HBPM devices. Two-thirds performed HBPM at least once weekly. The high acceptance of HBPM after interventions was probably related to clinicians' promotion and demonstrated of HBPM.²³ The RAMP-group scored significantly more full-mark in the HBPM knowledge checklist. Similar to previous studies, the finding supports group-intervention focused on HBPM knowledge and technique.^{8 19} The reason for better HBPM knowledge could be explained by interactive group education with information delivery, HBPM procedure demonstration and hands-on practice could improve and retain essential knowledge in HBPM. The professional-patient contact time was shortened in this study (total 3 hours) than previous successful intervention in USA (total 12 hours).¹⁸ Less contact time implied less healthcare cost, especially to the population with lower educational level or HL.

Promotion of regular HBPM practice is achievable by both interventions. The study findings further support interventions to increase in knowledge of treatment and complications of HT, the use of HBPM readings and the proper technique of HBPM.^{24 25}

There were multiple potential sources of HBPM measurement difficulties.¹⁴ Therefore, many patients, particularly those from lower socioeconomic classes, may experience more barriers to perform HBPM regularly and correctly.^{26 27} This study found patients were less likely to have regular HBPM if they were younger, working or with inadequate HL. In Ayala *et al* and Seidlerová *et al* have also found older age group and people with adequate HL were more likely to perform HBPM regularly.^{3 28} Interventions focusing on attitude, knowledge and technique of HBPM could further improve their HBPM knowledge compared with conventional counselling on overall HT self-management. The RAMP-group patients could share higher-quality HBPM reading to their clinicians, to design their daily HT management plan. The proven high knowledge in HBPM after focused education could encourage clinicians to trust patients' HBPM findings and promote HBPM in their practice.^{11 12}

This study's strength includes that it is a pragmatic result from a primary care cluster-randomised controlled trial. The response rate to follow-up 6 months postintervention was satisfactory. The HBPM knowledge score was developed according to international recommendations to patients. We have performed subgroup analysis which mainly looks into the difference in outcomes between patients with adequate and inadequate HL.

This study's limitation includes the practice of HBPM could only be verified by patients' reported habit with their consultation records. Although this method is widely accepted in a bigger population survey, there is a possibility of observer and recall bias.^{8 9 14} Nowadays, there were HBPM devices with electronically linked up with patients' mobile devices, or even healthcare professional's computer system so that the immediate HBPM records could be accurately achieved. In this study, the research subjects were mostly from a lower socioeconomic class, and they could not afford the more advanced HBPM models.

In conclusion, HL or educational level were not limiting factors to acquire HBPM knowledge. Patients could retain

Main messages

- ▶ Current research questions are: more patients who attended group education could retain better HBPM knowledge and regular HBPM than those attended conventional individual counselling 6 months post-interventions.
- ▶ Patients with better HL or higher educational level would have higher compliance with regular HBPM and better HBPM knowledge.
- ▶ In this study, more than 70% of participants continued HBPM 6 months after both types of interventions.
- ▶ Patients age more than 65, not working and with adequate health literacy were more likely to perform at least weekly HBPM.

What is already known on the subject

- ▶ Regular home blood pressure monitoring (HBPM) with additional education and feedback intervention was known to reduce blood pressure of patients with hypertension.
- ▶ High-quality HBPM could possibly be achieved in patients with low health literacy or educational level.

HBPM knowledge better possessed after focused group education than individual counselling on general HT management. Older, retired and patients with adequate HL were more likely to continue weekly HBPM 6 months after education. Healthcare professionals can focus continuous educational intervention for those younger, working-class and those with inadequate HL for regular HBPM.

Twitter Carlos K H Wong @CarlosWongHKU

Acknowledgements Ethical approval was obtained from the Research Ethics Committee of the Kowloon West Cluster, Hospital Authority of Hong Kong.

Contributors SNF, MCD, CKHW and BMYC designed the study. SNF and MCD collected the data. SNF, MCD and CKHW conducted the statistical analysis. SNF, MCD, CKHW and BMYC analyzed and interpreted the data. SNF, MCD, CKHW and BMYC wrote the manuscript. All authors critically reviewed and approved the final version of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This study was approved by the Hong Kong Hospital Authority, Kowloon West Cluster Clinical Research Ethics Committee, reference number: KW/EX-15-115(88-14).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. All data relevant to the study are included in the article or uploaded as online supplemental information. All data in this study are deidentified.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Sau Nga FU <http://orcid.org/0000-0002-6491-654X>

Man Chi Dao <http://orcid.org/0000-0002-8793-7167>

Carlos K H Wong <http://orcid.org/0000-0002-6895-6071>
Bernard M Y Cheung <http://orcid.org/0000-0001-9106-7363>

REFERENCES

- Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults. *J Am Coll Cardiol* 2018;71:e127–248.
- McCormack T, Krause T, O'Flynn N. Management of hypertension in adults in primary care: NICE guideline. *Br J Gen Pract* 2012;62:163–4.
- Ayala C, Tong X, Neeley E, et al. Home blood pressure monitoring among adults—American heart association cardiovascular health consumer survey, 2012. *J Clin Hypertens* 2017;19:584–91.
- Wang Y, Wang Y, Gu H, et al. Use of home blood pressure monitoring among hypertensive adults in primary care: Minhang community survey. *Blood Press Monit* 2014;19:140–4.
- Glynn LG, Murphy AW, Smith SM, et al. Self-Monitoring and other non-pharmacological interventions to improve the management of hypertension in primary care: a systematic review. *Br J Gen Pract* 2010;60:e476–88.
- Tucker KL, Sheppard JP, Stevens R, et al. Self-Monitoring of blood pressure in hypertension: a systematic review and individual patient data meta-analysis. *PLoS Med* 2017;14:1549–1277.
- Ringrose JS, Polley G, McLean D, et al. An assessment of the accuracy of home blood pressure monitors when used in device owners. *Am J Hypertens* 2017;30:683–9.
- Mondal S, Mondal H, Dutta R, et al. Competency in home blood pressure monitoring and effect of training program on competency. *Journal Of Clinical And Diagnostic Research* 2018;12:OC10–14.
- Sendra-Lillo J, Martínez-Martínez F, García-Corpas JP, et al. Validity of home blood pressure measurements manually registered by patients after an educational session provided by community pharmacists. *Blood Press Monit* 2015;20:232–6.
- Hwang KO, Thomas EJ, Petersen LA. Use of home blood pressure results for assessing the quality of care for hypertension. *JAMA* 2018;320:1753–4.
- Tirabassi J, Fang J, Ayala C. Attitudes of primary care providers and recommendations of home blood pressure monitoring—DocStyles, 2010. *J Clin Hypertens* 2013;15:224–9.
- Dugelay G, Kivits J, Desse L, et al. Implementation of home blood pressure monitoring among French GPs: a long and winding road. *PLoS One* 2019;14:e0220460.
- Fu SN, Dao MC, Wong CKH, et al. The association of health literacy with high-quality home blood pressure monitoring for hypertensive patients in outpatient settings. *Int J Hypertens* 2020;2020:7502468.
- Kallioinen N, Hill A, Horswill MS, et al. Sources of inaccuracy in the measurement of adult patients' resting blood pressure in clinical settings: a systematic review. *J Hypertens* 2017;35:421–41.
- SN F, Dao MC, Luk W. A cluster-randomized study on the risk assessment and management program for home blood pressure monitoring in an older population with inadequate health literacy. *J Clin Hypertens* 2020;22:1565–76.
- Leblanc M-E, Cloutier L, Veiga EV. Knowledge and practice outcomes after home blood pressure measurement education programs. *Blood Press Monit* 2011;16:265–9.
- Dawes MG, Kaczorowski J, Swanson G, et al. The effect of a patient education booklet and BP 'tracker' on knowledge about hypertension. A randomized controlled trial. *Fam Pract* 2010;27:472–8.
- Kim KB, Han H-R, Huh B, et al. The effect of a community-based self-help multimodal behavioral intervention in Korean American seniors with high blood pressure. *Am J Hypertens* 2014;27:1199–208.
- Dymek J, Skowron A, Polak W, et al. Assessment of knowledge and skills of patients with hypertension related to self-measurement of blood pressure (SBPM). *Arterial Hypertension* 2015;19:39–44.
- Shimbo D, Artinian NT, Basile JN, et al. Self-Measured blood pressure monitoring at home: a joint policy statement from the American heart association and American Medical association. *Circulation* 2020;142:e42–63.
- British Hypertension Society. Taking your blood pressure at home, 2020. Available: https://bihsoc.org/wp-content/uploads/2017/09/How_to_instructional_leaflet.pdf [Accessed 23 Sep 2020].
- Sharman JE, Howes F, Head GA, et al. How to measure home blood pressure: recommendations for healthcare professionals and patients. *Aust Fam Physician* 2016;45:31–4.
- Bancej CM, Campbell N, McKay DW, et al. Home blood pressure monitoring among Canadian adults with hypertension: results from the 2009 survey on living with chronic diseases in Canada. *Can J Cardiol* 2010;26:e152–7.
- Ostchega Y, Zhang G, Kit BK, et al. Factors associated with home blood pressure monitoring among US adults: National health and nutrition examination survey, 2011–2014. *Am J Hypertens* 2017;30:1126–32.
- Wake AD, Bekele DM, Tuiji TS. Knowledge and attitude of self-monitoring of blood pressure among adult hypertensive patients on follow-up at selected public hospitals in Arsi zone, Oromia regional state, Ethiopia: a cross-sectional study. *Integr Blood Press Control* 2020;13:1–13.
- Paldan K, Sauer H, Wagner N-F. *Promoting inequality? self-monitoring applications and the problem of social justice*. AI & society, 2018: 1–11.
- Paulsen MS, Andersen M, Munck AP, et al. Socio-Economic status influences blood pressure control despite equal access to care. *Fam Pract* 2012;29:503–10.
- Seidlerová J, Filipovský J, Wohlfahrt P, et al. Availability and use of home blood pressure measurement in the Czech general population. *Cor Vasa* 2014;56:e158–63.

Table S: Patients outcomes at 6-months post-interventions

Outcomes	Complete Data	Missing data	Percentage	Sex*	Age**
				Pearson Chi's Square P-value***	
Any change of HT Drugs	289	0	0.0%	NA	NA
Change of body weight (kg, mean \pm S.D.)	276	13	4.5%	0.265	0.008
Fasting Blood Sugar (mmol/l, mean \pm S.D.)	285	4	1.4%	0.138	0.729
Low Density Lipoprotein (mmol/l, mean \pm S.D.)	283	6	2.1%	0.162	0.670
Have their own BP devices (%)	289	0	0.0%	0.086	0.003
Measure BP at home in past 6 months (%)	289	0	0.0%	0.086	0.003
HBPM Assessment Score (%)	272	17	5.9%	0.372	0.161

BP = Blood Pressure; HT = hypertension; HBPM = Home Blood Pressure Monitoring; S.D. = standard deviation

* Sex = male: female; **Age = <65 : \geq 65

***P < 0.05 = significantly more missing data in the group of \geq 65