OPEN ACCESS

In situ simulation training for a better interprofessional team performance in transferring critically ill patients with COVID-19: a prospective randomised control trial

Sidharta Kusuma Manggala ,¹ Aida Rosita Tantri ,^{1,2} Adhrie Sugiarto ,¹ Imelda Rosalyn Sianipar ,^{2,3} Theddeus Octavianus Hari Prasetyono ,^{4,5,6}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/postgradmedj-2021-141426>).

For numbered affiliations see end of article.

Correspondence to

Aida Rosita Tantri,
Anesthesiology and Intensive
Care, Universitas Indonesia,
Cipto Mangunkusumo National
General Hospital, Jakarta,
Indonesia, Indonesia;
aidatantri@gmail.com

Received 7 December 2021
Accepted 15 January 2022

ABSTRACT

Background Transferring critically ill patients with COVID-19 is a challenging task; therefore, well-trained medical team is needed. This study aimed to determine the role of in situ simulation training during pandemic by using high-fidelity manikin to improve interprofessional communication, skills and teamwork in transferring critically ill patients with COVID-19.

Methods This single-blinded randomised control trial included 40 subjects allocated into standard low-fidelity simulator (LFS) and high-fidelity simulator (HFS) groups. Subjects, who were not members of multiprofessional team taking care of patients with COVID-19, in each group were assigned into small groups and joined an online interactive lecture session, two sessions of in-situ simulation and a debriefing session with strict health protocols. The first simulation aimed to teach participants the skills and steps needed. The second simulation aimed to assess transfer skills, communication and teamwork performance, that participants had learnt using a validated, comprehensive assessment tool. Data were analysed using unpaired t test or Mann-Whitney test.

Results The HFS group showed significantly better overall transfer and communication skills than LFS group (89.70 ± 4.65 vs 77.19 ± 3.6 , <0.05 and 100 vs 88.34 ($63.33-100$), $p=0.022$, respectively). The HFS group also demonstrated significantly better teamwork performance than the standard LFS group (90 ($80-900$) vs 80 ($70-90$), $p=0.028$).

Conclusion In situ simulation training using HFS significantly showed better performance than the standard training using LFS in regards to overall transfer and communication skills as well as teamwork performance. The training using HFS may provide a valuable adjunct to improve interprofessional skills, communication and teamwork performance in transferring critically ill patients with COVID-19. Trial registration number NCT05113823.

INTRODUCTION

Safe transportation of critically ill patients is challenging. It depends on patient selection, staff training, interprofessional teamwork, predefined hospital transport protocols and checklists, appropriate transport equipment availability and also transport timing.¹⁻⁵ Such complex system and

situation could be recreated in a simulated environment. Simulation-based learning has proven to have a good impact on the students' competence. An in situ simulation, conducted with multidisciplinary team, could recreate the circumstances similar to which they conduct their usual activities, thus further accelerate the students' knowledge, skills and safety attainment process.⁶ High-fidelity simulator (HFS) has also started to be used widely in simulation-based training aside from low-fidelity simulator (LFS). It supports a realistic yet safe environment, where participants can face a rare clinical situation and learn from mistakes without harming the patients.^{3,5}

Simulation may play an integral role in the medical education's response to the pandemic era. COVID-19 pandemic created an urgent need for targeted and adaptive training for all intensive care and medical emergency team members, especially in conducting safe intrahospital transportation of critically ill patients with COVID-19. Modifying interprofessional simulation-based training by implementing strict health protocols is necessary to initiate simulation-based training. This implementation enable us to continue improving knowledge, skills, communication and teamwork performance in managing and transferring patients with COVID-19 during pandemic era. This study aimed to determine the role of in situ simulation training by using HFS compared with the standard LFS to improve interprofessional skills, communication and teamwork performance in transferring critically ill patients with COVID-19.

METHODS

This single-blinded randomised control trial aims to study in situ simulation training using HFS compared with a standard use of LFS. The study was held in the High Care Unit Cipto Mangunkusumo Hospital and Simulation-Based Medical Education and Research Center (SIMUBEAR), IMERI Universitas Indonesia. The sample size was determined using a numeric analytical formula to achieve a 80% power with 5% error rate and an estimated drop out of 10%, resulting in 20 subjects in each group. After obtaining ethical approval from the Ethics Committee of the Faculty of Medicine, Universitas Indonesia/Cipto Mangunkusumo Hospital, achieving Clinical Trial approval, 40 subjects,

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Manggala SK, Tantri AR, Sugiarto A, et al. *Postgrad Med J* Epub ahead of print: [please include Day Month Year]. doi:10.1136/postgradmedj-2021-141426

comprised of 16 medical doctors and 24 nurses, who were in good physical condition, not detected positive from COVID-19 infection and had no history of involving COVID-19 patient care, were voluntarily recruited. All subjects provided informed consent. The study fully implemented the principles outlined in the Declaration of Helsinki and followed the CONSORT checklist (online supplemental file 1 online supplemental figure 1)

The simulation-based training consisted of an online interactive lecture session 1 day before simulation, two sessions of in situ simulation and debriefing at the end of each simulation session. Eight scenarios (online supplemental figure 2) were developed to prevent information bias, as the study was spread throughout 3 days. We divided eight groups into 3 days as there was a time limit on the venue. Reading materials of the interactive lecture topic were given to the subjects a day before. Subjects were obliged to complete an online 5-item multiple-choice pretest and post-test for cognitive evaluation.

Specific assessment tool was developed based on focus discussion group, after that it was developed and validated to evaluate interprofessional teamwork, communication and the team's ability to perform each step in the COVID-19 critically ill patient transportation checklist were used. Skills and communication assessment tools were developed from predefined hospital transport protocols and checklists. Interprofessional teamwork was assessed using interprofessional teamwork assessment tools that had been developed in the previous study.^{7–11} Each item was rated '0' for undone skill, '2' for a incompletely done skill and '5' for the skill that had been done completely. Validation of the study checklist was performed by measuring Cronbach's alpha by our senior anaesthesiologist consultants (online supplemental figure 3). The panel consisted of five senior anaesthesiologist consultants who were not part of the research team.

Subjects were randomly allocated into two large groups, the HFS and the LFS groups using a random allocator downloaded from www.randomizer.org. Each group was divided into smaller groups, consisting of two doctors and three nurses. All subjects joined the same interactive lecture session discussing patient transfer methods during COVID-19 and two sessions of in situ simulation according to their assigned group. Each group was given a 1-hour lecture session and an approximately 2-hour in situ simulation session. In situ simulation and debriefing were performed by implementing appropriate personal protective equipment (PPE) and social distancing. At the end of each simulation session, each group underwent a debriefing session conducted by an experienced instructor.

The first simulation was aimed to teach participants skills and steps in transporting critically ill patients with COVID-19 according to the hospital checklist. Meanwhile, the second simulation was aimed to assess skills, teamwork and communication that participants had learnt from the previous simulation by using the abovementioned assessment tools. In addition to comparing each point in the assessment tool between the two groups, the points earned were added to get the overall points for total skills, cooperation and communication score. At the end of the session, subjects gave feedback immediately online, which provided accountability for attendance, content learning and course evaluation in general.

All collected data were then analysed using SPSS V.26.0. Categorical data are presented in the form of numbers and percentages (n (%)). Numerical data are shown in the form of mean $\pm$ SD if the data distribution is normal or in the form of the median (minimum–maximum value) if the distribution is not normal. Unpaired t test and Mann-Whitney test were used to analyse the two numerical variables.

Table 1 Subject characteristics

Variables	HFS group (n=20)	LFS group (n=20)
Age (year)	27.5 (23–41)	30 (24–46)
Sex		
Male (n)	4	7
Female (n)	16	13
Working experience (Year)	3.5 (1–13)	8 (0.25–27)
Working unit		
Emergency doctor (n)	8	5
Inpatient unit (n)	4	4
Outpatient unit (n)	4	4
HCU and ICU (n)	4	4

*Value in percentage (%).

†Value in median (minimum–maximum).

HFS, high-fidelity simulator; ICU, Intensive Care Unit; LFS, low-fidelity simulator.

RESULTS

All the participants completed the study. The majority were woman (80% in the HFS group and 65% in the LFS group), and an average of 27.5 years and 30 years old in the HFS and LFS groups, respectively. Participants in the HFS group had median working experience of 3.5 years, and those in the LFS group had 8 years. The participant characteristic details are summarised in [table 1](#).

There was no significant difference in pretest and post-test cognitive scores between the two groups ([table 2](#)). The [table 2](#) also presents the overall transfer skill, communication skill and teamwork scores. The subsection analysis of the skill components is presented in [table 3](#).

DISCUSSION

Intrahospital transfer of critically ill patients with COVID-19 is challenging and requires proper strategies to maintain patient safety and prevent disease exposure to the medical team. Sixty-eight per cent of critically ill patients could experience unexpected events; and 9% could experience severe unexpected events, including hypotension, airway problems and increased intracranial pressure during transport.¹² In handling critically ill patients with COVID-19, medical teams require skills, communication, good teamwork and sufficient knowledge.

Since the pandemic started, there have been increasing needs in intrahospital transfer of critically ill patients with COVID-19. Two studies reported that 63.4% of transferred patients were confirmed for COVID-19.^{13 14} Cautious planning and prudent decision are needed to ensure the safety of the critically

Table 2 Comparison of cognitive, transfer skill, communication skill and teamwork scores between the groups

Variable	HFS group (n=4)	LFS group (n=4)	P
Cognitive			
Pre-test	72.99 $\pm$ 11.13	68.99 $\pm$ 14.55	0.335*
Post-test	87.67 $\pm$ 7.26	86.99 $\pm$ 9.79	0.889†
Transfer skills	89.70 $\pm$ 4.65	77.19 $\pm$ 3.61	0.000†
Communication	100	88.34 (63.33–100)	0.022†
Team work	90 (80–90)	80 (70–90)	0.028†

*Unpaired t-test; .

†Mann-Whitney test.

.HFS, high-fidelity simulation; LFS, low-fidelity simulation.

Table 3 Distribution of skill components of the two groups

Variable	HFS group (n=4)	LFS group (n=4)	P
Pre-transfer preparation skills			
Equipment preparation and donning	86.54±13.92	77.95±12.49	0.287*
Patient preparation	87.41±7.12	77.78±7.70	0.048*
Medical team preparation	100 (50–100)	100	1.000†
Skills during patient transfer	93.2 (83–98)	85 (83–97)	0.090*
Monitoring of patient and equipment	90 (73.33–90)	76.67 (56.67–90)	0.067*
PPE doffing	85.83±8.01	48.33±23.80	0.010†

*Mann-Whitney test.

†unpaired t-test.

HFS, high-fidelity simulation; LFS, low-fidelity simulation; PPE, personal protective equipment.

ill COVID-19 transferring process. Hence, there are urgent needs for targeted and adaptive training for all intensive care and medical emergency team members conducting intrahospital transportation of critically ill patients with COVID-19.

Doubts about the risk of COVID-19 infection have made many institutions withhold in-hospital training. Modifying interprofessional simulation-based training by implementing strict social distancing and appropriate PPE might be the solution to improve the knowledge, skills, communication and teamwork needed to manage and transfer patients with COVID-19. Additionally, simulation-based training using HFS might allow learners of all levels an opportunity to immerse themselves in a better realism-simulated clinical scenario, suspend their disbelief and engage more in the learning activity. Hence, this highly needed simulation base training can be carried out more effectively.

There were no differences in the baseline age and workplace characteristics between the HFS and LFS groups (table 1). Subjects in the LFS group had longer working experience compared with those in the HFS group. The outcome of the post-test supports the homogeneity of the participants between the groups. After the team training, there was an increase in participants' knowledge in both groups, but there was no significant difference found between both groups. The team-based simulation assessment revealed significant difference in the global skill performance scores ($p>0.05$) between the HFS (89.70 ± 4.5) and LFS (77.19 ± 3.61) groups. There were also significant differences in the global communication and teamwork performance scores ($p=0.022$ and $p=0.028$, respectively) in both groups. Adequate communication and good teamwork are critical to the safe transfer of a critically ill patient. The results are in line with the previous researches that HFS is preferable for skills training, stress exposure training and team training than LFS.^{7–11} Training using HFS enhances participants' skills, teamwork and leadership.⁹ Obviously, HFS contains features such as realistic physiological responses, the ability to communicate and interact with the manikin and various other feedback mechanisms. Hence, HFS enables low-risk, standardised training with a complex, immersive scenario and realistic feedback.^{15–18}

The subject characteristic may also play an essential role in this study. Most of the subjects participating in this training had work experience for 4–7 years. Based on the simulation training given, the participants achieve a different competency levels. A high degree of realism simulation training favours a higher clinical competence in the classical Miller pyramid of clinical competence assessment.

Both HFS and LFS simulations were performed as in situ simulations. The attendee had opportunities to identify hazards and deficiencies in their clinical systems, the environment and know another provider. Patients with COVID-19 are often highly complex Intensive Care Unit (ICU) patients with more than one organ system failure. Pretransfer preparation is needed to ensure patient's physiological stability than patient safety during transfer. Unstable patients and lack of potential events' anticipation during transfer can worsen patient outcomes. Physiological stability during transfer involves careful pretransfer assessment and care and preparation.¹⁹ Troncoso *et al* stated that approximately 25.4% of critically ill patients with COVID-19 need vasopressor, 13.1% were pharmacologically paralysed and 22% required change of ventilator setting during transfer.²⁰ Allen *et al* stated that 45% of patients with COVID-19 needed to be intubated, 40% required additional paralytic drugs and 40% of the patients required to be given vasopressors before transfer.¹³ Most patients require oxygen therapy ranging from high-flow nasal cannula, continuous positive airway pressure (CPAP), to mechanical ventilation. The success of transfer critically ill patients with COVID-19 is based on anticipation and prevention of potential complications and hazards to the patient and transfer team.¹⁶

When inspecting the individual section of the assessment tool, the statistically significant difference between HFS and LFS was noted for item patient preparation but not for equipment preparation, team preparation and donning items (table 3). Preparing patients before a transfer is a more complex procedure than equipment preparation, team preparation and donning. This procedure also requires good clinical reasoning, judgement and communication between team members. HFS was better than LFS in this case. A higher level of fidelity was needed in achieving higher intended learning goals.²¹ In the immersive and safe learning environment of HFS, participants can make mistakes, revise those mistakes in real time and learn from them without fear of compromising patient safety. Previous research has also shown that the effect of stress that participants gain from simulation-based high-fidelity training will be appraised as a challenge rather than a threat. This positive reaction will contribute to performance improvement.^{22 23}

Interestingly, both groups do not show significant difference in medical team preparation items, including donning items to ensure healthcare provider safety. The participants might have been accustomed to preparing themselves in this pandemic state, including wearing PPE and maintaining a good protocol.²⁴ Donning items included wearing appropriate masks and other PPE (gloves, fluid-repellent long-sleeved gown and eye protection devices), which were put in order based on the level of precautions required.

There was no significant difference in skill score during the transfer between both groups (table 2). This might be because the task to be fulfilled during transfer was not as difficult and complex as pretransfer preparation. If appropriate and precise measures have been taken pretransfer, there should be little requirement for active intervention during transport. During the transfer, medical teams should continue to reassess the patient's clinical status, ensuring that vascular access sites remain accessible, good teamwork and communication.

After transfer, the HFS group showed a higher result in PPE doffing (table 3). Doffing involves more complex and challenging steps than donning.²⁵ As mentioned before, HFS is better than LFS in achieving competence in a more complex and difficult task. In line with this result, deviation in doffing protocols was more often found than in donning protocols. Ensuring safe

Original research

practices for high-risk, highly potential COVID-19 exposure scenarios in minimising contamination risks requires specific training methods in doffing protocols compliance.

Limitation

Although this study tested the participants' team performances and attitudes using simulation-based assessments and validated tools, the self-reported attitude towards teamwork and collaboration of the attendee and the retention of team performances were not measured. Further studies can be conducted to gain insights into the more effective training methods and the use of the attendee's self-reported attitude in transferring critically ill patients with COVID-19.

CONCLUSION

In situ simulation training improved skills, communication and interprofessional team cooperation in transferring critically ill patients with COVID-19. HFS-based training showed better team performance compared with the standard LFS-based training.

Main messages

- ▶ In situ simulation training, both using high-fidelity simulator (HFS) and low-fidelity simulator (LFS), improved skills in transferring critically ill patients with COVID-19.
- ▶ In situ simulation training, both using HFS and LFS, improved communication in transferring critically ill patients with COVID-19.
- ▶ In situ simulation training, both using HFS and LFS, improved interprofessional team cooperation in transferring critically ill patients with COVID-19.

Current research questions

- ▶ Will in situ simulation training improve skills of its participants?
- ▶ Will in situ simulation training improve communication of its participants?
- ▶ Will in situ simulation training improve interprofessional team cooperation of its participants?

What is already known on the subject

- ▶ Subjects' Characteristics
- ▶ Subjects are health professionals that are usually handle hospital patient transfer

Author affiliations

¹Department of Anesthesiology and Intensive Care, Faculty of Medicine, Universitas Indonesia, Cipto Mangunkusumo National General Hospital, Jakarta, DKI Jakarta, Indonesia

²SIMUBEAR (Simulation Based Medical Education and Research Center), IMERI (Indonesian Medical Education and Research Institute), Faculty of Medicine, Universitas Indonesia, Jakarta, DKI Jakarta, Indonesia

³Department of Medical Physiology, Faculty of Medicine, Universitas Indonesia, Cipto Mangunkusumo National General Hospital, Jakarta, DKI Jakarta, Indonesia

⁴Department of Plastic Surgery, Departement of Surgery, Faculty of Medicine, Universitas Indonesia, Cipto Mangunkusumo National General Hospital, Jakarta, DKI Jakarta, Indonesia

⁵ICTEC (Indonesian Clinical Training and Education Center), Faculty of Medicine, Universitas Indonesia, Cipto Mangunkusumo National General Hospital, Jakarta, DKI Jakarta, Indonesia

⁶Medical Technology Cluste IMERI (Indonesian Medical Education and Research Institute), Faculty of Medicine, IMERI (Indonesian Medical Education and Research Institute), Faculty of Medicine, Universitas Indonesia, Jakarta, DKI Jakarta, Indonesia

Acknowledgements This study was a part of PUTI 2020 Grant from Universitas Indonesia (HIBAH PUTI UI Q3 2020: NKB-1879/UN2/RST/HKP.05/00/2020).

Contributors SKM and ART planned the study and conceived the study. AS conducted study survey. IRS and TOHP processed and analysed study data. TOHP supervised the study. SKM developed the theoretical framework. SKM, ART and AS wrote the manuscript. All authors discussed the results and commented on the manuscript. ART submitted the study. ART and TOHP revised the manuscript. SKM and ART are the guarantors of the study.

Funding This works was supported by the Universitas Indonesia Research Fund (HIBAH PUTI UI Q3 2020: NKB-1879/UN2/RST/HKP.05/00/2020)

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s)

Ethics approval This study involves human participants and was approved by Ethics Committee of Universitas Indonesia (Approval Number KET-591/UN2.F1/ETIK/PPM.00.02/2020). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. Not applicable.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Sidharta Kusuma Manggala <http://orcid.org/0000-0002-0974-5864>

Aida Rosita Tantri <http://orcid.org/0000-0002-5535-985X>

Adhrie Sugiarto <http://orcid.org/0000-0003-2542-1373>

Imelda Rosalyn Sianipar <http://orcid.org/0000-0002-1438-096X>

Theddeus Octavianus Hari Prasetyono <http://orcid.org/0000-0003-3675-4309>

REFERENCES

- 1 Eiding H, Kongsgaard UE, Braarud A-C. Interhospital transport of critically ill patients: experiences and challenges, a qualitative study. *Scand J Trauma Resusc Emerg Med* 2019;27:27.
- 2 Fanara B, Manzoni C, Barbot O, et al. Recommendations for the intra-hospital transport of critically ill patients. *Crit Care* 2010;14:R87.
- 3 Lahner D, Nikolic A, Marhofer P, et al. Incidence of complications in intrahospital transport of critically ill patients--experience in an Austrian university hospital. *Wien Klin Wochenschr* 2007;119:412-6.
- 4 Caruana M, Culp K. Intrahospital transport of the critically ill adult: a research review and implications. *Dimens Crit Care Nurs* 1998;17:146-56.
- 5 Baptista R, Pereira F, Martins J. Perception of nursing students on high-fidelity practices: a phenomenological study. *J Nurs Educ Pract* 2016;6:10-22.
- 6 Goldshtein D, Krensky C, Doshi S, et al. In situ simulation and its effects on patient outcomes: a systematic review. *BMJ Simulation and Technology Enhanced Learning* 2020;6:3-9.
- 7 Chang Y-C, Chou L-T, Lin H-L, et al. An interprofessional training program for intrahospital transport of critically ill patients: model Build-up and assessment. *J Interprof Care* 2019;00:1-5.
- 8 Jarden RJ, Quirke S. Improving safety and documentation in intrahospital transport: development of an intrahospital transport tool for critically ill patients. *Intensive Crit Care Nurs* 2010;26:101-7.
- 9 Intensive Care Society. *Guidance on: the transfer of the critically ill adult*, 2019: 1-40.
- 10 Williams P, Karupiah S, Greentree K, et al. A checklist for intrahospital transport of critically ill patients improves compliance with transportation safety guidelines. *Aust Crit Care* 2020;33:20-4.
- 11 da SR, Amante LN. Checklist for the intrahospital transport of patients admitted to the intensive care unit. *Enferm* 2015;24:539-47.
- 12 Goldshtein D, Krensky C, Doshi S, et al. In situ simulation and its effects on patient outcomes: a systematic review. *Bmj Stel* 2020;6:3-9.
- 13 Allen R, Wanersdorfer K, Zebley J, et al. Interhospital transfer of critically ill patients because of coronavirus disease 19-Related respiratory failure. *Air Med J* 2020;39:498-501.
- 14 Garfinkel E, Lopez S, Troncoso R, et al. A critical care transport program's innovative approach to safety during the coronavirus disease 2019 pandemic. *Air Med J* 2021;40:pp.112-4.

- 15 Lim WY, Ong J, Vimal V, *et al.* High-Fidelity simulation training with PPE may optimise resuscitation outcomes in the COVID-19 era. *Resuscitation* 2021;159:42–4.
- 16 Martin T. Transporting the adult critically ill patient. *Surgery* 2012;30:219–24.
- 17 Finan E, Bismilla Z, Whyte HE, *et al.* High-Fidelity simulator technology may not be superior to traditional low-fidelity equipment for neonatal resuscitation training. *J Perinatol* 2012;32:287–92.
- 18 Massoth C, Röder H, Ohlenburg H, *et al.* High-Fidelity is not superior to low-fidelity simulation but leads to overconfidence in medical students. *BMC Med Educ* 2019;19:29.
- 19 Gray A, Bush S, Whiteley S. Secondary transport of the critically ill and injured adult. *Emerg Med J* 2004;21:281–5.
- 20 Troncoso RD, Garfinkel EM, Leon D, *et al.* Decision making and interventions during Interfacility transport of High-Acuity patients with severe acute respiratory syndrome coronavirus 2 infection. *Air Med J* 2021;40:220–4.
- 21 Munshi F, Lababidi H, Alyousef S. Low- versus high-fidelity simulations in teaching and assessing clinical skills. *Journal of Taibah University Medical Sciences* 2015;10:12–15.
- 22 Tomaka J, Blascovich J, Kelsey RM, *et al.* Subjective, physiological, and behavioral effects of threat and challenge appraisal. *J Pers Soc Psychol* 1993;65:248–60.
- 23 Nicolaides M, Theodorou E, Emin EI, *et al.* Team performance training for medical students: low vs high fidelity simulation. *Ann Med Surg* 2020;55:308–15.
- 24 Tabah A, Ramanan M, Laupland KB, *et al.* Personal protective equipment and intensive care unit healthcare worker safety in the COVID-19 era (PPE-SAFE): an international survey. *J Crit Care* 2020;59:70–5.
- 25 Kwon JH, Burnham C-AD, Reske KA, *et al.* Assessment of healthcare worker protocol deviations and Self-Contamination during personal protective equipment Donning and Doffing. *Infect Control Hosp Epidemiol* 2017;38:1077–83.