

Table

<b>Inflammatory and Autoimmune markers</b>	
ANCA	Negative
ANA-Immunofluorescence	< 80
Anti-ds-DNA	< 25 (0-25 IU/mL)
Cryoglobulin	Negative
<b>Complement 3 (C3)</b>	<b>1.74 elevated than reference range (0.80-1.60 g/L)</b>
Complement 4 (C4)	0.41 (0.17-0.60 g/L)
Anti-HBc, Total	Non-Reactive
HBsAg	Non-Reactive
Anti-HCV	Non-Reactive
<b>Erythrocyte Sedimentation Rate (ESR)</b>	<b>48 elevated (3-15mm/hr)</b>
<b>C-Reactive Protein (CRP)</b>	<b>10.7 elevated (0-5mg/L)</b>
<b>Full Blood Count</b>	
White blood cell	<b>10.2 high (4-9.6 X10<sup>9</sup>/L)</b>
Haemoglobin	12.2 (11.8-14.6 g/dL)
Platelets	338 (150-360 X10 <sup>9</sup> /L)
Absolute Neutrophils	<b>7.42 elevated (1.90-6.6X10<sup>9</sup>/L)</b>
Absolute Lymphocytes	2.04 (1.10-3.10X10 <sup>9</sup> /L)
Absolute Monocytes	0.56 (0.20-0.70X10 <sup>9</sup> /L)
Absolute Eosinophils	0.05 (0.00-0.60X10 <sup>9</sup> /L)
Absolute Basophils	0.08 (0.00-0.10X10 <sup>9</sup> /L)
<b>Coagulation panel</b>	
APTT	31.4 s
PT	13.9 s
INR	1.1
<b>Urine Protein and creatinine ratio</b>	
Total Protein on admission	0.3 U
Creatinine admission	18.1 U
Prot:Crea Ratio on admission	14
<b>Urine Microscopy</b>	
WBC,Urine on admission	> 225 (0-6 cells/uL)
RBC,Urine on admission	18 (0-13 cells/uL)
<b>Histopathology</b>	
Punch biopsy of skin to the subcutis. The stratum corneum is basketweave. The epidermis is unremarkable. There is a superficial and deep perivascular infiltrate of lymphocytes with many eutrophils. There is nuclear dust and red cell extravasation. Fibrin exudation is seen around some upper dermal blood vessels. This is in keeping with the diagnosis of leukocytoclastic vasculitis.	
<b>Direct Immunofluorescence</b>	
IgM	Negative
IgG	Negative
C3	Negative
Fibrinogen	Negative

**Table 1:** Summary of patient's investigations.